

Transitivity of Heisenberg group extensions of hyperbolic systems

Ian Melbourne
Department of Mathematics
University of Surrey
Guildford, Surrey GU2 7XH, UK

Viorel Nițică*
Department of Mathematics
West Chester University
West Chester, PA 19383, USA

Andrei Török*
Department of Mathematics
University of Houston
Houston, TX 77204-3008, USA

May 12, 2010

Abstract

We show that among C^r extensions ($r > 0$) of a uniformly hyperbolic dynamical system with fiber the standard real Heisenberg group \mathcal{H}_n of dimension $2n + 1$, those that avoid an obvious obstruction to topological transitivity are generically topological transitive. Moreover, if one considers extensions with

*and Institute of Mathematics of the Romanian Academy, P.O. Box 1-764, RO-70700 Bucharest, Romania

fiber a connected nilpotent Lie group with compact commutator subgroup (for example, \mathcal{H}_n/\mathbb{Z}), among those that avoid the obvious obstruction, topological transitivity is open and dense.

1 Introduction

This paper is part of a sequence aiming to study topological transitivity in various classes of noncompact group-extensions of hyperbolic systems. Consider a continuous transformation $f : X \rightarrow X$, a Lie group Γ , and a continuous map $\beta : X \rightarrow \Gamma$ called a *cocycle*. These determine a *skew product*, or Γ -*extension*,

$$f_\beta : X \times \Gamma \rightarrow X \times \Gamma, \quad f_\beta(x, \gamma) = (fx, \gamma\beta(x)).$$

The Γ -extension f_β is called *topologically transitive*, or simply *transitive*, if it has a dense orbit. The problem of interest to us is whether noncompact Lie group extensions of a hyperbolic basic set are typically topologically transitive.

Let (M, d_M) be a smooth manifold endowed with a Riemannian metric. Let $f : M \rightarrow M$ be a smooth diffeomorphism and $X \subset M$ a compact and f -invariant subset of M . Let Df be the derivative of f . We recall that X is said to be *hyperbolic* if there exists a continuous Df -invariant splitting $E^s \oplus E^u$ of the tangent bundle $T_X M$ and constants $C_1 > 0, 0 < \lambda < 1$, such that for all $n \geq 0$ and $x \in X$ we have:

$$\begin{aligned} \|(Df^n)_x v\| &\leq C_1 \lambda^n \|v\|, \quad v \in E_x^s \\ \|(Df^{-n})_x v\| &\leq C_1 \lambda^n \|v\|, \quad v \in E_x^u. \end{aligned}$$

We say that X is *locally maximal* if there exists an open neighborhood U of X such that every compact f -invariant set of U is contained in X . A locally maximal hyperbolic set X is a *hyperbolic basic set* if $f : X \rightarrow X$ is transitive and X does not consist of a single periodic orbit.

Given a connected Lie group Γ and a C^r cocycle $\beta : X \rightarrow \Gamma, r > 0$, we consider the Γ -extension $f_\beta : X \times \Gamma \rightarrow X \times \Gamma$ given by $f_\beta(x, \gamma) = (fx, \gamma\beta(x))$. For brevity, we say that the cocycle β is *transitive* if the Γ -extension f_β is transitive. In [6] we proposed a general conjecture about transitivity: namely that modulo obstructions appearing from the fact that the range of the cocycle is included in a maximal semigroup with non-empty interior, the set of C^r transitive cocycles contains an open and dense subset. The conjecture is proved for various classes of Lie groups, mostly semidirect products of compact and Euclidean, in [2, 5, 6, 7, 10]. An important test case is

presented by the special Euclidean group $\Gamma = SE(n) = SO(n) \times \mathbb{R}^n$. It is shown in [5, 6, 7] that when n is even the set of cocycles that are transitive is open and dense.

In [8] we showed that for $SE(n)$, $n \geq 3$ odd, the transitive C^r cocycles form a residual subset of the space of all C^r cocycles for all $r > 0$. In other words, transitivity is generic for such extensions. More generally, we considered *Euclidean-type* groups of the form $\Gamma = G \times \mathbb{R}^n$ where G is a compact connected Lie group. The general case of the conjecture remains unsolved for $SE(n)$ if $n \geq 3$ odd.

Recently, [9] obtained examples of groups that are compact extensions of nilpotent (not abelian) Lie groups for which transitivity is open and dense. Recall that a *compact element* g in a Lie group is one for which the closure of the cyclic group generated by g is compact. The method used in [9] borrows from [6] as it relies on the existence of an open and dense set of compact elements in Γ . This approach cannot be applied to the case of a nilpotent Lie group due to the lack of compact elements.

In this paper we study the conjecture for certain 2-step nilpotent Lie groups.

Definition 1.1 For $n \geq 1$, let \mathcal{H}_n denote the Heisenberg group, consisting of matrices of the form

$$(a, b, c) := \begin{pmatrix} 1 & a^T & c \\ 0 & I_n & b \\ 0 & 0 & 1 \end{pmatrix} \in \text{Mat}_{n+2}(\mathbb{R}).$$

where $a, b \in \mathbb{R}^n, c \in \mathbb{R}$ and I_n is the n -dimensional identity matrix.

Remark 1.2 (a) We can identify \mathcal{H}_n with the space $\mathbb{R}^n \oplus \mathbb{R}^n \oplus \mathbb{R}$ endowed with the multiplication

$$(a, b, c)(a', b', c') = (a + a', b + b', c + c' + a^T b'). \quad (1.1)$$

(b) \mathcal{H}_1 is the standard 3-dimensional Heisenberg group.

The center of \mathcal{H}_n is $[\mathcal{H}_n, \mathcal{H}_n] = \{(0, 0, c)\} = \mathbb{R}$. Denote $\widehat{\mathcal{H}}_n = \mathcal{H}_n/\mathbb{R} \cong \mathbb{R}^{2n}$. If $\beta : X \rightarrow \mathcal{H}_n$ is a cocycle, denote by $\widehat{\beta} : X \rightarrow \mathbb{R}^{2n}$ the corresponding quotient cocycle. There is an obvious obstruction to transitivity, namely that $\widehat{\beta} : X \rightarrow \mathbb{R}^{2n}$ takes values in a half-space bounded by a hyperplane passing through the origin. To avoid repetition, we assume from now on that a *half-space* in a linear space is always bounded by a hyperplane passing through the origin. More generally, if $\widehat{\beta}$ is cohomologous to a cocycle with values in a half-space, then f_β is not transitive.¹

¹ We recall that $\beta, \beta' : X \rightarrow \mathbb{R}^d$ are *cohomologous* if there exists a map $P : X \rightarrow \mathbb{R}^d$ such that for all $x \in X$, $\beta'(x) = P(fx) + \beta(x) - P(x)$.

Remark 1.3 By [1], a Hölder \mathbb{R}^d -valued cocycle is cohomologous to one that takes values in a half-space if and only if its periodic data is in a half-space.

If $r > 0$, let $\mathcal{S}^r(X, \mathcal{H}_n)$ be the set of C^r cocycles $\beta : X \rightarrow \mathcal{H}_n$ for which $\widehat{\beta}$ is *not* cohomologous to a cocycle with values in a half-space. Our first main result is

Theorem 1.4 *Assume that X is a hyperbolic basic set for $f : X \rightarrow X$. Let $n \geq 1$, $r > 0$. Among cocycles $\beta \in \mathcal{S}^r(X, \mathcal{H}_n)$, those that are transitive are generic.*

There is a class of nilpotent groups for which we obtain stronger results, and for which the proofs are much simpler.

Let Γ be a nilpotent Lie group, with compact commutator subgroup $[\Gamma, \Gamma]$. Note that, by [13, Lemma 4], in this case $[\Gamma, \Gamma]$ is central, hence Γ is two-step nilpotent. For a cocycle $\beta : X \rightarrow \Gamma$, denote its image in the abelianization of Γ by $\widehat{\beta} : X \rightarrow \Gamma/[\Gamma, \Gamma]$. Since $\Gamma/[\Gamma, \Gamma]$ is an abelian Lie group, it is isomorphic to $\mathbb{R}^a \times \mathbb{T}^b$. Denote by $\widetilde{\beta} : X \rightarrow \mathbb{R}^a$ the image of $\widehat{\beta}$ in \mathbb{R}^a .

For $r > 0$, let $\mathcal{S}^r(X, \Gamma)$ be the set of C^r cocycles $\beta : X \rightarrow \Gamma$ for which $\widetilde{\beta}$ is *not* cohomologous to a cocycle with values in a half-space.

Theorem 1.5 *Let Γ be a nilpotent Lie group with $[\Gamma, \Gamma]$ compact, and $r > 0$. Assume that X is a hyperbolic basic set for $f : X \rightarrow X$. Among cocycles $\beta \in \mathcal{S}^r(X, \Gamma)$, those that are transitive contain an open and dense set.*

One group Γ for which the above Theorem applies is the quotient \mathcal{H}_n/\mathbb{Z} , where the central subgroup \mathbb{Z} is generated by $(0, 0, 1)$. In this case the abelianization is \mathbb{R}^{2n} . Other examples are products of such groups, and $(\mathcal{H}_n/\mathbb{Z}) \times \mathbb{R}^d$.

Remarks 1.6 1. The set \mathcal{S} of cocycles considered in Theorems 1.4 and 1.5 is exactly the one to which our conjecture [6] refers: those that are not cohomologous to a cocycle with values in a maximal semigroup with non-empty interior.

2. In order to prove Theorem 1.4, it is enough to show that $\mathcal{S}^r(X, \mathcal{H}_n)$ contains a dense set of cocycle that are transitive (see for example the introduction in [8]).

3. As was the case in [8] for certain Euclidean-type groups (e.g. $SE(3)$), replacing generic by open and dense in Theorem 1.4 remains an unsolved problem.

4. The proof of Theorem 1.4 develops further the techniques in [8]. In addition, results from the classical theory of Diophantine approximation come into play. It is not hard to extend Theorem 1.4 to products of Heisenberg groups. We

believe that this method works for general two-step nilpotent Lie groups (and perhaps also for nilpotent Lie groups of step higher than two) but there are several technical details that have to be clarified.

The remainder of the paper is organized as follows. In Section 2 we recall some general results from [6], in particular a general criterion for transitivity of extensions of hyperbolic systems. In Section 3 we review and extend results for \mathbb{R}^d -extensions and their periodic data. In Section 4 we prove Theorem 1.5. In Section 5 we recall a technical result from [8]. In Section 6 we specialize to the setting of nilpotent Lie groups and prove Theorem 1.4.

2 Criterion for transitivity

Let Γ be a connected Lie group with Lie algebra $L\Gamma$. We denote by e_Γ the identity element of Γ . Let Ad denote the adjoint representation of Γ on $L\Gamma$. Let $\|\cdot\|$ be a norm on $L\Gamma$. It is known that there is a metric d on Γ with the following properties:

1. $d(\gamma\gamma_1, \gamma\gamma_2) = d(\gamma_1, \gamma_2)$;
2. $d(\gamma_1\gamma, \gamma_2\gamma) \leq \|\text{Ad}(\gamma)\|d(\gamma_1, \gamma_2)$;

for any $\gamma, \gamma_1, \gamma_2 \in \Gamma$.

Definition 2.1 Let $f : X \rightarrow X$ be a map and $\beta : X \rightarrow \Gamma$ a cocycle. We write $f_\beta^k(x, \gamma) = (f^k x, \gamma\beta(k, x))$ where, for $k \geq 1$,

$$\beta(k, x) = \beta(x)\beta(fx) \cdots \beta(f^{k-1}x) = \prod_{j=0}^{k-1} \beta(f^j x).$$

The meaning of the product notation in the last term above is the middle expression.

If Q is a trajectory of f of length k (i.e. $Q = \{x, f(x), \dots, f^{k-1}(x)\}$ for some x), then we define the *height of β over Q* to be $\beta(Q) = \beta(k, x)$. In particular, if x is a periodic point of period ℓ , then the height of the corresponding periodic orbit P is $\beta(P) = \beta(\ell, x)$. The set of heights of β over all periodic orbits of f is referred to as *the periodic data* of β .

Definition 2.2 Given a cocycle $\beta : X \rightarrow \Gamma$ over $f : X \rightarrow X$, define $\mu \geq 1$ to be

$$\mu = \max \left\{ \limsup_{n \rightarrow \infty} \sup_{x \in X} \|\text{Ad}(\beta(n, x))\|^{1/n}, \limsup_{n \rightarrow \infty} \sup_{x \in X} \|\text{Ad}(\beta(n, x))^{-1}\|^{1/n} \right\}.$$

We say that the cocycle β has *subexponential growth* if $\mu = 1$.

Remark 2.3 The subexponential growth condition is automatically satisfied for any cocycle if the group Γ is compact, nilpotent, or a semidirect product of compact and nilpotent.

One of the key notions used in this paper was introduced in [6]:

Definition 2.4 Let Γ be a connected Lie group, X a hyperbolic basic set for $f : X \rightarrow X$, $\beta : X \rightarrow \Gamma$ a cocycle, and $f_\beta : X \times \Gamma \rightarrow X \times \Gamma$ the skew-extension. Given $x \in X$, let

$$\mathcal{L}_\beta(x) = \{\gamma \in \Gamma : \text{there exist } x_k \in X \text{ and } n_k > 0 \text{ such that} \\ x_k \rightarrow x \text{ and } f_\beta^{n_k}(x_k, e_\Gamma) \rightarrow (x, \gamma)\}.$$

That is, the set $\mathcal{L}_\beta(x)$ consists of the possible limits $\lim_{k \rightarrow \infty} \beta(n_k, x_k)$, subject to $x_k \rightarrow x$ and $f^{n_k}(x_k) \rightarrow x$. Note that we do not require that $n_k \rightarrow \infty$ or that $x_k \neq x$. Clearly $\mathcal{L}_\beta(x)$ is a closed subset of Γ .

The following theorem is a special case of [6, Lemma 3.1, Theorem 3.3].

Theorem 2.5 *Assume that X is a hyperbolic basic set for $f : X \rightarrow X$, that Γ is a connected Lie group and that $\beta : X \rightarrow \Gamma$ is a Hölder cocycle that has subexponential growth. Then*

- (a) $\mathcal{L}_\beta(x)$ is a closed semigroup of Γ for each $x \in X$.
- (b) β is a transitive cocycle if and only if there exists a point $x_0 \in X$ such that $\mathcal{L}_\beta(x_0) = \Gamma$.

Denote by $W^s(x)$ ($W_{loc}^s(x)$) and $W^u(x)$ ($W_{loc}^u(x)$) the (local) stable and (local) unstable leaves of f through x . The next lemma is a consequence of [11, Appendix A].

Lemma 2.6 *Assume that X is a hyperbolic basic set for $f : X \rightarrow X$, that Γ is a connected Lie group and that $\beta : X \rightarrow \Gamma$ is an α -Hölder cocycle that has subexponential growth. Then the Γ -extension $f_\beta : X \times \Gamma \rightarrow X \times \Gamma$ admits stable and unstable foliations which are α -Hölder and invariant under right multiplication by elements of Γ . The stable and unstable leaves of f_β through $(x, e_\Gamma) \in X \times \Gamma$ are the graphs of the functions*

$$\begin{aligned} \gamma_x^s : W^s(x) &\rightarrow \Gamma, & \gamma_x^s(y) &= \lim_{n \rightarrow \infty} \beta(n, x) \beta(n, y)^{-1}, \\ \gamma_x^u : W^u(x) &\rightarrow \Gamma, & \gamma_x^u(y) &= \lim_{n \rightarrow \infty} \beta(-n, x) \beta(-n, y)^{-1}. \end{aligned}$$

These functions are α -Hölder and vary continuously with the cocycle β in the following sense: if $\beta_k \rightarrow \beta$ in C^0 -topology and β_k remains C^α -bounded, then $\gamma_{k,x}^s \rightarrow \gamma_x^s$ on $W_{loc}^s(x)$ and $\gamma_{k,x}^u \rightarrow \gamma_x^u$ on $W_{loc}^u(x)$ in C^0 -topology.

We call the values of the functions γ_x^s, γ_x^u *holonomies along stable/unstable leaves*.

The following lemma is a special case of [6, Lemma 2.2].

Lemma 2.7 *Assume that X is a hyperbolic basic set for $f : X \rightarrow X$, that Γ is a connected Lie group and that $\beta : X \rightarrow \Gamma$ is an α -Hölder cocycle that has subexponential growth. Then there is a constant $C > 0$ with the following property.*

Given $\varepsilon > 0$ sufficiently small and $n \geq 1$, assume that there are two trajectories $x_k = f^k(x_0), y_k = f^k(y_0)$, such that $d_M(x_k, y_k) < \varepsilon$ for $0 \leq k \leq n - 1$. Then

$$d(\beta(n, x_0), \beta(n, y_0)) \leq C(\|Ad(\beta(n, x_0))\| + 1)\varepsilon^\alpha. \quad (2.1)$$

3 \mathbb{R}^d -valued cocycles and periodic data

In this section, we review and extend results on \mathbb{R}^d -extensions. As always, $X \subset M$ is a hyperbolic basic set for $f : M \rightarrow M$. Let $\mathcal{S}^r(X, \mathbb{R}^d)$ be the set of C^r cocycles $\beta : X \rightarrow \mathbb{R}^d$ that are *not* cohomologous to a cocycle with values in a half-space in \mathbb{R}^d . More generally, if K is a connected compact Lie group, denote by $\mathcal{S}^r(X, \mathbb{R}^d \times K)$ the C^r -cocycles $\beta : X \rightarrow \mathbb{R}^d \times K$ whose \mathbb{R}^d -component is not cohomologous to a cocycle with values in a half-space in \mathbb{R}^d . A necessary condition for transitivity is that $\beta \in \mathcal{S}^r(X, \mathbb{R}^d \times K)$.

First we recall a result of [10, 2, 7].

Theorem 3.1 *There is an open and dense set $\mathcal{U} \subset \mathcal{S}^r(X, \mathbb{R}^d \times K)$ such that if $\beta \in \mathcal{U}$, then $\beta : X \rightarrow \mathbb{R}^d \times K$ is transitive.*

Next, we prove that one can prescribe periodic data for Hölder \mathbb{R}^d -cocycles. This is needed in the proof of Theorem 1.4.

Proposition 3.2 *Suppose that $\beta : X \rightarrow \mathbb{R}^d$ is a Hölder cocycle and $U \subset X$ is an open set.*

- (a) *Assume that β is transitive. Then the periodic data of β on orbits that intersect U is dense in \mathbb{R}^d .*

(b) Assume that β is stably transitive. Then, given any $\varepsilon > 0$ and $a \in \mathbb{R}^d$, there exists a C^r cocycle $\beta' : X \rightarrow \mathbb{R}^d$ ε -close to β in the C^r -topology, such that β and β' differ only on U and a is the periodic data of β' on a periodic orbit intersecting U .

Proof (a) Pick an open set $V \subset U$ with $\bar{V} \subset U$. Let $a \in \mathbb{R}^d$ and $\delta > 0$. By the transitivity of β , there is a point $x_0 \in V$ and an orbit segment $x_0, f(x_0), \dots, f^N(x_0)$ such that $d_M(x_0, f^N(x_0)) < \delta$ and $\|\beta(N, x_0) - a\| < \delta$. Anosov's Closing Lemma [3] implies that there is a periodic orbit $y_0, f(y_0), \dots, f^N(y_0) = y_0$ such that $d_M(f^k(x_0), f^k(y_0)) < C\delta$ for $0 \leq k \leq N - 1$, where C depends only on f . The estimate (2.1) shows then that $\beta(N, y_0)$ is within $C'\delta^\alpha$ of a , where C' depends only on f and $\|a\|$. If δ is small enough then $y_0 \in U$, and the conclusion follows.

For part (b), fix a C^∞ bump function $g : X \rightarrow [0, 1]$ supported in U such that $g|_V \equiv 1$ for an open set $V \subset U$. Let $M = \|g\|_{C^r}$. By part (a), there exists a periodic orbit P intersecting V with height $\beta(P) = b$ such that $\|a - b\| < \varepsilon/M$. Let $c = g(P)$ be the height of g on P . Since P intersects V , we have $c \geq 1$. Define $\beta' = \beta + (a - b)g/c$. Then $\|\beta' - \beta\|_{C^r} \leq \|a - b\| \|g\|_{C^r} < \varepsilon$, and $\beta'(P) = \beta(P) + (a - b) = a$ as required. ■

4 Open and dense transitivity (compact commutator subgroup)

In this section we prove Theorem 1.5. We start with a preliminary result that is of independent interest.

Proposition 4.1 *Let Γ be a connected finite dimensional Lie group. Suppose that*

(a) *N is a compact normal subgroup of Γ with quotient group H and natural projection $\pi : \Gamma \rightarrow H$.*

(b) *The only closed connected subgroup $\Gamma' \subset \Gamma$ satisfying $\pi(\Gamma') = H$ is $\Gamma' = \Gamma$.*

Assume that the cocycle $\beta : X \rightarrow \Gamma$ has subexponential growth, and let $\beta_H : X \rightarrow H$ be the cocycle obtained after quotienting by N .

Then $\beta : X \rightarrow \Gamma$ is transitive if and only if $\beta_H : X \rightarrow H$ is transitive.

Proof We begin with the observation:

(*) Suppose that (i) Γ' is a closed *semigroup* in Γ and (ii) $\pi(\Gamma') = H$. Then $\Gamma' = \Gamma$.

For suppose that $\gamma \in \Gamma'$. By (ii), there exists $n \in N$ such that $\gamma^{-1}n \in \Gamma'$. Hence $n \in \Gamma'$. But n lies in the compact group N , so the closed semigroup generated by n is a group. In particular $n^{-1} \in \Gamma'$ and so $\gamma^{-1} \in \Gamma'$. Combined with (i), this shows that Γ' is a closed subgroup of Γ . By (ii), $\pi(\Gamma') = H$. By hypothesis (b), $\Gamma' = \Gamma$ as required.

Choose $x_0 \in X$. By Theorem 2.5(b), it suffices to show that $\mathcal{L}_\beta(x_0) = \Gamma$ if and only if $\mathcal{L}_{\beta_H}(x_0) = H$. Clearly, if $\mathcal{L}_\beta(x_0) = \Gamma$, then $\mathcal{L}_{\beta_H}(x_0) = H$. It remains to prove the converse.

So, suppose that $\mathcal{L}_{\beta_H}(x_0) = H$. We show that $\Gamma' = \mathcal{L}_\beta(x_0)$ satisfies the hypotheses of (*), and so $\mathcal{L}_\beta(x_0) = \Gamma$ as required. Now (i) is immediate by Theorem 2.5(a). Let $\gamma \in \Gamma$. Then $\gamma N \in H$ and there exists $x_k \in X$ and $m_k > 0$ such that $f^{m_k}x_k \rightarrow x_0$ and $\beta_H(x_k, m_k) \rightarrow \gamma N$. Hence $\beta(x_k, m_k) = \gamma_k n_k$ where $\gamma_k \rightarrow \gamma$ and $n_k \in N$. Since N is compact, we can pass to a subsequence so that $n_k \rightarrow n \in N$. Hence $\gamma n \in \mathcal{L}_\beta(x_0)$. Thus (ii) is satisfied. ■

Remark 4.2 If $A \subset G$ is a subgroup of a nilpotent group, then $A[G, G] = G$ implies that $A = G$ (see [4, Theorem 16.2.5]).

Proof of Theorem 1.5

Since Γ is nilpotent, cocycles $\beta : X \rightarrow \Gamma$ automatically have subexponential growth. By Remark 4.2, Γ and its compact commutator subgroup $N = [\Gamma, \Gamma]$ satisfy the hypotheses of Proposition 4.1. (This is not so hard to check directly for $\Gamma = \mathcal{H}_n/\mathbb{Z}$.)

Let $\beta \in \mathcal{S}^r(X, \Gamma)$. So the quotient cocycle $\tilde{\beta} : X \rightarrow \mathbb{R}^a$ is *not* cohomologous to a cocycle with values in a half-space in \mathbb{R}^a , where $\Gamma/[\Gamma, \Gamma] \cong \mathbb{R}^a \times \mathbb{T}^b$. By Theorem 3.1, there is an open and dense set $\mathcal{U} \subset \mathcal{S}^r(X, \Gamma)$ such that if $\beta \in \mathcal{U}$, then $\hat{\beta} : X \rightarrow \Gamma/[\Gamma, \Gamma]$ is transitive. By Proposition 4.1, $\beta : X \rightarrow \Gamma$ is also transitive. ■

5 Elements of \mathcal{L}_β

In this section we recall a method to obtain elements of \mathcal{L}_β introduced in [8]. Throughout, (M, d_M) is a Riemannian manifold, $X \subset M$ a hyperbolic basic set for $f : X \rightarrow X$, Γ a connected Lie group and $\beta : X \rightarrow \Gamma$ a Hölder cocycle that has subexponential growth.

Definition 5.1 By a *periodic heteroclinic cycle* we mean a cycle consisting of points p_1, \dots, p_k that are periodic for the map f , have disjoint trajectories, and such that p_j is transverse heteroclinic to p_{j+1} through a point $\zeta_j \in W^u(p_j) \cap W^s(p_{j+1})$, for $j = 1, \dots, k$ (where $p_{k+1} = p_1$).

Let P_1, \dots, P_k be the corresponding periodic orbits and denote the periods by ℓ_1, \dots, ℓ_k . Denote by O_j the heteroclinic trajectory from p_j to p_{j+1} (of the point ζ_j chosen above), and by H_j the holonomy along this heteroclinic connection (that is, along $W^u(p_j)$ from p_j to ζ_j and then along $W^s(p_{j+1})$ from ζ_j to p_{j+1}).

Replace the heteroclinic orbit O_j from p_j to p_{j+1} by the trajectory Q_j of length $\ell_j M_j + \ell_{j+1} M_{j+1}$ that spends time $\ell_j M_j$ in the first half of O_j and time $\ell_{j+1} M_{j+1}$ in the second half of O_j ; that is, $Q_j = \{f^n(\zeta_j) \mid -\ell_j M_j \leq n < 0\} \cup \{f^n(\zeta_j) \mid 0 \leq n < \ell_{j+1} M_{j+1}\}$. For the trajectory connecting p_k to p_{k+1} , we allow M_1 and M_{k+1} to be distinct. The positive integers M_j will be chosen later.

Consider the heights $\beta(P_j)$ and $\beta(Q_j)$ over the periodic orbits P_j and trajectories Q_j (see Definition 2.1).

Lemma 5.2 For $j = 1, \dots, k$, the limit

$$\lim_{M_j, M_{j+1} \rightarrow \infty} \beta(P_j)^{-M_j} \beta(Q_j) \beta(P_{j+1})^{-M_{j+1}} = H_j$$

exists and is the product of the holonomies along the unstable and stable leaves of O_j , from p_j to p_{j+1} .

Proof This follows from Lemma 2.6. ■

Definition 5.3 Consider a sequence of vectors $N(1), N(2) \dots \in \mathbb{N}^{k+1}$ whose entries are positive integers. Write $N(i) = (M_1(i), \dots, M_{k+1}(i))$. The sequence is *admissible* if there is a constant $C_2 \geq 1$ such that $M_p(i)/M_q(i) \leq C_2$ for all $p, q = 1, \dots, k+1$ and all $i \geq 1$.

If $N = (M_1, \dots, M_{k+1})$ is a sequence of vectors, we write $N \rightarrow \infty$ if $M_p \rightarrow \infty$ for each $p = 1, \dots, k+1$. (For an admissible sequence, this is equivalent to $M_p \rightarrow \infty$ for at least one value of p .)

The following result is [8, Theorem 3.4].

Theorem 5.4 Let $N = (M_1, \dots, M_{k+1}) \in \mathbb{N}^{k+1}$. Define

$$A(N) = \beta(P_1)^{M_1} H_1 \beta(P_2)^{2M_2} H_2 \dots \beta(P_k)^{2M_k} H_k \beta(P_1)^{M_{k+1}}.$$

If the limit $A = \lim_{N \rightarrow \infty} A(N)$ exists along an admissible sequence $N(1), N(2), \dots$, then $A \in \mathcal{L}_\beta(p_1)$.

6 Transitivity for $\Gamma = \mathcal{H}_n$

In this section, we return to the Heisenberg group $\Gamma = \mathcal{H}_n$ and prove our main result Theorem 1.4. First, we specialize the methods of Section 5 to this situation.

As in Section 1, we denote by $\widehat{\beta} : X \rightarrow \mathbb{R}^{2n}$ the projection of $\beta : X \rightarrow \mathcal{H}_n$.

By Theorem 3.1 and Proposition 3.2(b), we can assume without loss of generality that the following holds for β and some periodic orbit P_1 :

$$\begin{aligned} \beta \text{ is stably transitive} \\ \widehat{\beta}(P_1) = (0, 0) \in \mathbb{R}^{2n}. \end{aligned} \tag{6.1}$$

We will make a finite number of arbitrarily small, localized, changes to the cocycle. At each step of this process, the property (6.1) will be preserved. The main step is the following:

Lemma 6.1 *Let $\beta \in \mathcal{S}^r(X, \mathcal{H}_n)$ satisfying (6.1) and $\varepsilon > 0$. For any $U \subset X$ open and nonempty, and any vectors $e, f \in \mathbb{R}^n$ there exists $\beta' : X \rightarrow \mathcal{H}_n$ ε -close to β in the C^r topology and $h, k \in \mathbb{R}^n$ with the properties:*

- (a) *for all $u, v \in \mathbb{Z}$ there exists $c \in \mathbb{R}$ such that $(h + ue, k + vf, c) \in \mathcal{L}_{\beta'}(p_1)$;*
- (b) *β' satisfies (6.1);*
- (c) *the vector $(h, k) \in \mathbb{R}^{2n}$ is the sum of holonomies for $\widehat{\beta}'$ along a heteroclinic cycle; therefore, it can be modified by localized changes of the cocycle, while preserving properties (a) and (b) above;*
- (d) *β' differs from β only on U .*

Proof We consider a concatenation of six periodic orbits:

$$A(N) = \beta(P_1)^{M_1} H_1 \beta(P_2)^{M_2} H_2 \dots \beta(P_6)^{M_6} H_6 \beta(P_1)^{M_7}, \tag{6.2}$$

where P_1 is the orbit used in (6.1) and $N = (M_1, \dots, M_7) \in \mathbb{N}^7$. Using the notation in (1.1), write $\beta(P_i) = (a_i, b_i, c_i)$, $H_i = (h_i, k_i, \ell_i)$, $i = 1, \dots, 6$. Then

$$\beta(P_i)^{M_i} H_i = (M_i a_i + h_i, M_i b_i + k_i, M_i c_i + \ell_i + \frac{1}{2} M_i (M_i - 1) a_i^T b_i + M_i a_i k_i). \tag{6.3}$$

Let $h = h_1 + \dots + h_6$ and $k = k_1 + \dots + k_6$. Also it is convenient to write $(a_7, b_7, c_7) = (a_1, b_1, c_1)$. Using (6.2) and (6.3), we compute that $A(N) = (Z_1, Z_2, Z_3)$ where

$$Z_1 = \sum_{i=1}^7 a_i M_i + h, \quad Z_2 = \sum_{i=1}^7 b_i M_i + k, \quad (6.4)$$

and Z_3 is a quadratic polynomial in M_1, \dots, M_7 with leading terms

$$\begin{aligned} & \frac{1}{2} \sum_{j=1}^7 (a_j^T b_j) M_j^2 + \sum_{1 \leq i < j \leq 7} (a_i^T b_j) M_i M_j \\ &= \frac{1}{2} \left(\sum_{i=1}^7 a_i M_i \right)^T \left(\sum_{j=1}^7 b_j M_j \right) + \frac{1}{2} \sum_{1 \leq i < j \leq 7} (a_i^T b_j - a_j^T b_i) M_i M_j. \end{aligned} \quad (6.5)$$

By (6.1) and Proposition 3.2(b), we can prescribe the (a_i, b_i) components of the periodic data of P_2, P_3, P_4, P_5, P_6 by making arbitrarily C^r -small perturbations of β localized in U . We arrange the perturbation so that

$$(a_2, a_3, a_4, a_5, a_6) = (e, 0, -8e, 2e, e), \quad (b_2, b_3, b_4, b_5, b_6) = (0, f, -f, f, 0).$$

We claim that with this choice of $a_i, b_i \in \mathbb{R}^n$, for any $u, v \in \mathbb{Z}$ there is an admissible sequence $(M_1, \dots, M_7) \in \mathbb{N}^7$ that yields $A(N) = (Z_1, Z_2, Z_3) = (h + ue, k + vf, Z_3)$ with Z_3 bounded. Passing to a subsequence, we obtain that $A(N) \rightarrow (h + ue, k + vf, c)$ for some c .

It remains to verify the claim. First set

$$M_1 = M_7 = M_4, \quad M_2 = 4M_4 - M_5 + u, \quad M_3 = M_4 - M_5 + v, \quad M_6 = 4M_4 - M_5. \quad (6.6)$$

In particular, $\sum a_i M_i = ue$ and $\sum b_i M_i = vf$, hence Z_1 and Z_2 take the required values $Z_1 = h + ue$, $Z_2 = k + vf$.

Substituting (6.6) into (6.5) gives a quadratic polynomial in M_4, M_5 . We compute (see Remark 6.2 below) that

$$Z_3 = Q(M_4, M_5) = (e^T f)(4M_4^2 - 8M_4 M_5 + M_5^2) + D_1 M_4 + D_2 M_5 + E, \quad (6.7)$$

where D_1, D_2, E are real coefficients (depending on $a_i, b_i, c_i, h_i, k_i, \ell_i, u, v$).

The discriminant of $Q(M_4, M_5)$ is positive. Moreover the roots of $4x^2 - 8x + 1$ are $1 \pm \frac{1}{2}\sqrt{3}$ which are positive and irrational. We take $\theta = 1 + \frac{1}{2}\sqrt{3}$. By Corollary A.3

we obtain an infinite sequence $(M_4, M_5) \in \mathbb{N}^2$ such that $Z_3 = Q(M_4, M_5)$ is bounded as required and moreover that

$$M_4 \approx \theta M_5. \quad (6.8)$$

Since $\theta > 1$, it follows from (6.6) and (6.8) that the integers M_1, \dots, M_7 are eventually positive (for u and v fixed) and the sequence (M_1, \dots, M_7) is admissible. \blacksquare

Remark 6.2 The computation required to go from (6.5) and (6.6) to (6.7) is routine but a little tedious. It is convenient to organise the calculation as follows.

- (i) The first term in (6.5) reduces to $\frac{1}{2}uv(e^T f)$ and hence contributes only to the constant term E in (6.7).
- (ii) Since $a_1 = b_1 = a_7 = b_7 = 0$, $a_2 = a_6$, $b_2 = b_6$ and $M_2 - M_6 = u$, it is easily verified that the second term in (6.5) contributes to the quadratic part only $\frac{1}{2}\{(a_3^T b_4 - a_4^T b_3)M_3 M_4 + (a_3^T b_5 - a_5^T b_3)M_3 M_5 + (a_4^T b_5 - a_5^T b_4)M_4 M_5\}$ or $(e^T f)(4M_3 M_4 - M_3 M_5 - 3M_4 M_5)$.
- (iii) Substituting $M_3 = M_4 - M_5$ gives the required result.

Remark 6.3 We note certain properties of the perturbations in Lemma 6.1. There was an initial global perturbation which ensures that the quotient cocycle $\widehat{\beta} : X \rightarrow \mathbb{R}^{2n}$ is stably transitive. All further perturbations preserve this property. The next perturbation is local to the vertices p_2, \dots, p_6 of the heteroclinic cycle through p_1 and specifies the heights of $\widehat{\beta}$ at P_2, \dots, P_6 . The values of h, k can then be adjusted by a further perturbation local to the heteroclinic cycle but not affecting the heights at P_1, \dots, P_6 .

Evidently, we may repeat the construction in Lemma 6.1 for any number of heteroclinic cycles at p_1 with localised and hence independent perturbations. In particular, for any $s \geq 1$ and any $e_i, f_i \in \mathbb{R}^n$, $i = 1, \dots, s$, there exists $\beta' : X \rightarrow \mathcal{H}_n$ ε -close to β in the C^r topology, and $(h_i, k_i) \in \mathbb{R}^{2n}$, $i = 1, \dots, s$, given by the holonomies of each cycle, with the property:

$$\text{For all } i = 1, \dots, s, \text{ and all } u, v \in \mathbb{Z}, \text{ there exists } c \in \mathbb{R} \text{ such that} \\ (h_i + ue_i, k_i + vf_i, c) \in \mathcal{L}_{\beta'}(p_1).$$

Moreover, the central ℓ_i components of the holonomies can also be perturbed (independently of the (h_i, k_i) and of the heights $\beta(P_i)$) so that the values of c can be modified as necessary.

Proof of Theorem 1.4 By Remark 6.3, we can choose $(2n + 1)$ pairs $(e_i, f_i) \in \mathbb{R}^{2n}$ that do not lie in a half-space, and by Lemma 6.1 we obtain elements $(h_i + u_i e_i, k_i + v_i f_i, c_i) \in \mathcal{L}_\beta(p_1)$ with $u_i, v_i \in \mathbb{Z}$ arbitrary. In particular, we can choose u_i, v_i large and positive so that the vectors $y_i = (h_i + u_i e_i, k_i + v_i f_i)$ do not lie in a half-space in \mathbb{R}^{2n} . We can make a further perturbation (along the heteroclinic connections) to adjust the values of h_i, k_i so that the vectors y_1, \dots, y_{2n+1} generate \mathbb{R}^{2n} as a closed group and hence (e.g., by [10, Lemma 5] or [7, Lemma 2.12]) as a closed semigroup.

Let $x^* \in \mathbb{R}^n - \{0\}$; for definiteness we take $x^* = (1, 0, \dots, 0)$. By Remark 6.3, we can repeat Lemma 6.1 for another heteroclinic cycle; using the density of the semigroup generated by the y_i 's, we can make an arbitrarily small perturbation to the holonomy of the new cycle to arrange that $(x^*, x^*, c_1) \in \mathcal{L}_\beta(p_1)$ for some $c_1 \in \mathbb{R}$.

Repeat the above step three more times to obtain in $\mathcal{L}_\beta(p_1)$ four elements

$$g_1 = (x^*, x^*, c_1), \quad g_2 = (x^*, -x^*, c_2), \quad g_3 = (-x^*, x^*, c_3), \quad g_4 = (-x^*, -x^*, c_4),$$

where $c_i \in \mathbb{R}$ for $i = 1, \dots, 4$.

Let $c = c_1 + c_2 + c_3 + c_4$. A calculation shows that

$$g_1^m g_2^m g_4^m g_3^m = (0, 0, mc - 2m^2), \quad g_2^m g_1^m g_3^m g_4^m = (0, 0, mc + 2m^2).$$

For m large enough, this yields elements of the form $(0, 0, d_1)$, $(0, 0, d_2)$ with $d_1 < 0$, $d_2 > 0$. If necessary, we can perturb c_1 (say) so that d_1 and d_2 are incommensurate. Since $\mathcal{L}_\beta(p_1)$ is a closed semigroup, we deduce that $\mathcal{L}_\beta(p_1)$ contains the center \mathbb{R} .

Next we return to the elements $(y_1, t_1), \dots, (y_{2n+1}, t_{2n+1})$ where the vectors $y_i = (h_i + u_i e_i, k_i + v_i f_i) \in \mathbb{R}^{2n}$ do not lie in a half-space and generate (as a semigroup) a dense subset of \mathbb{R}^{2n} . Since $\mathcal{L}_\beta(p_1)$ is a closed semigroup containing $\{0\} \times \mathbb{R}$, it follows that $\mathbb{R}^{2n} \times \{0\} \subset \mathcal{L}_\beta(p_1)$ completing the proof. \blacksquare

A Bounds for indefinite binary quadratic polynomials

In this appendix, we recall classical results of Chebyshev and Minkowski that give bounds for indefinite binary quadratic polynomials along sequences of positive integers. A proof of the next theorem, due to Chebyshev, can be found in [12] (only a few steps of [12] are needed to prove (A.1) with 2 replaced by some bounded constant; that is sufficient for our purposes). Corollary A.2 is a weaker form of a result that Minkowski proved in 1901.

Theorem A.1 (Chebyshev, 1866) *Let θ be irrational and ω a real number. Then there exist an infinite number of integer pairs (x, y) , $y > 0$, for which the quantity $|x - \theta y - \omega|$ is made arbitrarily small while*

$$|x - \theta y - \omega| < \frac{2}{y}. \quad (\text{A.1})$$

Corollary A.2 *Let $\theta > 0$ be irrational and ω a real number. Then for any real numbers B_1, B_2, B_3 , there exist constants $\varepsilon \in (0, \theta)$ and $M = M(B_1, B_2, \theta) > 0$, and an infinite set of distinct pairs of positive integers (x, y) that satisfies*

$$\theta - \varepsilon < \frac{x}{y} < \theta + \varepsilon \quad (\text{A.2})$$

and

$$|(x - \theta y - \omega)(B_1 x + B_2 y + B_3)| < M. \quad (\text{A.3})$$

Proof Pick a sequence of integers (x, y) given by Theorem A.1. As $\theta > 0$, we can assume that x and y have the same signs, and thus are positive.

Consider now (A.1) and divide both sides by y . One has

$$\left| \frac{x}{y} - \theta - \frac{\omega}{y} \right| < \frac{2}{y^2},$$

which implies the existence of ε and (A.2).

To prove (A.3), use (A.1), (A.2) and take y sufficiently large so that

$$\begin{aligned} |(x - \theta y - \omega)(B_1 x + B_2 y + B_3)| &\leq 2 \left| B_1 \cdot \frac{x}{y} + B_2 + \frac{B_3}{y} \right| \\ &\leq 2|B_1|(\theta + \varepsilon) + 2|B_2| + 1 = M. \end{aligned}$$

■

Corollary A.3 *Let $Q(x, y) = Ax^2 + Bxy + Cy^2 + Dx + Ey + F$ be a quadratic polynomial with positive discriminant $B^2 - 4AC > 0$. Suppose that there exists $\theta > 0$ irrational such that $A\theta^2 + B\theta + C = 0$. Then there exist constants $\varepsilon \in (0, \theta)$, $M > 0$ and an infinite set of positive integers (x, y) such that*

$$\theta - \varepsilon < \frac{x}{y} < \theta + \varepsilon \quad \text{and} \quad |Q(x, y)| < M.$$

Proof Since $B^2 - 4AC \neq 0$, we can write $Q(x, y) = A\tilde{x}^2 + B\tilde{x}\tilde{y} + C\tilde{y}^2 + \tilde{F}$ where $\tilde{x} = x + a$, $\tilde{y} = y + b$ and $a, b, \tilde{F} \in \mathbb{R}$. Since $B^2 - 4AC > 0$, there are real roots θ_1, θ_2 such that $A\tilde{x}^2 + B\tilde{x}\tilde{y} + C\tilde{y}^2 = A(\tilde{x} - \theta_1\tilde{y})(\tilde{x} - \theta_2\tilde{y})$. Hence there exist $\omega_1, \omega_2 \in \mathbb{R}$ such that

$$Q(x, y) = A(x - \theta_1y - \omega_1)(x - \theta_2y - \omega_2) + \tilde{F}.$$

At least one of the roots, say θ_1 , is irrational and positive, and we set $\theta = \theta_1$. The result follows from Corollary A.2. ■

Acknowledgments: This research was supported in part by EPSRC Grant EP/F031807/1 (IM) and by NSF Grant DMS-0600927 (AT).

References

- [1] T. Bousch. *La condition de Walter*. Ann. Sci. Ecole Norm. Sup. **34** (2001) 287–311.
- [2] M. Field, I. Melbourne, A. Török. *Stable ergodicity for smooth compact Lie group extensions of hyperbolic basic sets*. Ergod. Th. & Dynam. Sys. **25** (2005) 517–551.
- [3] A. Katok, B. Hasselblatt, Introduction to the modern theory of dynamical systems, Cambridge University Press, 1995.
- [4] M. I. Kargapolov, Ju. I Merzljakov, Fundamentals of the theory of groups, Graduate Texts in Mathematics, **62**, Springer-Verlag, 1979.
- [5] I. Melbourne, M. Nicol. *Stable transitivity of Euclidean group extensions*. Ergod. Th. & Dynam. Sys. **23** (2003) 611–619.
- [6] I. Melbourne, V. Nițică, A. Török. *Stable transitivity of certain noncompact extensions of hyperbolic systems*. Annales Henri Poincaré **6** (2005) 725–746.
- [7] I. Melbourne, V. Nițică, A. Török. *A note about stable transitivity of noncompact extensions of hyperbolic systems*. Discrete & Contin. Dynam. Systems **14** (2006) 355–363.
- [8] I. Melbourne, V. Nițică, A. Török. *Transitivity of Euclidean-type extensions of hyperbolic systems*. Ergod. Th. & Dynam. Sys. **29** (2009) 1585–1602.

- [9] V. Nițică. *Stably transitivity for extensions of hyperbolic systems by semidirect products of compact and nilpotent Lie groups*. submitted to Discrete and Continuous Dynamical Systems (2010).
- [10] V. Nițică, M. Pollicott. *Transitivity of Euclidean extensions of Anosov diffeomorphisms*. Ergod. Th. & Dynam. Sys. **25** (2005) 257–269.
- [11] V. Nițică, A. Török. *An open and dense set of stably ergodic diffeomorphisms in a neighborhood of a non-ergodic one*. Topology **40** (2001) 259–278.
- [12] R. Q. Seale. *A new proof of Minkowski's theorem on the product of two linear forms*. Bull. Amer. Math. Soc. **41** (1935) 419–426.
- [13] J. Winkelmann. *Generic subgroups of Lie groups*. Topology **41** (2002) 163–181.