

# Operator renewal theory and mixing rates for dynamical systems with infinite measure

Ian Melbourne <sup>\*</sup>      Dalia Terhesiu <sup>†</sup>

24 August, 2010. Revised 10 August, 2011.  
Minor corrections 5 March, 2015.

## Abstract

We develop a theory of operator renewal sequences in the context of infinite ergodic theory. For large classes of dynamical systems preserving an infinite measure, we determine the asymptotic behaviour of iterates  $L^n$  of the transfer operator. This was previously an intractable problem.

Examples of systems covered by our results include (i) parabolic rational maps of the complex plane and (ii) (not necessarily Markovian) nonuniformly expanding interval maps with indifferent fixed points.

In addition, we give a particularly simple proof of pointwise dual ergodicity (asymptotic behaviour of  $\sum_{j=1}^n L^j$ ) for the class of systems under consideration.

In certain situations, including Pomeau-Manneville intermittency maps, we obtain higher order expansions for  $L^n$  and rates of mixing. Also, we obtain error estimates in the associated Dynkin-Lamperti arcsine laws.

This version includes minor corrections in Sections 10 and 11, and corresponding modifications of certain statements in Section 1. All main results are unaffected. In particular, Sections 2–9 are unchanged from the published version.

**AMS Classification codes:** 37A40, 37A25, 37A50, 60K05

---

<sup>\*</sup>Department of Mathematics, University of Surrey, Guildford, Surrey GU2 7XH, UK

<sup>†</sup>Department of Mathematics, University of Surrey, Guildford, Surrey GU2 7XH, UK

# 1 Introduction

In finite ergodic theory, much recent interest has focussed on the statistical properties of smooth dynamical systems with strong hyperbolicity (expansion/contraction) properties. Landmark results include the proof of exponential decay of correlations for certain classes of uniformly hyperbolic flows [8, 12, 30] and planar dispersing billiards [42]. The latter is part of a general scheme [42, 43] for estimating decay of correlations, or mixing rates, for discrete time dynamical systems.

For systems with subexponential decay of correlations, most approaches yielded only upper bounds for mixing rates. Sarig [37] introduced a powerful new technique, *operator renewal theory*, to obtain precise asymptotics and hence sharp mixing rates. This is an extension of scalar renewal theory from probability theory to the operator situation. The technique was substantially extended and refined by Gouëzel [18, 19].

Garsia & Lamperti [17] developed a theory of renewal sequences with infinite mean in the probabilistic setting. The techniques are very different from the finite mean case, and draw heavily on the theory of regular variation [7, 16, 27]. A natural question is to develop renewal operator theory in the infinite mean case.

**Renewal sequences** In probability theory, renewal sequences relate return probabilities to a specified “nice” set with first return probabilities. The analogue in ergodic theory arises in the study of first return maps.

Let  $(X, \mu)$  be a measure space (finite or infinite), and  $f : X \rightarrow X$  a conservative measure preserving map. Fix  $Y \subset X$  with  $\mu(Y) > 0$ . Let  $\varphi : Y \rightarrow \mathbb{Z}^+$  be the first return time  $\varphi(y) = \inf\{n \geq 1 : f^n y \in Y\}$  (finite almost everywhere by conservativity). Let  $L : L^1(X) \rightarrow L^1(X)$  denote the transfer operator (Perron-Frobenius operator) for  $f$  and define

$$T_n v = 1_Y L^n(1_Y v), \quad n \geq 0, \quad R_n v = 1_Y L^n(1_{\{\varphi=n\}} v), \quad n \geq 1.$$

Note that  $T_n$  and  $R_n$  can be viewed as operators on  $L^1(Y)$  with  $T_0 = I$ . Thus  $T_n$  corresponds to returns to  $Y$  and  $R_n$  corresponds to first returns to  $Y$ . The relationship  $T_n = \sum_{j=1}^n T_{n-j} R_j$  generalises the classical notion of renewal sequences in probability theory.

In the infinite mean setting of Garsia & Lamperti [17], a crucial requirement is that the first return probabilities have regularly varying tails. In our setting it is natural to assume that the return time probabilities have regularly varying tails. Indeed, many of the results in the infinite ergodic theory literature rely crucially on such an assumption [1, 39, 41, 45]. Under this assumption, and certain functional-analytic hypotheses on the operators  $R_n$ , we obtain detailed results (Theorems 2.1, 2.2, 2.3) on the asymptotic behaviour of the operators  $T_n$  as  $n \rightarrow \infty$ . This has strong ramifications for the asymptotics of the iterates  $L^n$ , and hence for the underlying dynamical system.

**Maps with indifferent fixed points** An important class of examples is provided by interval maps with indifferent fixed points, in particular the Pomeau-Manneville

intermittency maps [33] which are uniformly expanding away from an indifferent fixed point at 0. To fix notation, we focus on the version studied by Liverani *et al.* [31]:

$$fx = \begin{cases} x(1 + 2^\alpha x^\alpha), & 0 < x < \frac{1}{2} \\ 2x - 1 & \frac{1}{2} < x < 1 \end{cases}. \quad (1.1)$$

When  $\alpha = 0$ , this is the doubling map, and Lebesgue measure is invariant, ergodic and exponentially mixing. For  $\alpha \in (0, 1)$ , there is still a unique ergodic invariant probability measure  $\mu$  absolutely continuous with respect to Lebesgue measure, but the rate of mixing is polynomial:  $\int_X v w \circ f^n d\mu - \int_X v d\mu \int_X w d\mu \leq C_{v,w} n^{-(\beta-1)}$  where  $\beta = \frac{1}{\alpha}$  for all  $w \in L^\infty(X)$  and all  $v$  sufficiently regular (for example, Hölder continuous). Hu [25] proved that this decay rate is optimal; a special case of the theory of [18, 37].

For  $\alpha \geq 1$ , we are in the situation of infinite ergodic theory. There no longer exists an absolutely continuous invariant probability measure, but there is a unique (up to scaling)  $\sigma$ -finite, absolutely continuous invariant measure  $\mu$ . Previous studies established *pointwise dual ergodicity*:  $a_n^{-1} \sum_{j=1}^n L^j v \rightarrow \text{const} \int_X v d\mu$  almost everywhere for all  $v \in L^1(X)$ , where  $a_n = n^\beta$  for  $\beta \in (0, 1)$  and  $a_n = n/\log n$  when  $\beta = 1$ .

An important refinement is the limit theorem of Thaler [38] where the convergence of  $a_n^{-1} \sum_{j=1}^n L^j v$  is shown to be uniform on compact subsets of  $(0, 1]$  for all observables of the form  $v = u/h$  where  $u$  is Riemann integrable and  $h$  is the density.

The results of [38] are formulated for Markov maps of the interval with indifferent fixed points. Zweimüller [44, 45] relaxed the Markov condition and systematically studied a large class of non-Markovian nonuniformly expanding interval maps, called *AFN maps*. (See Section 11.3 for a precise definition of AFN map.) In particular, [44] obtained a spectral decomposition into basic (conservative and ergodic) sets and proved that for each basic set there is a  $\sigma$ -finite absolutely continuous invariant measure, unique up to scaling. The results in [38] on uniform pointwise dual ergodicity were extended in [45] to the class of AFN maps.

Understanding the asymptotics of  $L^n$ , rather than  $\sum_{j=1}^n L^j$ , turns out to be a much more difficult problem, even for (1.1). Previously, the sole success in this direction was obtained by Thaler [40]. However, the class of systems covered by [40] is quite restrictive and includes the family (1.1) only for  $\beta = 1$ .

It is this situation that we have sought to redress in this paper. It is convenient to describe our main results in the setting of AFN maps  $f : X \rightarrow X$ , though our general theory goes much further, as described later on. Let  $X' \subset X$  denote the complement of the indifferent fixed points. For any compact subset  $A \subset X'$ , the construction in [44] yields a suitable first return set  $Y$  containing  $A$ . Fix such a set  $Y$  with first return time function  $\varphi : Y \rightarrow \mathbb{Z}^+$ . Then we assume that the tail probabilities are regularly varying:  $\mu(\varphi > n) = \ell(n)n^{-\beta}$  where  $\beta \in (0, 1]$  and  $\ell(x)$  is slowly varying ( $\ell : (0, \infty) \rightarrow (0, \infty)$  is measurable and  $\lim_{x \rightarrow \infty} \ell(\lambda x)/\ell(x) = 1$  for all  $\lambda > 0$ ). (For (1.1),  $\ell$  is asymptotically constant and  $\beta = \frac{1}{\alpha}$ .)

Now we can state our results for AFN maps. Set  $d_\beta = \frac{1}{\pi} \sin \beta\pi$  for  $\beta \in (0, 1)$  and  $d_1 = 1$ . Define  $m(x) = \ell(x)$  for  $\beta \in (0, 1)$  and  $m(x) = \sum_{j=1}^{\lfloor x \rfloor} \ell(j)j^{-1}$  for  $\beta = 1$ .

**Theorem 1.1** *Suppose that  $f : X \rightarrow X$  is a topologically mixing AFN map with  $\sigma$ -finite absolutely continuous invariant measure  $\mu$  (with density  $h$ ) and regularly varying tail probabilities.*

(a) *If  $\beta \in (\frac{1}{2}, 1]$ , then  $\lim_{n \rightarrow \infty} m(n)n^{1-\beta}L^n v = d_\beta \int_X v d\mu$  uniformly on compact subsets of  $X'$  for all  $v : X \rightarrow \mathbb{R}$  of the form  $v = u/h$  where  $u$  is Riemann integrable on  $X$ .*

(b) *If  $\beta \in (0, 1]$ , then for  $v : X \rightarrow \mathbb{R}$  that is  $\mu$ -integrable and Riemann integrable, there is a subset  $E \subset \mathbb{N}$  of zero density such that  $\lim_{n \rightarrow \infty, n \notin E} m(n)n^{1-\beta}L^n v = d_\beta \int_X v d\mu$  pointwise on  $X'$ .*

*Moreover, if  $v \geq 0$ , then  $\liminf_{n \rightarrow \infty} m(n)n^{1-\beta}L^n v = d_\beta \int_X v d\mu$  pointwise on  $X'$ .*

(c) *If  $\beta \in (0, \frac{1}{2})$ , then  $L^n v = O(\ell(n)n^{-\beta})$  uniformly on compact subsets of  $X'$  for all measurable  $v : X \rightarrow \mathbb{R}$  such that  $vh$  is bounded.*

**Remark 1.2** (i) It is known that the asymptotic behaviour of  $L^n$  might be complicated. Chung [9, Section I.10] gives an example of a null recurrent Markov chain for which the ratio of  $n$ -step transition probabilities  $p_{ij}^n/p_{kl}^{(n)}$  has no limit as  $n \rightarrow \infty$  (the regular variation assumption on the return time probabilities is violated). Hajian & Kakutani [21] (see also [3, Proposition 1.4.7]) prove that there always exist *weakly wandering* sets  $W$  of positive measure. For such sets,  $\int_W L^n 1_W d\mu = 0$  for infinitely many  $n$ . (Such indicator functions  $1_W$  do not lie in our class of observables  $v$ ).

(ii) In the special case of the family (1.1), Theorem 1.1(a) recovers the result of Thaler [40] for  $\beta = 1$  and the cases  $\beta < 1$  are new. Parts (b,c) are probably not optimal for (1.1) but are the best one can expect in the general setting, see Remark 2.4.

(iii) In addition to yielding convergence results for  $L^n$  (rather than  $\sum_{j=1}^n L^j$ ) our methods also cover much wider classes of observables than was previously possible. An indicative example is the family (1.1) where  $\beta \in (\frac{1}{2}, 1]$ . There is a constant  $C > 1$  such that  $C^{-1}x^{-\frac{1}{\beta}} \leq h(x) \leq Cx^{-\frac{1}{\beta}}$ . Consider observables of the form  $v(x) = x^q$ . Whereas the results of [38, 45] yield uniform convergence of  $\sum_{j=1}^n L^j v$  (and of  $L^n v$  when  $\beta = 1$ ) on compact subsets of  $(0, 1]$  if and only if  $\beta q \geq 1$ , our results apply for  $\beta(q+1) > 1$  (see Theorem 11.14).

(iv) An immediate consequence of Theorem (1.1)(a) is that if  $\beta \in (\frac{1}{2}, 1]$ ,  $v$  is of the required form  $v = u/h$ , and  $w \in L^1(X)$  is supported on a compact subset of  $X'$ , then  $\lim_{n \rightarrow \infty} m(n) \int_X v w \circ f^n d\mu = d_\beta \int_X v d\mu \int_X w d\mu$ .

(v) The situation changes considerably if  $\int_X v d\mu = 0$ . We have the following result which has no counterpart in standard renewal theory, though Gouëzel [18] proves an analogous (and equally unexpected) result in the case of finite ergodic theory:

**Theorem 1.3** *Suppose that  $f : X \rightarrow X$  is a topologically mixing AFN map with regularly varying tail probabilities,  $\beta \in (0, 1)$ . Suppose that  $v$  is of bounded variation and is supported on a compact subset of  $X'$ . If  $\int_X v d\mu = 0$ , then  $L^n v = O(\ell(n)n^{-\beta})$  uniformly on compact subsets of  $X'$ .*

**Second order asymptotics and rates of mixing** In certain situations, including the family (1.1), the tail probabilities satisfy  $\mu(\varphi > n) = cn^{-\beta} + O(n^{-q})$  for some  $q > 1$ ,  $c > 0$ . It is then possible to obtain higher order expansions of  $L^n v$  on compact sets for bounded variation observables  $v$  supported on a compact subset of  $(0, 1]$ . For example, in the specific case of (1.1),  $\beta \in (\frac{1}{2}, 1)$ , we prove that  $|n^{1-\beta} \int_0^1 v w \circ f^n d\mu - \int_0^1 v d\mu \int_0^1 w d\mu| \leq Cn^{-\gamma}$  where  $\gamma = \min\{1 - \beta, \beta - \frac{1}{2}\}$ . The rate of mixing is sharp for  $\beta \geq \frac{3}{4}$ , and we obtain precise second order asymptotics provided  $\beta > \frac{3}{4}$ .

These and related results such as error rates in the Dynkin-Lamperti arcsine laws [14, 29] are discussed in Section 9.

The remainder of the paper is organised as follows. In Section 2, we describe the general framework for our results on the renewal operators  $T_n$ . Sections 3, 4 and 5 contain the proofs for  $\frac{1}{2} < \beta < 1$ . In Section 6, we cover the case  $\beta = 1$ . In Section 7, we give a particularly elementary, self-contained, proof of pointwise dual ergodicity for all  $\beta$ . In Section 8, we prove our results for  $0 < \beta \leq \frac{1}{2}$ . In Section 9, we formulate and prove results on higher order asymptotics. In Section 10, we show how to pass from  $T_n v = 1_Y L^n(1_Y v)$  to  $L^n$ . Finally in Section 11, we show that our theory applies to large classes of examples including AFN maps (in particular, we prove Theorem 1.1) and systems for which the first return map is Gibbs-Markov. The latter includes parabolic rational maps of the complex plane [6].

**Notation** We use “big O” and  $\ll$  notation interchangeably, writing  $A_n = O(a_n)$  or  $A_n \ll a_n$  as  $n \rightarrow \infty$  if there is a constant  $C > 0$  such that  $\|A_n\| \leq Ca_n$  for all  $n \geq 1$  (for  $A_n$  operators and  $a_n \geq 0$  scalars). We write  $A_n \sim c_n A$  as  $n \rightarrow \infty$  if  $\lim_{n \rightarrow \infty} \|A_n/c_n - A\| = 0$  (for  $A_n, A$  operators and  $c_n > 0$  scalars).

## 2 General framework

Let  $(X, \mu)$  be an infinite measure space, and  $f : X \rightarrow X$  a conservative measure preserving map. Fix  $Y \subset X$  with  $\mu(Y) \in (0, \infty)$  and rescale  $\mu$  so that  $\mu(Y) = 1$ . Let  $\varphi : Y \rightarrow \mathbb{Z}^+$  be the first return time  $\varphi(y) = \inf\{n \geq 1 : f^n y \in Y\}$  and define the first return map  $F = f^\varphi : Y \rightarrow Y$ .

The return time function  $\varphi : Y \rightarrow \mathbb{Z}^+$  satisfies  $\int_Y \varphi d\mu = \infty$ . Our crucial assumption is that the tail probabilities are regularly varying:

$\mu(y \in Y : \varphi(y) > n) = \ell(n)n^{-\beta}$  where  $\ell$  is slowly varying and  $\beta \in (0, 1]$ .

Recall that the transfer operator  $R : L^1(Y) \rightarrow L^1(Y)$  for the first return map  $F : Y \rightarrow Y$  is defined via the formula  $\int_Y Rv w d\mu = \int_Y v w \circ F d\mu$ ,  $w \in L^\infty(Y)$ . Let  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  and  $\bar{\mathbb{D}} = \{z \in \mathbb{C} : |z| \leq 1\}$ . Given  $z \in \bar{\mathbb{D}}$ , we define  $R(z) : L^1(Y) \rightarrow L^1(Y)$  to be the operator  $R(z)v = R(z^\varphi v)$ . Also, for each  $n \geq 1$ , we define  $R_n : L^1(Y) \rightarrow L^1(Y)$ ,  $R_n v = R(1_{\{\varphi=n\}}v)$ . It is easily verified that  $R(z) = \sum_{n=1}^{\infty} R_n z^n$ .

Our assumptions on the first return map  $F : Y \rightarrow Y$  are functional-analytic in nature. We assume that there is a function space  $\mathcal{B} \subset L^\infty(Y)$  containing constant functions, with norm  $\|\cdot\|$  satisfying  $\|v\|_\infty \leq \|v\|$  for  $v \in \mathcal{B}$ , such that for some constant  $C > 0$ :

(H1) For all  $n \geq 1$ ,  $R_n : \mathcal{B} \rightarrow \mathcal{B}$  is a bounded linear operator with  $\|R_n\| \leq C\mu(\varphi = n)$ .

In particular,  $z \mapsto R(z)$  is a continuous family of bounded linear operators on  $\mathcal{B}$  for  $z \in \bar{\mathbb{D}}$ . Since  $R(1) = R$  and  $\mathcal{B}$  contains constant functions, 1 is an eigenvalue of  $R(1)$ . We require:

- (H2) (i) The eigenvalue 1 is simple and isolated in the spectrum of  $R(1)$ .  
(ii) For  $z \in \bar{\mathbb{D}} \setminus \{1\}$ , the spectrum of  $R(z)$  does not contain 1.

Denote the spectral projection corresponding to the simple eigenvalue 1 for  $R(1)$  by  $P$ . Then  $(Pv)(y) \equiv \int_Y v d\mu$ .

## 2.1 Asymptotics of $T_n$

We state our main results for the operators  $T_n$ . Since  $T_n v = 1_Y L^n(1_Y v)$ , we obtain precise results for the convergence of  $L^n v$  on  $Y$  for observables  $v$  supported on  $Y$ . The restriction to  $Y$  is lifted in Section 10. Throughout, we assume regularly varying tails  $\mu(\varphi > n) = \ell(n)n^{-\beta}$  and hypotheses (H1) and (H2). Set  $d_\beta = \frac{1}{\pi} \sin \beta\pi$  for  $\beta \in (0, 1)$  and  $d_1 = 1$ . Define  $m(n) = \ell(n)$  for  $\beta \in (0, 1)$  and  $m(n) = \sum_{j=1}^n \ell(j)j^{-1}$  for  $\beta = 1$ .

In some of our statements, the observable  $v$  is not mentioned. Here, we are speaking of convergence of operators on the Banach space  $\mathcal{B}$ . So for example, Theorem 2.1 states that  $\sup_{v \in \mathcal{B}, \|v\|=1} \|m(n)n^{1-\beta}T_n v - d_\beta \int_Y v d\mu\| \rightarrow 0$  as  $n \rightarrow \infty$ . Since  $\mathcal{B}$  is embedded in  $L^\infty(Y)$ , this immediately implies almost sure convergence (at a uniform rate) on  $Y$ . Redefining sequences on a set of measure zero, we obtain uniform convergence on  $Y$ . For brevity, we will speak of uniform convergence throughout this paper without further comment.

**Theorem 2.1** *If  $\beta \in (\frac{1}{2}, 1]$ , then  $\lim_{n \rightarrow \infty} m(n)n^{1-\beta}T_n = d_\beta P$ .*

The next result gives upper bounds on the decay rate of  $T_n$  for  $\beta \leq \frac{1}{2}$ , and an improved upper bound for  $\beta \geq \frac{1}{2}$  when the observable is of mean zero.

**Theorem 2.2** (a) If  $\beta = \frac{1}{2}$ , then  $T_n \ll \ell(n)n^{-\frac{1}{2}} \int_{1/n}^{\pi} \ell(1/\theta)^{-2}\theta^{-1} d\theta$ .

(b) If  $\beta \in (0, \frac{1}{2})$ , then  $T_n \ll \ell(n)n^{-\beta}$ .

(c) If  $\beta \in (0, 1)$ , and  $v \in \mathcal{B}$  satisfies  $\int_Y v d\mu = 0$ , then  $T_n v \ll \ell(n)n^{-\beta}$ .

As indicated in [17], the estimate in Theorem 2.2(b) is essentially optimal. However, we recover certain aspects of Theorem 2.1 even for  $\beta \leq \frac{1}{2}$ . Recall that  $E \subset \mathbb{N}$  has *density zero* if  $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n 1_E(j) = 0$ .

**Theorem 2.3** Let  $\beta \in (0, \frac{1}{2}]$  and  $v \in \mathcal{B}$ .

(a) For all  $y \in Y$ , there exists a set  $E$  of zero density such that  $\lim_{n \rightarrow \infty, n \notin E} \ell(n)n^{1-\beta}(T_n v)(y) = d_\beta \int_Y v d\mu$ .

(b) If  $v \geq 0$ , then  $\liminf_{n \rightarrow \infty} \ell(n)n^{1-\beta}T_n v = d_\beta \int_Y v d\mu$  pointwise on  $Y$ .

**Remark 2.4** In general, Theorem 2.1 fails for  $\beta \leq \frac{1}{2}$ . However, there is the possibility of proving the result for all  $\beta$  under additional hypotheses. Indeed, Gouëzel [20] informs us that he is able to prove Theorem 2.1 for all  $\beta \in (0, 1)$  under the additional assumption that  $\mu(\varphi = n) = O(\ell(n)n^{-(\beta+1)})$ . In particular, Gouëzel's result applies to the family (1.1).

It is worth recalling the situation from the scalar case (where the  $T_n$  are probabilities instead of operators). Under the additional assumption  $\mu(\varphi = n) = O(\ell(n)n^{-(\beta+1)})$ , Garsia & Lamperti [17] were able to extend Theorem 2.1 to the case  $\beta \in (\frac{1}{4}, \frac{1}{2})$ . This is the only part of [17] that we are unable to generalise to the operator setting. However, an argument of Doney [13] applies to all  $\beta \in (0, 1)$  and according to Gouëzel [20] this argument can be extended to the operator case.

**Remark 2.5** An immediate consequence of Theorem 2.1 is that  $Y$  is a Darling-Kac set whenever  $\beta > \frac{1}{2}$ . We refer to Aaronson [3, Chapter 3] and Aaronson, Denker & Urbanski [6, Section 1] for definitions and numerous consequences. Other consequences include the Dynkin-Lamperti arcsine laws, see Thaler [39]. Indeed our main result significantly simplifies the derivation of the arcsine laws, see [40].

## 2.2 Preliminaries

For convenience, we state Karamata's Theorem on the integration of regularly varying sequences [7, 16].

**Proposition 2.6** Suppose that  $\ell$  is slowly varying.

(a) If  $p > -1$ , then  $\sum_{j=1}^n \ell(j)j^p \sim \frac{1}{p+1}\ell(n)n^{p+1}$  as  $n \rightarrow \infty$ .

(b) The function  $\tilde{\ell}(x) = \sum_{j=1}^{\lfloor x \rfloor} \ell(j)j^{-1}$  is slowly varying and  $\lim_{n \rightarrow \infty} \ell(n)/\tilde{\ell}(n) = 0$ . ■

The following consequence of (H1) and regularly varying tails is standard.

**Proposition 2.7** *There is a constant  $C > 0$  such that  $\|R(\rho e^{i(\theta+h)}) - R(\rho e^{i\theta})\| \leq Cm(1/h)h^\beta$  and  $\|R(\rho) - R(1)\| \leq Cm(\frac{1}{1-\rho})(1-\rho)^\beta$  for all  $\theta \in [0, 2\pi)$ ,  $\rho \in (0, 1]$ ,  $h > 0$ .*

**Proof** We sketch the calculation for the first estimate. By Proposition 2.6,

$$R(\rho e^{i(\theta+h)}) - R(\rho e^{i\theta}) \ll h \sum_{j=1}^N j\mu(\varphi=j) + \sum_{j>N} \mu(\varphi=j) \ll hm(N)N^{1-\beta} + m(N)N^{-\beta},$$

so the result follows with  $N = [h^{-1}]$ .  $\blacksquare$

By (H2), there exists  $\epsilon > 0$  such that  $R(z)$  has a continuous family of simple eigenvalues  $\lambda(z)$  for  $z \in \bar{\mathbb{D}} \cap B_\epsilon(1)$  with  $\lambda(1) = 1$ . Let  $P(z) : \mathcal{B} \rightarrow \mathcal{B}$  denote the corresponding family of spectral projections, with complementary projections  $Q(z) = I - P(z)$  and  $P(1) = P$ . Also, let  $v(z) \in \mathcal{B}$  denote the corresponding family of eigenfunctions normalised so that  $\int_Y v(z) d\mu = 1$  for all  $z$ . In particular,  $v(1) \equiv 1$ .

**Corollary 2.8** *The estimates for  $R(z)$  in Proposition 2.7 are inherited by the families  $P(z)$ ,  $Q(z)$ ,  $\lambda(z)$  and  $v(z)$ , where defined.*

**Proof** This is a standard consequence of perturbation theory for analytic families of operators [28].  $\blacksquare$

We have defined the bounded linear operators  $T_n, R_n : \mathcal{B} \rightarrow \mathcal{B}$  given by

$$T_nv = 1_Y L^n(1_Y v), \quad n \geq 0, \quad R_nv = 1_Y L^n(1_{\{\varphi=n\}}v) = R(1_{\{\varphi=n\}}v), \quad n \geq 1.$$

The power series

$$T(z) = \sum_{n=0}^{\infty} T_n z^n, \quad z \in \mathbb{D}, \quad R(z) = \sum_{n=1}^{\infty} R_n z^n, \quad z \in \bar{\mathbb{D}},$$

are analytic on the open unit disk  $\mathbb{D}$ , and  $R(z)$  is continuous on  $\bar{\mathbb{D}}$  by (H1). We have the usual relation  $T_n = \sum_{j=1}^n T_{n-j} R_j$  for  $n \geq 1$ , and it follows that  $T(z) = I + T(z)R(z)$  on  $\mathbb{D}$ . Hence  $T(z) = (I - R(z))^{-1}$  on  $\mathbb{D}$ . By (H2)(ii),  $T(z)$  extends continuously to  $\bar{\mathbb{D}} - \{1\}$  via the formula  $T(z) = (I - R(z))^{-1}$ .

**Proposition 2.9** *There exists  $\epsilon, C > 0$  such that  $\|T(z) - (1 - \lambda(z))^{-1}P(z)\| \leq C$  for  $z \in \bar{\mathbb{D}} \cap B_\epsilon(1)$ ,  $z \neq 1$ , and  $\|T(z)\| \leq C$  for  $z \in \bar{\mathbb{D}} \setminus B_\epsilon(1)$ .*

**Proof** Choose  $\epsilon > 0$  so that the family of simple eigenvalues  $\lambda(z)$  is defined on  $\bar{\mathbb{D}} \cap B_\epsilon(1)$ . For  $z \in \bar{\mathbb{D}} \cap B_\epsilon(1)$ , we can write  $R(z) = \lambda(z)P(z) + R(z)Q(z)$ . If in addition  $z \neq 1$ , then we have (in an obvious notation)

$$T(z) = (1 - \lambda(z))^{-1}P(z) + (I - R(z))^{-1}Q(z).$$



By (H2), the second term is uniformly bounded in the operator norm, and  $T(z)$  is uniformly bounded for  $z \in \bar{\mathbb{D}} \setminus B_\epsilon(1)$ .  $\blacksquare$

## 2.3 Strategy of the proof of Theorem 2.1

Our aim is to compute the operators  $T_n$  defined above. Most of the analysis is carried out on the unit circle  $S^1$ , so it is convenient to abuse notation, writing  $T(\theta)$  instead of  $T(e^{i\theta})$  and so on. For  $\beta \in (\frac{1}{2}, 1)$ , our treatment follows Garsia & Lamperti [17] but there are some significant differences in two of the three steps.

The first step, Section 3, is to study the singularity for  $T(\theta)$  at  $\theta = 0$ . The argument in [17] is scalar and similar results can be found in [15, 22]. In our situation, the key is to use the fact that the return time  $\varphi$  lies in the domain of a stable law, and a Nagaev-type argument due to Aaronson & Denker [4] shows that  $(1 - \lambda(\theta))^{-1} \sim \text{const } \ell(1/\theta)^{-1} \theta^{-\beta}$  when  $\beta \in (0, 1)$ . By Corollary 2.8 and Proposition 2.9,  $T(\theta) \sim \text{const } \ell(1/\theta)^{-1} \theta^{-\beta} P$ .

In particular  $T(\theta) \in L^1$  with Fourier coefficients  $\widehat{T}_n$ . The second step is to relate  $T_n$  to  $\widehat{T}_n$ . In the scalar case, [17] invokes ideas of Herglotz [24] on analytic functions with positive real part. A different approach is required here since we are working with operators. In Section 4, we verify that  $T_n = \widehat{T}_n$ .

In the final step, Section 5, we investigate the behaviour of  $\widehat{T}_n$  as  $n \rightarrow \infty$ , directly following [17].

## 2.4 Tower extensions

The following tower construction will be required in Sections 7 and 10.

Starting from the first return map  $F = f^\varphi : Y \rightarrow Y$ , there is a standard way of constructing an extension  $f_\Delta : \Delta \rightarrow \Delta$  of the underlying map  $f : X \rightarrow X$ . Define the *tower*  $\Delta = \{(y, j) \in Y \times \mathbb{Z} : 0 \leq j < \varphi(y)\}$  and the *tower map*  $f_\Delta : \Delta \rightarrow \Delta$  given by  $f_\Delta(y, j) = (y, j + 1)$  for  $j \leq \varphi(y) - 2$  and  $f_\Delta(y, j) = (Fy, 0)$  for  $j = \varphi(y) - 1$ .

The base of the tower  $\{(y, 0) : y \in Y\}$  is naturally identified with  $Y$  and so we may regard  $Y$  as a subset of both  $X$  and  $\Delta$ . Let  $\mu_\Delta$  be the unique  $f_\Delta$ -invariant measure on  $\Delta$  that agrees with the underlying measure  $\mu$  on the common subset  $Y$ .

Define the projection  $\pi : \Delta \rightarrow X$ ,  $\pi(y, j) = f^j y$ . Then  $\pi f_\Delta = f \pi$  and  $\pi_* \mu_\Delta = \mu$ . Thus  $f_\Delta$  is an extension of  $f$  with the same first return map  $F : Y \rightarrow Y$  and return time function  $\varphi : Y \rightarrow \mathbb{Z}^+$  as the original map.

## 3 Asymptotics of $T(\theta)$

In this section, we obtain an asymptotic expression for  $T(\theta)$  as  $\theta \rightarrow 0^+$ . Throughout,  $\beta \in (0, 1)$ . The main part of the analysis is to understand the asymptotics of the

leading eigenvalue  $\lambda(\theta)$ . In certain situations, we obtain a higher order expansion. Let  $c_\beta = -i \int_0^\infty e^{i\sigma} \sigma^{-\beta} d\sigma$ .

**Lemma 3.1** *Let  $\beta \in (0, 1)$ . As  $\theta \rightarrow 0^+$ ,*

$$(a) \lambda(\theta) = 1 - c_\beta \ell(1/\theta) \theta^\beta (1 + o(1)).$$

$$(b) T(\theta) - (1 - \lambda(\theta))^{-1} P = O(1).$$

$$(c) T(\theta) = c_\beta^{-1} \ell(1/\theta)^{-1} \theta^{-\beta} (1 + o(1)) P + O(1).$$

**Proof** (a) This is part of [4, Theorem 5.1]. (The main ideas of the proof are reproduced later in the proof of Lemma 3.2 and Lemma 4.1.)

(b) By Proposition 2.9,

$$T(\theta) = (1 - \lambda(\theta))^{-1} P(\theta) + O(1) = (1 - \lambda(\theta))^{-1} P + (1 - \lambda(\theta))^{-1} (P(\theta) - P) + O(1).$$

By Corollary 2.8,  $P(\theta) - P \ll \ell(1/\theta) \theta^\beta$ , so the result follows from (a).

(c) is immediate by (a) and (b). ■

The following expansion for  $\lambda(\theta)$  will be used in proving results on second order asymptotics (Section 9).

**Lemma 3.2** *Suppose that  $\mu(\varphi > n) = c(n^{-\beta} + H(n))$ , where  $c > 0$  and  $H(n) = O(n^{-q})$ ,  $q > 1$ . Let  $H_1(x) = [x]^{-\beta} - x^{-\beta} + H([x])$  and set  $c_H = \int_0^\infty H_1(x) dx$ . Then*

$$\lambda(\theta) = 1 - cc_\beta \theta^\beta + icc_H \theta + O(\theta^{2\beta}, \theta^q), \text{ as } \theta \rightarrow 0^+.$$

**Proof** We follow [4, Theorem 5.1]. Recall that  $v(\theta)$  is the eigenfunction corresponding to  $\lambda(\theta)$  normalised so that  $\int v(\theta) d\mu = 1$ . Since  $v(0) \equiv 1$ , it follows from Corollary 2.8 that  $|v(\theta) - 1|_\infty \ll \theta^\beta$ . Let  $\mathcal{F}_0$  denote the  $\sigma$ -algebra generated by  $\varphi$ . Define the step function  $\hat{v}(\theta) : [0, \infty) \rightarrow \mathbb{C}$  given by  $\hat{v}(\theta) \circ \varphi = E(v(\theta) | \mathcal{F}_0)$  and the distribution function  $G(x) = \mu(\varphi \leq x)$ . Then

$$\begin{aligned} \lambda(\theta) &= \int_Y R(\theta) v(\theta) d\mu = \int_Y e^{i\theta\varphi} v(\theta) d\mu = 1 + \int_Y (e^{i\theta\varphi} - 1) v(\theta) d\mu \\ &= 1 + \int_0^\infty (e^{i\theta x} - 1) \hat{v}_\theta(x) dG(x), \end{aligned}$$

where  $\sup_{x \geq 0} |\hat{v}_\theta(x) - 1| \ll \theta^\beta$  and  $1 - G(x) = c(x^{-\beta} + H_1(x))$ . Here  $H_1(x) = O(x^{-q'})$  as  $x \rightarrow \infty$ , where  $q' = \min\{q, \beta + 1\} > 1$ . In particular,  $H_1$  is integrable.

Write  $\hat{v}_\theta = 1 + v_\theta^1 - v_\theta^2 + iv_\theta^3 - iv_\theta^4$ , where  $v_\theta^s \geq 0$  and  $\sup_x |v_\theta^s(x)| \ll \theta^\beta$  for  $s = 1, 2, 3, 4$ . Define the positive measure  $dG_\theta^s = v_\theta^s dG$ . Then

$$\begin{aligned} \lambda(\theta) &= 1 + \int_0^\infty (e^{i\theta x} - 1) dG(x) + \sum_{s=1}^4 q_s \int_0^\infty (e^{i\theta x} - 1) dG_\theta^s(x) \\ &= 1 + i\theta \int_0^\infty e^{i\theta x} (1 - G(x)) dx + \sum_{s=1}^4 q_s i\theta \int_0^\infty e^{i\theta x} g_\theta^s(x) (1 - G(x)) dx, \end{aligned}$$

where  $q_1 = 1, q_2 = -1, q_3 = i, q_4 = -i, g_\theta^s(x) = \int_x^\infty v_\theta^s(u) dG(u) / \int_x^\infty dG(u) \ll \theta^\beta$ .

Now,

$$\begin{aligned} i\theta \int_0^\infty e^{i\theta x}(1 - G(x)) dx &= ic\theta \int_0^\infty e^{i\theta x}x^{-\beta} dx + ic\theta \int_0^\infty e^{i\theta x}H_1(x) dx \\ &= ic\theta^\beta \int_0^\infty e^{i\sigma}\sigma^{-\beta} d\sigma + ic\theta \int_0^\infty H_1(x) dx + ic\theta \int_0^\infty (e^{i\theta x} - 1)H_1(x) dx \\ &= -cc_\beta\theta^\beta + icc_H\theta + ic\theta A \end{aligned}$$

where

$$\begin{aligned} A &= \int_0^\infty (e^{i\theta x} - 1)H_1(x) dx = \int_0^{1/\theta} (e^{i\theta x} - 1)H_1(x) dx + \int_{1/\theta}^\infty (e^{i\theta x} - 1)H_1(x) dx \\ &\ll \theta \int_0^{1/\theta} xH_1(x) dx + \int_{1/\theta}^\infty H_1(x) dx \ll \theta \int_0^{1/\theta} x^{1-q'} dx + \int_{1/\theta}^\infty x^{-q'} dx \ll \theta^{q'-1}. \end{aligned}$$

It remains to estimate the terms  $\int_0^\infty e^{i\theta x}g_\theta^s(x)(1 - G(x)) dx$ . We give the details for  $\int_0^\infty \sin \theta x g_\theta^s(x)(1 - G(x)) dx$ ; the case with  $\sin$  replaced by  $\cos$  is identical. Since  $x \mapsto g_\theta^s(x)(1 - G(x))$  is decreasing for each fixed  $\theta$ , we can write

$$\begin{aligned} \int_0^\infty \sin \theta x g_\theta^s(x)(1 - G(x)) dx &\leq \int_0^{\pi/\theta} \sin \theta x g_\theta^s(x)(1 - G(x)) dx \\ &\ll \theta^\beta \int_0^{\pi/\theta} x^{-\beta} dx \ll \theta^{2\beta-1}, \end{aligned}$$

giving the required upper bound, and the lower bound is obtained in the same way. ■

## 4 Identification of the Fourier coefficients

Let  $\widehat{R}_n$  and  $\widehat{T}_n$  denote the Fourier coefficients of  $R(\theta)$  and  $T(\theta)$ . By (H1),  $R$  is uniformly absolutely summable on  $S^1$ . Therefore  $\widehat{R}_n = R_n$ . In this section, we verify that  $\widehat{T}_n = T_n$  for all  $n \geq 0$ . Throughout,  $\beta \in (0, 1)$ .

**Lemma 4.1** *There exists  $\epsilon, C > 0$  such that  $|1 - \lambda(z)| \geq C\ell(1/\theta)\theta^\beta$  for all  $z = \rho e^{i\theta} \in \mathbb{D} \cap B_\epsilon(1)$ .*

**Proof** We start off by mimicking the proof of Lemma 3.2. Consider functions  $v_z : [0, \infty) \rightarrow [0, \infty)$  satisfying either (i)  $v_z \equiv 1$  or (ii)  $|v_z|_\infty = o(1)$  as  $z \rightarrow 1$ . Write  $z = e^{-u+i\theta}$ ,  $0 \leq u, \theta < \epsilon$ . Then the expansion  $\lambda(z) - 1$  leads to a linear combination of five integrals of the form

$$I = \int_0^\infty (e^{(-u+i\theta)x} - 1)v_z(x) dG(x) = (-u + i\theta) \int_0^\infty e^{(-u+i\theta)x}g_z(x)(1 - G(x))dx, \quad (4.1)$$

where  $g_z \geq 0$  and either (i)  $g_z \equiv 1$  or (ii)  $|g_z|_\infty = o(1)$  as  $z \rightarrow 1$ . Moreover there is one integral of type (i) and we show that this satisfies the desired lower bound, whilst the four integrals of type (ii) are negligible.

Recall that  $1 - G(x) = \mu(\varphi > x) = \ell([x])[x]^{-\beta} = x^{-\beta}h(x)$  where  $h(x) = \ell(x)(1 + o(1))$ . Substituting  $\sigma = \theta x$ ,

$$I = (-u + i\theta)\ell(1/\theta)\theta^{\beta-1} \int_0^\infty e^{-\sigma y} e^{i\sigma} g_z(\sigma/\theta) \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma,$$

where  $y = u/\theta$ . We estimate the oscillatory integrals in the same way that alternating series are estimated in Leibnitz's theorem, making extensive use of the fact that  $\sigma \mapsto e^{-\sigma y} g_z(\sigma/\theta) \sigma^{-\beta} h(\sigma/\theta)$  is decreasing for each fixed  $\theta, y$ .

Write  $|I| = (u^2 + \theta^2)^{1/2} \ell(1/\theta) \theta^{\beta-1} A$ , where in cases (i) and (ii) respectively,

$$\begin{aligned} A_{(i)} &\geq \int_0^{3\pi/2} \cos \sigma e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma, \\ A_{(ii)} &\leq \left( \int_0^{\pi/2} \cos \sigma + \int_0^\pi \sin \sigma \right) e^{-\sigma y} \sigma^{-\beta} g_z(\sigma/\theta) h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma. \end{aligned}$$

We divide the region  $y > 0$  into the regions  $y \in (0, 1/\delta]$  and  $y \geq 1/\delta$  where  $\delta$  is chosen sufficiently small. We have the Potter's bounds [34], [7, Theorem 1.5.6]:  $C^{-1}\sigma^\beta \leq h(\sigma/\theta)\ell(1/\theta)^{-1} \leq C\sigma^{-1}$  uniformly in  $\theta > 0, \sigma \in (0, 2\pi]$ , where  $C$  is a constant. Hence

$$\begin{aligned} A_{(i)} &\geq \frac{\sqrt{2}}{2} \int_0^{\pi/4} e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma + O\left(\int_{\pi/4}^{3\pi/2} e^{-\sigma y} d\sigma\right) \\ &= \frac{\sqrt{2}}{2} \int_0^{\pi/4} e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma + O(y^{-1}e^{-(\pi/4)y}), \end{aligned}$$

$$\begin{aligned} A_{(ii)} &\leq 2|g_z|_\infty \int_0^{\pi/4} e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma + O\left(\int_{\pi/4}^\pi e^{-\sigma y} d\sigma\right) \\ &\ll |g_z|_\infty A_{(i)} + O(y^{-1}e^{-(\pi/4)y}). \end{aligned}$$

Since  $|g_z|_\infty = o(1)$ , we can choose  $\epsilon$  sufficiently small that

$$|1 - \lambda(z)| \gg u\ell(1/\theta)\theta^{\beta-1} \left\{ \int_0^{\pi/4} e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma + O(y^{-1}e^{-(\pi/4)y}) \right\}.$$

Furthermore,

$$\int_0^{\pi/4} e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma \gg \int_0^{\pi/4} e^{-\sigma y} d\sigma = y^{-1} + O(y^{-1}e^{-(\pi/4)y}).$$

For  $y \geq 1/\delta$  with  $\delta$  sufficiently small, the terms  $O(y^{-1}e^{-(\pi/4)y})$  are negligible so that  $1 - \lambda(z) \gg u\ell(1/\theta)\theta^{\beta-1}y^{-1} = \ell(1/\theta)\theta^\beta$ .

It remains to consider the complementary region  $y \in (0, 1/\delta]$ . Note that  $\int_0^{3\pi/2} e^{-\sigma y} \cos \sigma \sigma^{-\beta} d\sigma$  depends continuously on  $y$  and is positive for all  $y \geq 0$ . It follows by compactness that  $\int_0^{3\pi/2} e^{-\sigma y} \cos \sigma \sigma^{-\beta} d\sigma$  is bounded away from zero for  $y \in [0, 1/\delta]$ . Moreover there exists  $b \in (0, 3\pi/2)$  such that  $\int_b^{3\pi/2} e^{-\sigma y} \cos \sigma \sigma^{-\beta} d\sigma$  is bounded away from zero for  $y \in [0, 1/\delta]$ . By uniform convergence of slowly varying functions [7, Theorem 1.2.1], we can shrink  $\epsilon$  if necessary so that  $|h(\sigma/\theta)/\ell(1/\theta) - 1|$  is as small as desired, uniformly in  $\sigma \in [b, 3\pi/2]$  and  $\theta \in (0, \epsilon]$ . Hence  $A_{(i)} \geq \int_0^{3\pi/2} \cos \sigma e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma$  which is bounded away from zero for  $y \in (0, 1/\delta]$ .

Also, by Potter's bounds for any  $\beta' \in (\beta, 1)$ ,

$$A_{(ii)} \leq 2|g_z|_\infty \int_0^\pi e^{-\sigma y} \sigma^{-\beta} h(\sigma/\theta) \ell(1/\theta)^{-1} d\sigma \ll |g_z|_\infty \int_0^\pi \sigma^{-\beta'} d\sigma \ll |g_z|_\infty = o(1).$$

Hence  $A_{(ii)}$  is negligible relative to  $A_{(i)}$  and we obtain  $1 - \lambda(z) \gg \theta \ell(1/\theta) \theta^{\beta-1} = \ell(1/\theta) \theta^\beta$  completing the proof.  $\blacksquare$

It is convenient in the next result (and crucial in Section 6) to discuss the real part of an operator. We recall that the operators  $T_n$  are defined on the real Banach space  $\mathcal{B}$ . Passing to the complexification, there is a natural conjugation  $u+iv \mapsto u-iv$  on  $\mathcal{B}$ . Given an operator  $A : \mathcal{B} \rightarrow \mathcal{B}$ , define the conjugate  $\bar{A} : \mathcal{B} \rightarrow \mathcal{B}$  by setting  $\bar{A}v = \overline{Av}$ , and the real part  $\operatorname{Re} A = \frac{1}{2}(A + \bar{A})$ . In the case of the operator  $T(z) = \sum_{n=0}^\infty T_n z^n$ , this coincides with the definitions  $\overline{T(z)} = T(\bar{z})$  and  $\operatorname{Re} T(z) = \sum_{n=0}^\infty T_n \operatorname{Re}(z^n)$ .

**Corollary 4.2**  $T_n = \widehat{T}_n = \frac{1}{\pi} \operatorname{Re} \int_0^\pi T(e^{i\theta}) e^{-in\theta} d\theta$  for all  $n \geq 0$ .

**Proof** Since  $T(z) = \sum_{j=0}^\infty T_j z^j$  is analytic on the open unit disk  $\mathbb{D}$ ,  $T_n = \frac{1}{2\pi} \rho^{-n} \int_0^{2\pi} T(\rho e^{i\theta}) e^{-in\theta} d\theta$ , for all  $\rho \in (0, 1)$ .

By Proposition 2.9,  $T(z) = O(1)$  on  $\bar{\mathbb{D}} \setminus B_\epsilon(1)$ . Further, on  $\bar{\mathbb{D}} \cap B_\epsilon(1)$ ,  $T(z) = (1 - \lambda(z))^{-1} P(z) + O(1) \ll (1 - \lambda(z))^{-1} + O(1)$ . By Lemma 4.1,  $\|T(\rho e^{i\theta})\| \ll \ell(1/\theta)^{-1} \theta^{-\beta}$  for  $z = \rho e^{i\theta} \in \bar{\mathbb{D}}$  uniformly in  $\rho$ . Since  $\ell(1/\theta)^{-1} \theta^{-\beta}$  is integrable, it follows from the dominated convergence theorem (as  $\rho \rightarrow 1$ ) that  $T_n = \frac{1}{2\pi} \int_0^{2\pi} T(e^{i\theta}) e^{-in\theta} d\theta = \widehat{T}_n$ . Since  $T(\bar{z}) = \overline{T(z)}$ , we obtain the expression  $\frac{1}{\pi} \operatorname{Re} \int_0^\pi T(e^{i\theta}) e^{-in\theta} d\theta$ .  $\blacksquare$

## 5 Convergence for $\beta \in (\frac{1}{2}, 1)$

In this section, we prove Theorem 2.1 for  $\beta \in (\frac{1}{2}, 1)$ .

**Lemma 5.1** *Let  $\beta \in (\frac{1}{2}, 1)$ . Let  $n \geq 1$ ,  $a \in [1, n]$ . Then for any  $\beta' \in (0, \beta)$ ,*

$$\ell(n)n^{1-\beta} \int_{a/n}^{\pi} T(\theta)e^{-in\theta} d\theta \ll a^{-(2\beta'-1)}.$$

*If  $\ell$  is asymptotically constant, then the result holds with  $\beta' = \beta$ .*

**Proof** By Lemma 3.1(c), we have the estimate  $\|T(\theta)\| \ll \ell(1/\theta)^{-1}\theta^{-\beta}$ . The proof uses this fact together with Proposition 2.7, and follows Garsia & Lamperti [17, p. 231]. We give the details partly for completeness and partly because we want to make explicit certain estimates that will be used in Section 9.

First, write

$$I = \int_{a/n}^{\pi} T(\theta)e^{-in\theta} d\theta = - \int_{(a+\pi)/n}^{\pi+\pi/n} T(\theta - \pi/n)e^{-in\theta} d\theta,$$

so

$$2I = \int_{a/n}^{\pi} T(\theta)e^{-in\theta} d\theta - \int_{(a+\pi)/n}^{\pi+\pi/n} T(\theta - \pi/n)e^{-in\theta} d\theta = I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \int_{\pi}^{\pi+\pi/n} T(\theta - \pi/n)e^{-in\theta} d\theta, & I_2 &= \int_{a/n}^{(a+\pi)/n} T(\theta - \pi/n)e^{-in\theta} d\theta, \\ I_3 &= \int_{(a+\pi)/n}^{\pi} \{T(\theta) - T(\theta - \pi/n)\}e^{-in\theta} d\theta. \end{aligned}$$

Clearly,  $I_1 \ll 1/n$ , while

$$\begin{aligned} I_2 &\ll \int_{a/n}^{(a+\pi)/n} \ell(1/\theta)^{-1}\theta^{-\beta} d\theta \ll \ell(n)^{-1}n^{-(1-\beta)} \int_a^{a+\pi} [\ell(n)/\ell(n/\sigma)]\sigma^{-\beta} d\sigma \\ &\ll \ell(n)^{-1}n^{-(1-\beta)} \int_a^{a+\pi} \sigma^{-\beta'} d\sigma = \ell(n)^{-1}n^{-(1-\beta)} a^{1-\beta'} \{(1 + \pi/a)^{1-\beta'} - 1\} \\ &\ll \ell(n)^{-1}n^{-(1-\beta)} a^{-\beta'}. \end{aligned}$$

By the resolvent identity and Proposition 2.7 (with  $m(x) = \ell(x)$ ),

$$\begin{aligned} I_3 &\ll \int_{(a+\pi)/n}^{\pi} \|T(\theta)\| \|T(\theta - \pi/n)\| \|R(\theta) - R(\theta - \pi/n)\| d\theta \\ &\ll \ell(n/\pi)n^{-\beta} \int_{(a+\pi)/n}^{\pi} \ell(1/\theta)^{-1}\ell(1/(\theta - \pi/n))^{-1}\theta^{-\beta}(\theta - \pi/n)^{-\beta} d\theta \\ &= \ell(n/\pi)n^{-\beta} \int_{a/n}^{\pi-\pi/n} \ell(1/(\theta + \pi/n))^{-1}\ell(1/\theta)^{-1}(\theta + \pi/n)^{-\beta}\theta^{-\beta} d\theta. \end{aligned}$$

By Potter's bounds,  $\ell(1/(\theta + \pi/n))^{-1} \ll \ell(1/\theta)^{-1}$  for  $n\theta \geq 1$ . Hence,

$$\begin{aligned} I_3 &\ll \ell(n)n^{-\beta} \int_{a/n}^{\pi} \ell(1/\theta)^{-2}\theta^{-2\beta} d\theta = \ell(n)^{-1}n^{-(1-\beta)} \int_a^{n\pi} [\ell(n)/\ell(n/\sigma)]^2 \sigma^{-2\beta} d\sigma \\ &\ll \ell(n)^{-1}n^{-(1-\beta)} \int_a^{n\pi} \sigma^{-2\beta'} d\sigma \ll \ell(n)^{-1}n^{-(1-\beta)} a^{-(2\beta'-1)}. \end{aligned}$$

Altogether, we obtain  $\ell(n)n^{1-\beta}I \ll n^{-\beta} + a^{-\beta'} + a^{-(2\beta'-1)} \ll a^{-(2\beta'-1)}$  as required. ■

**Lemma 5.2** *Let  $\beta \in (0, 1)$ . Let  $n \geq 1$ ,  $a \in (0, \epsilon n)$ . Then*

$$\lim_{a \rightarrow \infty} \lim_{n \rightarrow \infty} \ell(n)n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta = d'_\beta,$$

where  $d'_\beta = i \int_0^\infty e^{-i\sigma} \sigma^{-\beta} d\sigma / \int_0^\infty e^{i\sigma} \sigma^{-\beta} d\sigma$ .

**Proof** This is identical to [17, Lemma 3.4.1] and we give the proof only for completeness. By Lemma 3.1, we can write  $(1 - \lambda(\theta))^{-1} = c_\beta^{-1} \ell(1/\theta)^{-1} \theta^{-\beta} h(\theta)$  where  $c_\beta = -i \int_0^\infty e^{i\sigma} \sigma^{-\beta} d\sigma$  and  $\lim_{\theta \rightarrow 0} h(\theta) = 1$ . Hence

$$\begin{aligned} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta &= n^{-1} \int_0^a (1 - \lambda(\sigma/n))^{-1} e^{-i\sigma} d\sigma \\ &= c_\beta^{-1} n^{-(1-\beta)} \int_0^a e^{-i\sigma} \sigma^{-\beta} \ell(n/\sigma)^{-1} h(\sigma/n) d\sigma, \end{aligned}$$

so that

$$\ell(n)n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta = c_\beta^{-1} \int_0^a e^{-i\sigma} \sigma^{-\beta} [\ell(n/\sigma)] h(\sigma/n) d\sigma.$$

For fixed  $a$ , it follows from the dominated convergence theorem that

$$\lim_{n \rightarrow \infty} \ell(n)n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta = c_\beta^{-1} \int_0^a e^{-i\sigma} \sigma^{-\beta} d\sigma,$$

and the result follows. ■

**Proof of Theorem 2.1**,  $\beta \in (\frac{1}{2}, 1)$ . By Section 4,  $T_n = \frac{1}{\pi} \operatorname{Re} \int_0^\pi T(\theta) e^{-in\theta} d\theta$ . Let

$$\begin{aligned} D(a, n) &= \int_0^\pi T(\theta) e^{-in\theta} d\theta - \int_0^{a/n} (1 - \lambda(\theta))^{-1} P e^{-in\theta} d\theta \\ &= \int_0^{a/n} \{T(\theta) - (1 - \lambda(\theta))^{-1} P\} e^{-in\theta} d\theta + \int_{a/n}^\pi T(\theta) P e^{-in\theta} d\theta, \end{aligned}$$

so  $D(a, n) \ll a/n + \ell(n)^{-1} n^{-(1-\beta)} a^{-(2\beta'-1)}$  by Lemma 3.1(b) and Lemma 5.1. Hence  $\lim_{a \rightarrow \infty} \lim_{n \rightarrow \infty} \ell(n)n^{1-\beta} D(a, n) = 0$ . By Lemma 5.2,  $\lim_{a \rightarrow \infty} \lim_{n \rightarrow \infty} \ell(n)n^{1-\beta} T_n = \frac{1}{\pi} \operatorname{Re} d'_\beta = d_\beta$ . The result follows since  $T_n$  is independent of  $a$ . ■

## 6 Convergence for $\beta = 1$

In this section, we prove Theorem 2.1 in the case  $\beta = 1$ . There are several differences from the case  $\beta \in (\frac{1}{2}, 1)$ . First,  $T(e^{i\theta}) \notin L^1$ ; instead it is shown below that  $\operatorname{Re} T(e^{i\theta})$  is integrable.

Estimating  $\operatorname{Re}\{(1 - \lambda(z))^{-1}\}$  is slightly easier than in Section 4 but estimating  $\operatorname{Re} T(z)$  is harder since  $\operatorname{Re}\{(1 - \lambda(z))^{-1}(P(z) - P)\}$  is not dominated by  $\operatorname{Re}\{(1 - \lambda(z))^{-1}\}$ . As a consequence,  $\operatorname{Re} T(z) = \operatorname{Re} T(\rho e^{i\theta})$  is not dominated by a single integrable function of  $\theta$ , see Lemma 6.4 below.

We have  $\mu(\varphi > n) = \ell(n)n^{-1}$ , where  $\ell$  is slowly varying and  $\sum \ell(n)n^{-1} = \infty$ . Let  $\tilde{\ell}(x) = m(x) = \sum_{j=1}^{\lfloor x \rfloor} \ell(j)j^{-1}$ . Then  $\tilde{\ell}$  is monotone increasing and  $\lim_{n \rightarrow \infty} \tilde{\ell}(n) = \infty$ . By Proposition 2.6(b),  $\tilde{\ell}$  is slowly varying and  $\ell(n)/\tilde{\ell}(n) \rightarrow 0$  as  $n \rightarrow \infty$ . Up to asymptotic equivalence, we have the alternative definitions  $\tilde{\ell}(x) = \int_1^x \ell(y)y^{-1} dy$  and  $\tilde{\ell}(x) = \int_0^x (1 - G(y)) dy$  where  $G(x) = \mu(\varphi \leq x)$ .

**Proposition 6.1** 
$$\int_0^{1/\theta} \frac{\ell(1/\theta)}{\theta(\tilde{\ell}(1/\theta))^2} d\theta = \int_y^\infty \frac{\ell(x)}{x(\tilde{\ell}(x))^2} dx \sim \frac{1}{\tilde{\ell}(y)}.$$

**Proof** Note that  $-(\tilde{\ell}(x))^{-1}$  is an antiderivative of  $\ell(x)x^{-1}(\tilde{\ell}(x))^{-2}$ . ■

### 6.1 Identification of Fourier coefficients

Write  $z = e^{-u+i\theta}$ ,  $u \in [0, 1]$ ,  $\theta \in [0, \pi]$ . Given a function  $g_\theta(x) \geq 0$  with  $|g_\theta|_\infty \leq C\theta^{1-\epsilon}$  for constants  $C > 0$ ,  $\epsilon \in (0, 1)$ , such that  $x \rightarrow g_\theta(x)(1 - G(x))$  is decreasing for each fixed  $\theta$ , define

$$J_C = \int_0^\infty e^{-ux} \cos \theta x g_\theta(x)(1 - G(x)) dx, \quad J_S = \int_0^\infty e^{-ux} \sin \theta x g_\theta(x)(1 - G(x)) dx.$$

Let  $I_C$  and  $I_S$  be the corresponding integrals in the case  $g_\theta \equiv 1$ .

**Proposition 6.2** *As  $u, \theta \rightarrow 0^+$ ,*

$$\begin{aligned} |I_S| &\ll \theta u^{-1} \ell(1/u), & I_C &= \tilde{\ell}(1/u)(1 + o(1)) + O(\theta u^{-1} \ell(1/u)), \\ |I_S| &\ll \ell(1/\theta), & I_C &= \tilde{\ell}(1/\theta)(1 + o(1)) + O(u\theta^{-1} \ell(1/\theta)), \\ |J_S| &\ll \theta^{2-\epsilon} u^{-1} \ell(1/u), & |J_C| &\ll \theta^{2-\epsilon} u^{-1} \ell(1/u) + \theta^{1-\epsilon} \tilde{\ell}(1/u), \\ |J_S| &\ll \theta^{1-2\epsilon}, & |J_C| &\ll \theta^{1-2\epsilon} + u\theta^{-2\epsilon}. \end{aligned}$$

**Proof** First,

$$\begin{aligned} |J_S| &\ll \theta |g_\theta|_\infty \int_0^\infty e^{-ux} \left| \frac{\sin \theta x}{\theta x} \right| \ell(x) dx \ll \theta |g_\theta|_\infty \int_0^\infty e^{-ux} \ell(x) dx \\ &= \theta |g_\theta|_\infty u^{-1} \ell(1/u) \int_0^\infty e^{-\sigma} \frac{\ell(\sigma/u)}{\ell(1/u)} d\sigma \ll \theta |g_\theta|_\infty u^{-1} \ell(1/u). \end{aligned}$$



This gives the first estimate for  $J_S$  and taking  $g_\theta = 1$  we obtain the first estimate for  $I_S$ . Alternatively, we make the substitution  $\sigma = \theta x$ . Using the oscillation of  $\sin \sigma$  and the fact that  $\sigma \mapsto e^{-u\sigma/\theta} g_\theta(\sigma/\theta)(1 - G(\sigma/\theta))$  is decreasing,

$$\begin{aligned} 0 \leq J_S &= \theta^{-1} \int_0^\infty e^{-u\sigma/\theta} \sin \sigma g_\theta(\sigma/\theta)(1 - G(\sigma/\theta)) d\sigma \\ &\leq \theta^{-1} \int_0^\pi e^{-u\sigma/\theta} \sin \sigma g_\theta(\sigma/\theta)(1 - G(\sigma/\theta)) d\sigma, \end{aligned}$$

and so  $|J_S| \leq |g_\theta|_\infty \int_0^\pi \sin \sigma \ell(\sigma/\theta) \sigma^{-1} d\sigma \ll |g_\theta|_\infty \ell(1/\theta)$ , yielding the remaining estimates for  $I_S$  and  $J_S$ .

In the estimates for  $J_C$  and  $I_C$ , we use the fact that  $\int_0^x (1 - G(x)) dx = \tilde{\ell}(x)(1 + o(1))$ . Note that

$$\left| \int_{1/u}^\infty e^{-ux} \cos \theta x g_\theta(x)(1 - G(x)) dx \right| \leq |g_\theta|_\infty \int_1^\infty e^{-\sigma} \ell(\sigma/u) \sigma^{-1} d\sigma \ll |g_\theta|_\infty \ell(1/u).$$

For the integral over  $[0, 1/u]$ , write  $e^{-ux} \cos \theta x = \{e^{-ux}(\cos \theta x - 1)\} + \{e^{-ux} - 1\} + 1$ . This yields three integrals, the first of which is estimated by  $|g_\theta|_\infty \theta u^{-1} \ell(1/u)$  (like  $J_S$ ) and the second by  $|g_\theta|_\infty \ell(1/u)$ . This leaves  $\int_0^{1/u} g_\theta(x)(1 - G(x)) dx \ll |g_\theta|_\infty \int_0^{1/u} (1 - G(x)) dx = |g_\theta|_\infty \tilde{\ell}(1/u)(1 + o(1))$  completing the first estimate for  $J_C$ . Setting  $g_\theta = 1$  yields the first asymptotic expression for  $I_C$ . The remaining estimate for  $J_C$  is obtained by splitting the range of integration into  $[0, 1/\theta]$  and  $[1/\theta, \infty)$  and combining the above arguments for  $J_C$  (first estimate) and  $J_S$  (second estimate). Again the final expression for  $I_C$  follows by setting  $g_\theta = 1$ .  $\blacksquare$

**Corollary 6.3** *Let  $z = e^{-u+i\theta} \in B_\epsilon(1)$ ,  $\epsilon$  sufficiently small,  $u > 0$ ,  $\theta \geq 0$ . Then*

$$\begin{aligned} |1 - \lambda(e^{-u+i\theta})|^{-1} &\ll \frac{1}{u\tilde{\ell}(1/u)}, \quad \text{for } \theta \in [0, u], \\ |1 - \lambda(e^{-u+i\theta})|^{-1} &\ll \frac{1}{(u + \theta)\tilde{\ell}(1/\theta)}, \quad \text{for } \theta \geq u. \\ |\operatorname{Re}\{(1 - \lambda(e^{-u+i\theta}))^{-1}\}| &\ll \frac{u}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)} + \frac{\theta\ell(1/\theta)}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)^2}, \quad \text{for } \theta \geq u. \end{aligned}$$

**Proof** Recall from the proof of Lemma 4.1 (in particular (4.1)) that  $1 - \lambda(z)$  is the sum of five integrals of the form

$$K = (u - i\theta) \int_0^\infty e^{(-u+i\theta)x} g_z(x)(1 - G(x)) dx,$$

where  $g_z \geq 0$  and either (i)  $g_z \equiv 1$  or by Corollary 2.8 (ii)  $|g_z|_\infty = O(m(1/\theta)\theta)$  as  $\theta \rightarrow 0$ . In case (i),  $K = uI_C + \theta I_S + i(uI_S - \theta I_C)$ . In case (ii),  $K$  consists of terms of the form  $uJ_C$ ,  $uJ_S$ ,  $\theta J_C$ ,  $\theta J_S$ . Hence

$$\operatorname{Re}(1 - \lambda(e^{-u+i\theta})) = uI_C + \theta I_S + E_1, \quad \operatorname{Im} \lambda(e^{-u+i\theta}) = uI_S - \theta I_C + E_2,$$

where  $E_j = O((|u| + |\theta|)(|J_C| + |J_S|))$ ,  $j = 1, 2$ .

For  $\theta \in [0, u]$ , we use the first estimates in Proposition 6.2 to obtain  $\operatorname{Re}(1 - \lambda(e^{-u+i\theta})) \sim u\tilde{\ell}(1/u)$ . Hence  $|1 - \lambda(e^{-u+i\theta})| \geq |\operatorname{Re}(1 - \lambda(e^{-u+i\theta}))| \gg u\tilde{\ell}(1/u)$ .

For  $\theta \geq u$  we use the second estimates in Proposition 6.2 to obtain

$$\operatorname{Re}(1 - \lambda(e^{-u+i\theta})) \sim u\tilde{\ell}(1/\theta) + O(\theta\ell(1/\theta)), \quad \operatorname{Im} \lambda(e^{-u+i\theta}) \sim -\theta\tilde{\ell}(1/\theta).$$

Hence  $|1 - \lambda(e^{-u+i\theta})| \gg (u + \theta)\tilde{\ell}(1/\theta)$ . Finally,

$$\operatorname{Re}\{(1 - \lambda(e^{-u+i\theta}))^{-1}\} = \frac{\operatorname{Re}\{1 - \lambda(e^{-u+i\theta})\}}{|1 - \lambda(e^{-u+i\theta})|^2} \ll \frac{u\tilde{\ell}(1/\theta) + \theta\ell(1/\theta)}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)^2},$$

completing the proof. ■

**Lemma 6.4** For  $u \in [0, 1]$ ,  $\theta \in [0, \pi]$ , we have  $\operatorname{Re} T(e^{-u+i\theta}) \ll h_u(\theta) + g(\theta)$  where

$$h_u(\theta) = \frac{u\tilde{\ell}(1/u)}{u + \theta} + \frac{1}{u\tilde{\ell}(1/u)} 1_{[0, u]}(\theta) + \frac{1}{\tilde{\ell}(1/\theta)} \frac{u}{u^2 + \theta^2}, \quad g(\theta) = \frac{\ell(1/\theta)}{\tilde{\ell}(1/\theta)^2} \frac{1}{\theta}.$$

**Proof** By Proposition 2.9, for  $z = e^{-u+i\theta} \in B_\epsilon(1)$ ,  $\epsilon$  sufficiently small,

$$T(z) = (1 - \lambda(z))^{-1}P + (1 - \lambda(z))^{-1}(P(z) - P) + O(1).$$

By Corollary 6.3,

$$\begin{aligned} \operatorname{Re}\{(1 - \lambda(z))^{-1}\} &\ll \frac{1}{u\tilde{\ell}(1/u)} 1_{[0, u]} + \left\{ \frac{u}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)} + \frac{\theta\ell(1/\theta)}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)^2} \right\} 1_{[u, \epsilon]} \\ &\leq \frac{1}{u\tilde{\ell}(1/u)} 1_{[0, u]} + \frac{u}{(u^2 + \theta^2)\tilde{\ell}(1/\theta)} + \frac{\ell(1/\theta)}{\theta\tilde{\ell}(1/\theta)^2}. \end{aligned}$$

By Corollary 2.8,  $P(e^{-u+i\theta}) - P(e^{-u}) \ll \tilde{\ell}(1/\theta)\theta$  uniformly in  $u$ , and  $P(e^{-u}) - P(1) \ll \tilde{\ell}(1/u)u$ . Combining this with the estimates for  $(1 - \lambda(e^{-u+i\theta}))^{-1}$ ,

$$\begin{aligned} (1 - \lambda(z))^{-1}(P(z) - P(1)) &\ll \left( \frac{1}{u\tilde{\ell}(1/u)} 1_{[0, u]} + \frac{1}{(u + \theta)\tilde{\ell}(1/\theta)} 1_{[u, \epsilon]} \right) (\theta\tilde{\ell}(1/\theta) + u\tilde{\ell}(1/u)) \\ &\ll 1 + \frac{1}{u\tilde{\ell}(1/u)} 1_{[0, u]} + \frac{u\tilde{\ell}(1/u)}{u + \theta}. \end{aligned}$$

This proves the result. ■

**Remark 6.5** By similar but much simpler calculations, we obtain the estimates  $|\operatorname{Re}\{(1 - \lambda(e^{i\theta}))^{-1}\}| \ll g(\theta)$  for  $\theta \in (0, \epsilon)$  and  $\operatorname{Re} T(e^{i\theta}) \ll g(\theta)$  for  $\theta \in (0, \pi]$ .

**Corollary 6.6** For  $n \geq 1$ ,  $\lim_{u \rightarrow 0} \int_0^\pi \cos n\theta \operatorname{Re} T(e^{-u+i\theta}) d\theta = \int_0^\pi \cos n\theta \operatorname{Re} T(e^{i\theta}) d\theta$ .

**Proof** The function  $g(\theta) = \frac{\ell(1/\theta)}{\tilde{\ell}(1/\theta)^2} \frac{1}{\theta}$  lies in  $L^1$  by Proposition 6.1. Note that  $\operatorname{Re} T(e^{-u+i\theta}) \rightarrow \operatorname{Re} T(e^{i\theta})$  and  $h_u(\theta) \rightarrow 0$  pointwise (for all  $\theta \neq 0$ ). We claim that  $\int_0^\pi h_u(\theta) d\theta \rightarrow 0$ . The result then follows from the dominated convergence theorem (more precisely the extended version stated in [35, p. 92]).

The claim is easy to check for the first two terms in  $h_u$ . For the third term  $k_u(\theta) = \frac{1}{\tilde{\ell}(1/\theta)} \frac{u}{u^2 + \theta^2}$ , we compute for  $b \in (0, \pi)$  that

$$\begin{aligned} \int_0^\pi k_u(\theta) d\theta &= \int_0^b k_u(\theta) d\theta + \int_b^\pi k_u(\theta) d\theta \ll \frac{1}{\tilde{\ell}(1/b)} u^{-1} b + u \int_b^\pi \theta^{-2} d\theta \\ &\ll \frac{1}{\tilde{\ell}(1/b)} u^{-1} b + u b^{-1}, \end{aligned}$$

where the implied constant is independent of  $b$  and  $u$ . Define  $b = b(u)$  such that  $u = b(\tilde{\ell}(1/b))^{-1/2}$ . In particular,  $b \rightarrow 0$  as  $u \rightarrow 0$  and so  $\int_0^\pi k_u(\theta) d\theta \ll (\tilde{\ell}(1/b))^{-1/2} \rightarrow 0$  as required.  $\blacksquare$

**Corollary 6.7**  $\operatorname{Re} T \in L^1$  and  $T_n = \frac{2}{\pi} \int_0^\pi \cos n\theta \operatorname{Re} T(e^{i\theta}) d\theta$  for all  $n \geq 1$ .

**Proof** The function  $\theta \mapsto T(\rho e^{i\theta})$  is integrable for each fixed  $\rho < 1$ . Moreover, the power series for  $T(z)$  is uniformly convergent on compact subsets of  $\mathbb{D}$ , so we obtain

$$\int_0^\pi \cos n\theta \operatorname{Re} T(\rho e^{i\theta}) d\theta = \sum_{j=0}^\infty T_j \rho^j \int_0^\pi \cos n\theta \cos j\theta d\theta = \frac{\pi}{2} T_n \rho^n.$$

By Corollary 6.6,  $T_n = \frac{2}{\pi} \rho^{-n} \int_0^\pi \cos n\theta \operatorname{Re} T(\rho e^{i\theta}) d\theta \rightarrow \frac{2}{\pi} \int_0^\pi \cos n\theta \operatorname{Re} T(e^{i\theta}) d\theta$ , as  $\rho = e^{-u} \rightarrow 1$ .  $\blacksquare$

## 6.2 Asymptotics of $T_n$

The calculations in this subsection are restricted to the unit circle, so we revert to writing  $T(\theta)$  instead of  $T(e^{i\theta})$  and so on. First we determine the asymptotics of  $\operatorname{Re}\{(1 - \lambda(\theta))^{-1}\}$  (see also [4]).

**Lemma 6.8**  $\operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} = \frac{\pi}{2} g(\theta)(1 + o(1))$  as  $\theta \rightarrow 0^+$ , where  $g(\theta) = \frac{\ell(1/\theta)}{\theta(\tilde{\ell}(1/\theta))^2}$ .

**Proof** By Remark 6.5,  $\operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} \ll g(\theta)$ . We claim that  $I_S \sim \frac{\pi}{2} \ell(1/\theta)$  from which the result follows easily.

Let  $M \geq 3\pi$ . Since  $\sigma \mapsto \ell(\sigma)/\sigma$  is decreasing, we have the oscillatory integral estimate

$$\frac{1}{\ell(1/\theta)} I_S = \int_0^\infty \frac{\sin \sigma}{\sigma} \frac{\ell(\sigma/\theta)}{\ell(1/\theta)} d\sigma = \int_0^M \frac{\sin \sigma}{\sigma} \frac{\ell(\sigma/\theta)}{\ell(1/\theta)} d\sigma + F_M,$$

where

$$|F_M| \leq 2 \sup_{\sigma \in [M-2\pi, M+2\pi]} \frac{\ell(\sigma/\theta)}{\sigma \ell(1/\theta)}.$$

By Potter's bounds, for any  $\delta > 0$ ,  $F_M = O(1/M^{1-\delta})$ . Hence,  $\lim_{\theta \rightarrow 0} \ell(1/\theta)^{-1} I_S = \int_0^M \frac{\sin \sigma}{\sigma} d\sigma + O(1/M^{1-\delta})$ . Let  $M \rightarrow \infty$  to verify the claim.  $\blacksquare$

**Corollary 6.9** *Let  $a > 0$ . Then  $\lim_{n \rightarrow \infty} \tilde{\ell}(n) \int_0^{a/n} \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta = \frac{\pi}{2}$ .*

**Proof** By Lemma 6.8, we can write  $\operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} = \frac{\pi}{2} g(\theta)(1 + h(\theta))$  where  $h(\theta) = o(1)$  as  $\theta \rightarrow 0^+$ . Let  $H(n) = \sup_{\theta \in [0, a/n]} |h(\theta)|$ , so  $H(n) = o(1)$  as  $n \rightarrow \infty$ . Then

$$\int_0^{a/n} \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta = \frac{\pi}{2} \int_0^{a/n} g(\theta) d\theta + O\left(H(n) \int_0^{a/n} g(\theta) d\theta\right).$$

By Proposition 6.1,  $\int_0^{a/n} g(\theta) d\theta = \tilde{\ell}(n/a)^{-1}$ . Hence

$$\tilde{\ell}(n) \int_0^{a/n} \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta = \frac{\pi}{2} \frac{\tilde{\ell}(n)}{\tilde{\ell}(n/a)} (1 + o(1)) \rightarrow \frac{\pi}{2},$$

as  $n \rightarrow \infty$ .  $\blacksquare$

**Proof of Theorem 2.1**,  $\beta = 1$ . By Remark 6.5 and Proposition 2.6(b),  $T(\theta) \ll (\theta \tilde{\ell}(1/\theta))^{-1}$ . Let  $\delta > 0$ . By the argument in the proof of Lemma 5.1, we obtain

$$\tilde{\ell}(n) \int_{a/n}^{\pi} \cos n\theta T(\theta) d\theta \ll a^{-(1-\delta)}, \quad (6.1)$$

for  $a \in [1, n]$ ,  $n \geq 1$ . Also, we have

$$\tilde{\ell}(n) \int_0^{a/n} \cos n\theta T(\theta) d\theta = \tilde{\ell}(n) \int_0^{a/n} \cos n\theta (1 - \lambda(\theta))^{-1} d\theta P + O(an^{-(1-\delta)}). \quad (6.2)$$

By Lemma 6.8,

$$\begin{aligned} & \tilde{\ell}(n) \int_0^{a/n} (\cos n\theta - 1) \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta \\ & \ll \tilde{\ell}(n) \int_0^a (\cos \sigma - 1) \ell(n/\sigma) (\tilde{\ell}(n/\sigma))^{-2} \sigma^{-1} d\sigma \ll \ell(n) (\tilde{\ell}(n))^{-1} a. \end{aligned} \quad (6.3)$$

Combining estimates (6.1), (6.2) and (6.3) with Corollary 6.9, we obtain

$$\lim_{a \rightarrow \infty} \lim_{n \rightarrow \infty} \tilde{\ell}(n) \int_0^{\pi} \cos n\theta \operatorname{Re} T(\theta) d\theta = \frac{\pi}{2} P,$$

and hence  $\tilde{\ell}(n) T_n \rightarrow P$  by Corollary 6.7.  $\blacksquare$

## 7 Pointwise dual ergodicity

In this section, we give an elementary proof of pointwise dual ergodicity for the class of systems under consideration for all  $\beta \in (0, 1]$ . We assume our general framework from Section 2, except that we do not require H2(ii).

**Proposition 7.1**  $T(s) \sim \begin{cases} \tilde{\ell}(\frac{1}{1-s})^{-1}(1-s)^{-1}P, & \beta = 1, \\ \Gamma(1-\beta)^{-1}\ell(\frac{1}{1-s})^{-1}(1-s)^{-\beta}P, & \beta \in (0, 1), \end{cases}$  as  $s \rightarrow 1^-$ .

**Proof** This is similar to the proof of Lemma 3.1, but much simpler since the integrals are absolutely convergent. By Proposition 2.9, for  $s \in (1-\epsilon, 1]$ ,

$$T(s) = (1 - \lambda(s))^{-1}P + (1 - \lambda(s))^{-1}(P(s) - P) + O(1).$$

By Corollary 2.8,  $P(s) - P \ll m(\frac{1}{1-s})(1-s)^\beta$ , so it suffices to establish the desired asymptotic expression for  $(1 - \lambda(s))^{-1}$ .

Setting  $s = e^{-u}$  we have  $\lambda(s) = 1 + \int_0^\infty (e^{-ux} - 1)\hat{v}_u(x)dG(x)$ , where  $G(x) = \mu(\varphi \leq x)$  and  $|\hat{v}_u - 1|_\infty = o(1)$  as  $u \rightarrow 0$ . Writing  $d\hat{G}_u = v_u dG$  and integrating by parts,

$$\lambda(s) = 1 + \int_0^\infty (e^{-ux} - 1)d\hat{G}_u(x) = 1 - u \int_0^\infty e^{-ux}g_u(x)(1 - G(x)) dx, \quad (7.1)$$

where  $|g_u(x) - 1|_\infty = o(1)$  as  $u \rightarrow 0$ .

If  $\beta \in (0, 1)$ , then

$$\lambda(s) = 1 - \ell(1/u)u^\beta \int_0^\infty e^{-\sigma}g_u(\sigma/u)\{\ell(\sigma/u)\ell(1/u)^{-1}\}\sigma^{-\beta} du.$$

By the dominated convergence theorem,  $\lambda(s) = 1 - \Gamma(1-\beta)\ell(1/u)u^\beta(1+o(1))$ . The result follows since  $u = -\log s = 1 - s + O((1-s)^2)$ .

If  $\beta = 1$ , then picking up from (7.1),

$$\begin{aligned} \int_0^\infty e^{-ux}g_u(x)(1 - G(x)) dx &= \int_0^{1/u} g_u(x)(1 - G(x)) dx \\ &+ \int_0^{1/u} (e^{-ux} - 1)g_u(x)(1 - G(x)) dx + \int_{1/u}^\infty e^{-ux}g_u(x)(1 - G(x)) dx. \end{aligned}$$

The last two integrals are  $O(\ell(1/u))$ , and  $\int_0^{1/u} g_u(x)(1 - G(x)) dx = (1+o(1)) \int_0^{1/u} (1 - G(x)) dx \sim \tilde{\ell}(1/u)$  by definition of  $\tilde{\ell}$ .  $\blacksquare$

**Theorem 7.2** Let  $v \in L^1(X)$  and  $\beta \in (0, 1]$ . Then

$$\lim_{n \rightarrow \infty} m(n)n^{-\beta} \sum_{j=1}^n L^j v = \beta^{-1}d_\beta \int_X v d\mu, \text{ almost everywhere on } X.$$

**Proof** By [3, Proposition 3.7.6] (see also the proof of Proposition 10.2), it suffices to prove pointwise dual ergodicity on the tower  $\Delta$  defined in Section 2.4. Let  $L_\Delta : L^1(\Delta) \rightarrow L^1(\Delta)$  denote the transfer operator on  $\Delta$ . Note that  $T_n v = 1_Y L_\Delta^n(1_Y v)$  coincides with our usual  $T_n$ . By the Hurewicz ergodic theorem [26], it is enough to prove that  $m(n)n^{-\beta} \sum_{j=0}^{n-1} L_\Delta^j v \rightarrow \beta^{-1} d_\beta \int_\Delta v d\mu$  almost everywhere on  $\Delta$  for the particular choice  $v = 1_Y$ .

Let  $y \in Y$ . For  $\beta \in (0, 1)$ , Proposition 7.1 gives  $(T(s)v)(y) \sim \Gamma(1-\beta)\ell(\frac{1}{1-s})^{-1}(1-s)^{-\beta} \int_Y v d\mu$  as  $s \rightarrow 1^-$ . By (the discrete version of) the Karamata Tauberian Theorem [27] [16, p. 445], [41, Proposition 4.2], it follows that  $\sum_{j=1}^n (T_j v)(y) \sim \beta^{-1} d_\beta \ell(n)^{-1} n^\beta \int_Y v d\mu$  as  $n \rightarrow \infty$ . Similarly for  $\beta = 1$ .

Finally, let  $p = (y, j)$  be a general point in  $\Delta$ . Then  $(L_\Delta^n v)(p) = (L_\Delta^{n-j} v)(y, 0) = (T_{n-j} v)(y)$  for all  $n > j$ .  $\blacksquare$

An immediate consequence is the Darling-Kac law [10]. Recall that a random variable  $\mathcal{M}_\beta$  on  $(0, \infty)$  has the *normalised Mittag-Leffler distribution of order  $\beta$*  if  $E(e^{z\mathcal{M}_\beta}) = \sum_{p=0}^\infty \Gamma(1+\beta)^p z^p / \Gamma(1+p\beta)$  for all  $z \in \mathbb{C}$ .

**Corollary 7.3** *Let  $v \in L^1(X)$ ,  $v \geq 0$ ,  $\int_X v d\mu = 1$ , and let  $\beta \in (0, 1]$ . Then*

$$m(n)n^{-\beta} \sum_{j=1}^n v \circ f^j \rightarrow_d \mathcal{M}_\beta \text{ as } n \rightarrow \infty.$$

*The convergence is in the sense of strong distributional convergence: convergence in distribution under any probability measure absolutely continuous w.r.t.  $\mu$ .*

**Proof** This follows from Theorem 7.2 by Aaronson [1], [3, Corollary 3.7.3].  $\blacksquare$

To conclude the section, we mention a simple consequence of Theorem 2.1 which gives uniform convergence on  $Y$  in the pointwise dual ergodic theorem for  $\beta > \frac{1}{2}$  for sufficiently regular observables.

**Proposition 7.4** *If  $\beta \in (\frac{1}{2}, 1]$ , then  $\lim_{n \rightarrow \infty} m(n)n^{-\beta} \sum_{j=1}^n T_j = \beta^{-1} d_\beta P$ .*

**Proof** By Theorem 2.1,  $T_n = m(n)^{-1} n^{-(1-\beta)} d_\beta P + S_n$  where  $\|S_n\| = o(m(n)^{-1} n^{-(1-\beta)})$ . Hence

$$m(n)n^{-\beta} \sum_{j=1}^n T_j = m(n)n^{-\beta} \sum_{j=1}^n m(j)^{-1} j^{-(1-\beta)} d_\beta P + m(n)n^{-\beta} \sum_{j=1}^n S_j. \quad (7.2)$$

By Proposition 2.6(a),  $\sum_{j=1}^n m(j)^{-1} j^{-(1-\beta)} \sim \beta^{-1} m(n)^{-1} n^\beta$ , so the first term on the RHS of (7.2) converges to the desired limit  $\beta^{-1} d_\beta P$ .

Let  $\delta > 0$ , and choose  $n_0$  such that  $\|S_n\| \leq \delta m(n)^{-1} n^{-(1-\beta)}$  for  $n > n_0$ . Then  $\sum_{j=1}^n \|S_j\| \leq \sum_{j=1}^{n_0} \|S_j\| + \sum_{j=n_0+1}^n \delta m(j)^{-1} j^{-(1-\beta)}$ . Applying Proposition 2.6(a) once more, we obtain  $\limsup_{n \rightarrow \infty} m(n)n^{-\beta} \sum_{j=1}^n \|S_j\| \leq \beta^{-1} \delta$ . Since  $\delta > 0$  is arbitrary, the second term on the RHS of (7.2) converges to zero.  $\blacksquare$

## 8 Results for $\beta \in (0, \frac{1}{2}]$

In this section, we prove Theorems 2.2 and 2.3.

**Proof of Theorem 2.2** If  $\beta \in (0, \frac{1}{2}]$ , then the proof of Theorem 2.1 breaks down only in the estimation of  $I_3$  in Lemma 5.1.

(a) When  $\beta = \frac{1}{2}$ , it is evident from the proof of Lemma 5.1 that  $I_3 \ll \ell(n)n^{-\frac{1}{2}} \int_{1/n}^{\pi} \ell(1/\theta)^{-2}\theta^{-1} d\theta$ . The remaining estimates are  $O(\ell(n)^{-1}n^{-\frac{1}{2}})$  as before. By Proposition 2.6(b),  $\ell(n)^2 \int_{1/n}^{\pi} \ell(1/\theta)^{-2}\theta^{-1} d\theta \rightarrow \infty$  as  $n \rightarrow \infty$ . Hence the estimate for  $I_3$  is the dominant one.

(b) For  $\beta \in (0, \frac{1}{2})$ ,  $I_3 \ll \ell(n)n^{-\beta} \int_0^{\pi} \ell(1/\theta)^{-2}\theta^{-2\beta} d\theta \ll \ell(n)n^{-\beta}$ . The remaining estimates are  $O(\ell(n)^{-1}n^{-(1-\beta)})$  as before.

(c) If  $Pv = 0$ , then  $\|T(\theta)v\| \ll \|v\|$ . Hence the resolvent identity

$$\{T(\theta) - T(\theta - \pi/n)\}v = T(\theta)(R(\theta) - R(\theta - \pi/n))T(\theta - \pi/n)v$$

yields  $\|\{T(\theta) - T(\theta - \pi/n)\}v\| \ll \ell(1/\theta)^{-1}\theta^{-\beta}\ell(n)n^{-\beta}$ . It follows that

$$2T_nv = \int_0^{2\pi} \{T(\theta) - T(\theta - \pi/n)\}e^{-in\theta} d\theta \ll \ell(n)n^{-\beta}\|v\|,$$

as required. ■

Next, we establish the lower bound in Theorem 2.3(b).

**Proposition 8.1** *If  $\beta \in (0, 1)$ , then  $\liminf_{n \rightarrow \infty} \ell(n)n^{1-\beta}T_nv \geq d_\beta \int_Y v d\mu$  pointwise on  $Y$  for all  $v \geq 0$ .*

**Proof** For any  $m \geq 1$ , we can write

$$T = (I - R)^{-1} = I + R + \cdots + R^{m-1} + T^{(m)}, \quad T^{(m)} = R^m(I - R)^{-1}.$$

Since  $R$  is a positive operator, we deduce that  $(T_nv)(y) \geq (T_n^{(m)}v)(y)$  for all  $v \geq 0$ ,  $y \in Y$ ,  $m \geq 1$ . Choosing  $m = b_n \sim b\ell(n)^{-1}n^\beta$ ,  $b > 0$ , as in [17, Theorem 3.6.1], we obtain  $\ell(n)n^{1-\beta}T_n^{(b_n)} \sim d_b P$ , where  $d_b \rightarrow d_\beta$  as  $b \rightarrow 0$ , and the result follows. ■

The following result is well-known (see [7, Theorem 2.9.1], [17]) but stated in a slightly different form, so we provide the proof for completeness.

**Proposition 8.2** *Let  $f_n$  be a sequence in  $\mathbb{R}$  and let  $A \in \mathbb{R}$ . Suppose that  $\beta \in (0, 1)$ , that  $\ell(n)$  is slowly varying, and that*

$$(a) \liminf_{n \rightarrow \infty} \ell(n)n^{1-\beta}f_n \geq A,$$

$$(b) \lim_{n \rightarrow \infty} \ell(n)n^{-\beta} \sum_{j=1}^n f_j = \beta^{-1}A.$$

Then there exists a set  $E$  of density zero such that  $\lim_{n \rightarrow \infty, n \notin E} \ell(n)n^{1-\beta} f_n = A$ .  
 In particular,  $\liminf_{n \rightarrow \infty} \ell(n)n^{1-\beta} f_n = A$ .

**Proof** Our proof is modelled on [32, p. 65, Lemma 6.2].

By Proposition 2.6(a),  $\sum_{j=1}^n \ell(j)^{-1} j^{-(1-\beta)} \sim \beta^{-1} \ell(n)^{-1} n^\beta$ . Let  $\hat{f}_n = f_n - \ell(n)^{-1} n^{(\beta-1)} A$ . Then (b) is equivalent to  $\lim_{n \rightarrow \infty} \ell(n)n^{-\beta} \sum_{j=1}^n \hat{f}_j = 0$ . Hence we may suppose without loss that  $A = 0$ . In addition, there is a monotone increasing function  $g(n)$  such that  $\ell(n)n^{1-\beta} \sim g(n)$  (see for example [7, Theorem 1.5.3]). Hence we may suppose that  $\ell(n)n^{1-\beta}$  is increasing.

Define the nested sequence of sets  $E_q = \{n \geq 1 : \ell(n)n^{1-\beta} f_n > 1/q\}$ . We claim that each  $E_q$  has density zero. Let  $\delta > 0$ . By (a), there exists  $n_0 \geq 1$  such that  $\ell(n)n^{1-\beta} f_n > -\delta$  for all  $n \geq n_0$ . Hence

$$\begin{aligned} \frac{1}{n} \sum_{j=1}^n 1_{E_q}(j) &\leq \ell(n)n^{-\beta} \sum_{j=1}^n \{\ell(j)j^{(1-\beta)}\}^{-1} 1_{E_q}(j) \leq q\ell(n)n^{-\beta} \sum_{j=1}^n f_j 1_{E_q}(j) \\ &= q\ell(n)n^{-\beta} \left( \sum_{j=1}^n f_j - \sum_{j=1}^{n_0} f_j 1_{E_q^c}(j) - \sum_{n_0+1}^n f_j 1_{E_q^c}(j) \right) \\ &\leq q\ell(n)n^{-\beta} \sum_{j=1}^n f_j + q\ell(n)n^{-\beta} \sum_{j=1}^{n_0} |f_j| + q\ell(n)n^{-\beta} \sum_{n_0+1}^n \ell(j)^{-1} j^{\beta-1} \delta \\ &= q\ell(n)n^{-\beta} \sum_{j=1}^n f_j + O(\ell(n)n^{-\beta}) + O(\delta). \end{aligned}$$

By (b),  $\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n 1_{E_q}(j) = O(\delta)$  and the claim follows since  $\delta$  is arbitrary.

By the claim, there exist  $1 = i_0 < i_1 < i_2 < \dots$  such that  $\frac{1}{n} \sum_{j=1}^n 1_{E_q}(j) < 1/q$  for  $n \geq i_{q-1}$ ,  $q \geq 2$ . Let  $E = \bigcup_{q=1}^{\infty} E_q \cap (i_{q-1}, i_q)$ . If  $n \in E$  and  $n \leq i_q$ , then  $n \in E_q$ . Hence for  $i_{q-1} \leq n \leq i_q$ , we have  $\frac{1}{n} \sum_{j=1}^n 1_E(j) \leq \frac{1}{n} \sum_{j=1}^n 1_{E_q}(j) \leq 1/q$ , verifying that  $E$  has density zero.

On the other hand, if  $n \notin E$  and  $i_{q-1} < n < i_q$ , then  $n \notin E_q$  and so  $\ell(n)n^{1-\beta} f_n \leq 1/q$ . Hence  $\limsup_{n \rightarrow \infty, n \notin E} \ell(n)n^{1-\beta} f_n \leq 0$ . Combined with assumption (a), we deduce that  $\lim_{n \rightarrow \infty, n \notin E} \ell(n)n^{1-\beta} f_n = 0$  and the last statement follows immediately.  $\blacksquare$

**Proof of Theorem 2.3** Part (b) is stated for  $v \geq 0$  and in part (a) we can break  $v$  into positive and negative parts. Hence without loss we may suppose that  $v \geq 0$ .

By Proposition 8.1 and Theorem 7.2, we have verified the hypotheses of Proposition 8.2 with  $f_n = (T_n v)(y)$  and  $A = d_\beta \int_Y v d\mu$ . The result is immediate.  $\blacksquare$



## 9 Second order asymptotics

In this section we prove results on second order asymptotics and higher order asymptotic expansions under assumptions on the asymptotics of  $\mu(\varphi > n)$ . Throughout, we suppose that  $\ell(n)$  is asymptotically constant (and that  $\beta > \frac{1}{2}$ ). In Subsection 9.1, we consider the case when  $\beta \in (\frac{1}{2}, 1)$ . The case  $\beta = 1$  is covered in Subsection 9.2. Error terms in the Dynkin-Lamperti arcsine law are obtained in Subsection 9.3.

### 9.1 Second order asymptotics for $\beta \in (\frac{1}{2}, 1)$

We assume that  $\mu(\varphi > n) = c(n^{-\beta} + H(n))$ , where  $H(n) = O(n^{-2\beta})$  and  $c > 0$ . (It is easy to relax this to the more general hypothesis that  $H(n) = O(n^{-q})$ ,  $q > 1$ . However the formulas become more complicated and our assumption is satisfied by (1.1).)

Recall that  $c_H = \int_0^\infty H_1(x) dx$  where  $H_1(x) = [x]^{-\beta} - x^{-\beta} + H([x])$ . Define  $\xi_p^\pm = \int_0^\infty e^{\pm i\sigma} \sigma^{-p} d\sigma$ ,  $0 < p < 1$ , so  $c_\beta = -i\xi_\beta^+$ , and recall that  $e_0 = ic_H/c_\beta$ .

Set  $d'_{\beta,j} = e_0^j \xi_{(j+1)\beta-j}^- / c_\beta = ie_0^j \xi_{(j+1)\beta-j}^- / \xi_\beta^+$  and  $d_{\beta,j} = \frac{1}{\pi} \operatorname{Re} d'_{\beta,j}$ .

We note that  $d_{\beta,0} = d_\beta = \frac{1}{\pi} \sin \beta\pi > 0$ , and that either  $d_{\beta,j} = 0$  for all  $j \geq 1$  or  $d_{\beta,j} \neq 0$  for all  $j \geq 1$ . Moreover, the latter situation is typical.

**Theorem 9.1** *Suppose that  $\beta \in (\frac{1}{2}, 1)$  and that  $\mu(\varphi > n) = c(n^{-\beta} + H(n))$ , where  $H(n) = O(n^{-2\beta})$  and  $c > 0$ . Let  $\gamma = \min\{1 - \beta, \beta - \frac{1}{2}\}$ . Then*

$$n^{1-\beta} T_n = c^{-1} d_\beta P + O(n^{-\gamma}).$$

Moreover, if  $\beta \in (\frac{3}{4}, 1)$ , then  $\lim_{n \rightarrow \infty} n^{1-\beta} \{n^{1-\beta} T_n - c^{-1} d_\beta P\} = d_{\beta,1} P$ .

**Remark 9.2** For  $\beta$  close to 1, we obtain higher order asymptotic expansions. There exist constants  $d_{\beta,j} \in \mathbb{R}$ ,  $j \geq 0$  with  $d_{\beta,0} = d_\beta$ , such that for each  $q = 0, 1, 2, \dots$ ,

$$n^{1-\beta} T_n = c^{-1} \left\{ \sum_{j=0}^q d_{\beta,j} n^{-j(1-\beta)} + O(n^{-(q+1)(1-\beta)}) \right\} P + O(n^{-(\beta-\frac{1}{2})}).$$

Thus,  $n^{1-\beta} T_n = \begin{cases} c^{-1} d_\beta P + O(n^{-(\beta-\frac{1}{2})}), & \beta \in (\frac{1}{2}, \frac{3}{4}] \\ c^{-1} d_\beta P + c^{-1} d_{\beta,1} n^{-(1-\beta)} P + O(n^{-(\beta-\frac{1}{2})}), & \beta \in (\frac{3}{4}, \frac{5}{6}] \end{cases}$  and so on.

**Corollary 9.3**  $n^{-\beta} \sum_{j=1}^n T_j = c^{-1} \beta^{-1} d_\beta P + O(n^{-\gamma})$  uniformly on  $Y$ .

**Proof** Specialising the proof of Proposition 7.4, we have

$$n^{-\beta} \sum_{j=1}^n T_j = n^{-\beta} \sum_{j=1}^n j^{-(1-\beta)} c^{-1} d_\beta P + n^{-\beta} \sum_{j=1}^n S_j,$$

where  $S_j = O(j^{-(1-\beta+\gamma)})$ . Now  $\sum_{j=1}^n j^{-(1-\beta)} = \int_1^n x^{-(1-\beta)} dx + O(1) = \beta^{-1}n^\beta + O(1)$ , and  $\sum_{j=1}^n S_j = O(n^{\beta-\gamma})$ .  $\blacksquare$

In the remainder of this subsection, we prove Theorem 9.1.

**Proposition 9.4**  $(1 - \lambda(\theta))^{-1} = c^{-1}c_\beta^{-1} \sum_j e_0^j \theta^{-((j+1)\beta-j)} + O(1)$ , where the sum is over those  $j \geq 0$  with  $(j+1)\beta - j > 0$ .

**Proof** By Lemma 3.2,  $1 - \lambda(\theta) = cc_\beta\theta^\beta(1 - e_0\theta^{1-\beta} + O(\theta^\beta))$ . Now invert and note that  $(1 - e_0\theta^{1-\beta} + O(\theta^\beta))^{-1} = \sum_{j \geq 0} e_0^j \theta^{j(1-\beta)} + O(\theta^\beta)$ .  $\blacksquare$

**Proposition 9.5** Let  $n \geq 1$ ,  $0 < a < \epsilon n$ . Then

$$\begin{aligned} n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta \\ = c^{-1} \sum_j d'_{\beta,j} n^{-j(1-\beta)} + O\left(\sum_j n^{-j(1-\beta)} a^{-((j+1)\beta-j)}\right) + O(an^{-\beta}), \end{aligned}$$

and the sums are over those  $j \geq 0$  with  $(j+1)\beta - j > 0$ .

**Proof** By Proposition 9.4,

$$\begin{aligned} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta &= c^{-1}c_\beta^{-1} \sum_j e_0^j \int_0^{a/n} \theta^{-((j+1)\beta-j)} e^{-in\theta} d\theta + O(a/n) \\ &= c^{-1}c_\beta^{-1} \sum_j e_0^j n^{-j(1-\beta)} \int_0^a \sigma^{-((j+1)\beta-j)} e^{-i\sigma} d\sigma + O(a/n). \end{aligned}$$

Hence

$$\begin{aligned} n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta \\ = c^{-1} \sum_j d'_{\beta,j} n^{-j(1-\beta)} - c^{-1}c_\beta^{-1} \sum_j e_0^j n^{-j(1-\beta)} \int_a^\infty \sigma^{-((j+1)\beta-j)} e^{-i\sigma} d\sigma + O(an^{-\beta}), \end{aligned}$$

yielding the required result.  $\blacksquare$

**Proof of Theorem 9.1** By Lemma 5.1,  $n^{1-\beta} \int_{a/n}^\pi T(\theta) e^{-in\theta} d\theta \ll a^{-(2\beta-1)}$ . For  $\theta \in [0, a/n] \subset [0, \epsilon]$ ,  $T(\theta) = (1 - \lambda(\theta))^{-1}P + O(1)$  by Lemma 3.1(b). Hence,

$$n^{1-\beta} \int_0^\pi T(\theta) e^{-in\theta} d\theta = n^{1-\beta} \int_0^{a/n} (1 - \lambda(\theta))^{-1} e^{-in\theta} d\theta P + O(an^{-\beta}) + O(a^{-(2\beta-1)}).$$

Taking  $a = n^{1/2}$  we obtain the error term  $O(n^{-(\beta-\frac{1}{2})})$  and the result follows from Proposition 9.5 and Corollary 4.2.  $\blacksquare$

## 9.2 Second order asymptotics for $\beta = 1$

**Theorem 9.6** *Suppose that  $\mu(\varphi > n) = c(n^{-1} + H(n))$  where  $c > 0$  and  $H(n) = O(n^{-q})$ ,  $q > 1$ . Let  $H_1(x) = [x]^{-1} - x^{-1} + H([x])$ ,  $x \geq 1$  and  $H_1(x) = \frac{1}{c}$ ,  $x \in [0, 1)$ . Then*

$$(\log n)T_n = c^{-1}\left\{1 - \int_0^\infty H_1(x) dx (\log n)^{-1} + O((\log n)^{-2})\right\}P + O((\log n)^{\frac{1}{2}}n^{-\frac{1}{2}}).$$

**Corollary 9.7**  $(\log n)n^{-1} \sum_{j=1}^n T_j = c^{-1}P + O((\log n)^{-1})$  uniformly on  $Y$ .

**Proof** As in the proof of Corollary 9.3, we have

$$(\log n)n^{-1} \sum_{j=1}^n T_j = (\log n)n^{-1} \sum_{j=1}^n (\log j)^{-1}c^{-1}P + (\log n)n^{-1} \sum_{j=1}^n S_j,$$

where  $\|S_j\| = O((\log j)^{-2})$ . Integration by parts yields  $\sum_{j=1}^n (\log j)^{-1} = n(\log n)^{-1} + \int_2^n (\log x)^{-2} dx + O(1) = n(\log n)^{-1} + O(n(\log n)^{-2})$  while  $\sum_{j=1}^n S_j \ll n(\log n)^{-2}$ . ■

In the remainder of this subsection, we prove Theorem 9.6.

**Proposition 9.8** *Let  $c_H = \int_0^1 (\cos \sigma - 1)\sigma^{-1} d\sigma + \int_1^\infty \cos \sigma \sigma^{-1} d\sigma + \int_0^\infty H_1(x) dx$ . Then*

$$\operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} = c^{-1}\frac{\pi}{2}\theta^{-1}(\log \frac{1}{\theta})^{-2} - c^{-1}c_H\pi\theta^{-1}(\log \frac{1}{\theta})^{-3} + O(\theta^{-1}(\log \frac{1}{\theta})^{-4}).$$

**Proof** Without loss of generality, we may suppose that  $q \in (1, 2)$ . Recall that  $G(x) \equiv 0$  for  $x \in [0, 1)$  and  $1 - G(x) = c(x^{-1} + H_1(x))$  for  $x \geq 1$  where  $H_1(x) = O(x^{-q})$ . In particular,  $H_1 \in L^1$ . Write

$$I_C = \int_0^\infty \cos \theta x (1 - G(x)) dx = c \int_1^\infty \cos \theta x x^{-1} dx + c \int_0^\infty \cos \theta x H_1(x) dx.$$

Now,

$$\begin{aligned} \int_1^\infty \cos \theta x x^{-1} dx &= \int_1^{1/\theta} x^{-1} dx + \int_1^{1/\theta} (\cos \theta x - 1)x^{-1} dx + \int_{1/\theta}^\infty \cos \theta x x^{-1} dx \\ &= \log \frac{1}{\theta} + \int_0^1 (\cos \sigma - 1)\sigma^{-1} d\sigma + \int_1^\infty \cos \sigma \sigma^{-1} d\sigma + O(\theta), \end{aligned}$$

and

$$\begin{aligned} \int_0^\infty \cos \theta x H_1(x) dx &= \int_0^\infty H_1(x) dx + \int_0^{1/\theta} (\cos \theta x - 1)H_1(x) dx \\ &\quad + \int_{1/\theta}^\infty (\cos \theta x - 1)H_1(x) dx = \int_0^\infty H_1(x) dx + O(\theta^{q-1}). \end{aligned}$$

Hence  $I_C = c \log \frac{1}{\theta} + cc_H + O(\theta^{q-1})$ . Also,

$$\begin{aligned} I_S &= \int_0^\infty \sin \theta x (1 - G(x)) dx = c \int_1^\infty \sin \theta x x^{-1} dx + c \int_0^\infty \sin \theta x H_1(x) dx \\ &= c \int_\theta^\infty \sin \sigma \sigma^{-1} d\sigma + O(\theta^{q-1}) = \frac{c\pi}{2} + O(\theta^{q-1}). \end{aligned}$$

Hence

$$\operatorname{Re}(1 - \lambda(\theta)) = \frac{c\pi}{2}\theta(1 + O(\theta^{q-1})), \quad \operatorname{Im} \lambda(\theta) = c\theta(\log \frac{1}{\theta})(1 + c_H(\log \frac{1}{\theta})^{-1} + O((\log \frac{1}{\theta})^{-2})),$$

and the result follows.  $\blacksquare$

**Lemma 9.9** *If  $a = O(n^{1-\delta})$  for some  $\delta > 0$ , then*

$$(i) \quad (\log n) \int_{a/n}^{\pi} \cos n\theta T(\theta) d\theta \ll a^{-1}.$$

$$(ii) \quad (\log n) \int_0^{a/n} \cos n\theta \{T(\theta) - (1 - \lambda(\theta))^{-1}P\} d\theta \ll an^{-1} \log n.$$

**Proof** (i) Since  $T(\theta) = (1 - \lambda(\theta))^{-1}P + O(1)$ , we have  $\operatorname{Re} T(\theta) \ll \theta^{-1}(\log \theta)^{-2}$  for  $\theta \in [0, \epsilon]$  and  $T(\theta) \ll 1$  for  $\theta \in [\epsilon, \pi]$ . The integral splits up into three parts as in Lemma 5.1. As usual  $I_1 \ll n^{-1}$ . Next,

$$\begin{aligned} I_2 &\ll \int_{a/n}^{(a+\pi)/n} \operatorname{Re} T(\theta) d\theta \ll \int_{a/n}^{(a+\pi)/n} \theta^{-1}(\log \theta)^{-2} d\theta \\ &\ll (\log(a/n))^{-1} - (\log((a+\pi)/n))^{-1} = \frac{\log((a+\pi)/n) - \log(a/n)}{\log(a/n) \log((a+\pi)/n)}. \end{aligned}$$

Since  $a = O(n^{1-\delta})$ , we deduce that  $I_2 \ll \log(1 + \pi/a) (\log n)^{-2} \ll a^{-1}(\log n)^{-2}$ .

Finally,  $R(\theta + h) - R(\theta) \ll h^{-1} \log h$  by Proposition 2.7, so

$$I_3 \ll \int_{(a+\pi)/n}^{\pi} \|T(\theta)\| \|T(\theta - \pi/n)\| \|R(\theta) - R(\theta - \pi/n)\| d\theta \ll (\log n)n^{-1}(1 + A)$$

where

$$\begin{aligned} A &= \int_{(a+\pi)/n}^{\epsilon} \|T(\theta)\| \|T(\theta - \pi/n)\| d\theta \ll \int_{a/n}^{\epsilon} (\theta \log \theta)^{-2} d\theta \\ &\ll \int_{a/n}^b (\theta \log \theta)^{-2} d\theta + \int_b^{\epsilon} (\theta \log \theta)^{-2} d\theta \ll (\log b)^{-2} \int_{a/n}^b \theta^{-2} d\theta + (b \log b)^{-2} \int_b^{\epsilon} 1 d\theta \\ &\ll (\log b)^{-2} n/a + (b \log b)^{-2}. \end{aligned}$$

Taking  $b = n^{-\frac{1}{2}\delta}$  say, we obtain  $I_3 \ll (\log n)^{-1}(a^{-1} + n^{-(1-\delta)}) \ll (\log n)^{-1}a^{-1}$ .

(ii) This is immediate since  $T(\theta) = (1 - \lambda(\theta))^{-1}P + O(1)$ .  $\blacksquare$

**Proof of Theorem 9.6** We use Proposition 9.8 to estimate  $\int_0^{a/n} \cos n\theta \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta$ , discarding all terms that are  $O((\log n)^{-3})$ , bearing in mind our eventual choice  $a = n^{\frac{1}{2}}$ .

First we note that for  $j \geq 2$ ,

$$\int_0^{a/n} \theta^{-1}(\log \frac{1}{\theta})^{-j} d\theta = \frac{1}{j-1}(\log \frac{n}{a})^{-(j-1)} \ll (\log n)^{-(j-1)}. \quad (9.1)$$

In particular, taking  $j = 4$  disposes of the  $O(\theta^{-1}(\log \frac{1}{\theta})^{-4})$  term.

Next we consider the  $\theta^{-1}(\log \frac{1}{\theta})^{-3}$  term. Using properties of oscillatory integrals,

$$\int_{1/n}^{a/n} \cos n\theta \theta^{-1} (\log \frac{1}{\theta})^{-3} d\theta = \int_1^a \cos \sigma \sigma^{-1} (\log \frac{n}{\sigma})^{-3} d\sigma \ll (\log n)^{-3},$$

and

$$\int_0^{1/n} (\cos n\theta - 1) \theta^{-1} (\log \frac{1}{\theta})^{-3} d\theta = \int_0^1 (\cos \sigma - 1) \sigma^{-1} (\log \frac{n}{\sigma})^{-3} d\sigma \ll (\log n)^{-3}.$$

Taking  $j = 3$  and  $a = 1$  in equation (9.1), we deduce that

$$\int_0^{a/n} \cos n\theta \theta^{-1} (\log \frac{1}{\theta})^{-3} d\theta = \frac{1}{2} (\log n)^{-2} + O((\log n)^{-3}).$$

To deal with the  $\theta^{-1}(\log \frac{1}{\theta})^{-2}$  term, we use the identity  $\frac{\log n}{\log \frac{n}{\sigma}} = 1 + \frac{\log \sigma}{\log \frac{n}{\sigma}}$ . So

$$\begin{aligned} \int_{1/n}^{a/n} \cos n\theta \theta^{-1} (\log \frac{1}{\theta})^{-2} d\theta &= (\log n)^{-2} \int_1^a \cos \sigma \sigma^{-1} \{\log n / \log \frac{n}{\sigma}\}^2 d\sigma \\ &= (\log n)^{-2} \int_1^a \cos \sigma \sigma^{-1} d\sigma + O((\log n)^{-3}) = (\log n)^{-2} \int_1^\infty \cos \sigma \sigma^{-1} d\sigma + O((\log n)^{-3}), \end{aligned}$$

and

$$\begin{aligned} \int_0^{1/n} (\cos n\theta - 1) \theta^{-1} (\log \frac{1}{\theta})^{-2} d\theta &= (\log n)^{-2} \int_0^1 (\cos \sigma - 1) \sigma^{-1} \{\log n / \log \frac{n}{\sigma}\}^2 d\sigma \\ &= (\log n)^{-2} \int_0^1 (\cos \sigma - 1) \sigma^{-1} d\sigma + O((\log n)^{-3}). \end{aligned}$$

Taking  $j = 2$  and  $a = 1$  in equation (9.1), we deduce that

$$\int_0^{a/n} \cos n\theta \theta^{-1} (\log \frac{1}{\theta})^{-2} d\theta = (\log n)^{-1} + A(\log n)^{-2} + O((\log n)^{-3}),$$

where  $A = \int_0^1 (\cos \sigma - 1) \sigma^{-1} d\sigma + \int_1^\infty \cos \sigma \sigma^{-1} d\sigma$ .

Combining these results, we obtain

$$\frac{2}{\pi} \int_0^{a/n} \cos n\theta \operatorname{Re}\{(1 - \lambda(\theta))^{-1}\} d\theta = c^{-1} - c^{-1} \int_1^\infty H_1(x) dx (\log n)^{-1} + O((\log n)^{-2}),$$

which combined with Lemma 9.9 (taking  $a = n^{\frac{1}{2}}$ ) gives the required result.  $\blacksquare$

### 9.3 Convergence rates in the arcsine law

As mentioned in Remark 2.5, a consequence of Theorem 2.1 is that the Dynkin-Lamperti arcsine law for waiting times holds when  $\beta > \frac{1}{2}$ . In fact, the arcsine law holds for AFN maps for all  $\beta$  [45]. See also [39, 41] for more general transformations. Here we show that our results on second order asymptotics yield a convergence rate.

For  $x \in \bigcup_{j=0}^n f^{-j}Y$ ,  $n \geq 1$ , let

$$Z_n(x) = \max\{0 \leq j \leq n : f^j x \in Y\},$$

denote the time of the last visit of the orbit of  $x$  to  $Y$  during the time interval  $[0, n]$ .

Let  $\zeta_\beta$  denote a random variable distributed according to the  $\mathcal{B}(1 - \beta, \beta)$  distribution:

$$\mathbb{P}(\zeta_\beta \leq t) = d_\beta \int_0^t \frac{1}{u^{1-\beta}} \frac{1}{(1-u)^\beta} du, \quad t \in [0, 1],$$

where  $d_\beta = \frac{1}{\pi} \sin \beta\pi$ .

**Corollary 9.10** *Suppose that  $\beta \in (\frac{1}{2}, 1)$  and that  $\mu(\varphi > n) = cn^{-\beta} + O(n^{-2\beta})$ , where  $c > 0$ . Let  $\gamma = \min\{1 - \beta, \beta - \frac{1}{2}\}$ .*

*Let  $\nu$  be an absolutely continuous probability measure on  $Y$  with density  $g \in \mathcal{B}$ . Then there is a constant  $C > 0$  independent of  $\nu$  such that*

$$|\nu\{\frac{1}{n}Z_n \leq t\} - \mathbb{P}(\zeta_\beta \leq t)| \leq C\|g\|n^{-\gamma}.$$

**Proof** Following Thaler [40], we notice that

$$\nu\{\frac{1}{n}Z_n \leq t\} = \sum_{0 \leq j \leq nt} \nu(f^{-j}\{\varphi > n - j\}), \quad (9.2)$$

$$\begin{aligned} \nu(f^{-j}\{\varphi > n - j\}) &= \int_X 1_{\{\varphi > n - j\}} \circ f^j 1_Y g d\mu = \int_X 1_{\{\varphi > n - j\}} L^j(1_Y g) d\mu \\ &= \int_Y 1_{\{\varphi > n - j\}} T_j g d\mu. \end{aligned}$$

By Theorem 9.1,  $T_j g = c^{-1}d_\beta j^{-(1-\beta)}(1 + O(j^{-(1-\beta)}) + O(\|g\|j^{-(\beta-\frac{1}{2})}))$  uniformly on  $Y$ . Combined with the assumption on  $\mu(\varphi > n)$ , we obtain

$$\begin{aligned} \nu(f^{-j}\{\varphi > n - j\}) &= d_\beta j^{-(1-\beta)}(n - j)^{-\beta} (1 + O(j^{-(1-\beta)}) + O(\|g\|j^{-(\beta-\frac{1}{2})})) (1 + O((n - j)^{-\beta})). \end{aligned}$$

Since functions of the form  $s^{-a}(n - s)^{-b}$  have only one turning point, replacing the sum in (9.2) by an integral introduces only three errors all of order  $\|g\|n^{-1}$  and so  $\mu\{\frac{1}{n}Z_n \leq t\} = d_\beta I + O(\|g\|n^{-1})$ , where

$$\begin{aligned} I &= \int_0^{nt} s^{-(1-\beta)}(n - s)^{-\beta} (1 + O(s^{-(1-\beta)}) + O(\|g\|s^{-(\beta-\frac{1}{2})})) (1 + O((n - s)^{-\beta})) ds \\ &= \int_0^t u^{-(1-\beta)}(1 - u)^{-\beta} du + O(n^{-(1-\beta)}) + O(\|g\|n^{-(\beta-\frac{1}{2})}), \end{aligned}$$

as required. ■

**Remark 9.11** (a) The proof shows that for any  $q \geq 0$ ,

$$\nu(\{\frac{1}{n}Z_n \leq t\}) = \sum_{k=0}^q b_{\beta,k} \mathbb{P}(\zeta_{\beta,k+1} \leq t) n^{-k(1-\beta)} + O(n^{-(q+1)(1-\beta)}) + O(\|g\| n^{-(\beta-\frac{1}{2})}).$$

where  $\zeta_{\beta,k}$  is the random variable with density proportional to  $u^{-k(1-\beta)}(1-u)^{-\beta}$  and  $b_{\beta,k} = d_{\beta,k} / \int_0^1 u^{-(k+1)(1-\beta)}(1-u)^{-\beta} du$ . Here  $b_{\beta,0} = 1$  and  $\zeta_{\beta,1} = \zeta_{\beta}$ .

Thus for  $\beta$  close to 1, we obtain asymptotic expansions to arbitrarily high order, and the error rate  $n^{-\gamma}$  is optimal for  $\beta \geq \frac{3}{4}$ .

(b) For  $x \in X$ , let  $Y_n(x) = \min\{k > n : f^k x \in Y\}$ . Then  $Y_k > n$  if and only if  $Z_n \leq k$  so that the arcsine law is equivalent to strong distributional convergence of  $\frac{1}{n}Y_n$  to  $\zeta_{\beta}^{-1}$  (see for example [39]). It is easily verified that the convergence rate in Corollary 9.10 holds also for  $\frac{1}{n}Y_n$ .

## 10 Convergence results for $L^n$

Sections 2 to 9 were concerned with the analysis of the sequence of renewal operators  $T_n$  given by  $T_n v = 1_Y L^n(1_Y v)$ . An important issue is to study the iterates  $L^n$  themselves. In Subsection 10.1, we show how convergence on  $Y$  implies convergence almost everywhere on  $X$ . In Subsection 10.2, we consider observables not supported on  $Y$ .

### 10.1 Convergence on $X$

Theorem 2.1 gives (uniform) convergence results on  $Y$  for observables  $v \in \mathcal{B}$ . Recall that  $Y$  can be regarded as a first return set for both the underlying system  $f : X \rightarrow X$  and the tower map  $f_{\Delta} : \Delta \rightarrow \Delta$  introduced in Subsection 2.4. We now show that observables  $v \in \mathcal{B}$  enjoy pointwise convergence everywhere on  $\Delta$  and almost everywhere on  $X$ .

**Proposition 10.1** *Let  $v \in L^1(\Delta)$ ,  $w(n) \in \mathbb{R}$ ,  $A \in \mathbb{R}$ . Suppose that  $w(n)L_{\Delta}^n v \rightarrow A$  pointwise on  $Y$ . Then  $w(n)L_{\Delta}^n v \rightarrow A$  pointwise on  $\Delta$ .*

**Proof** Let  $p = (y, j) \in \Delta$ . Then  $f_{\Delta}^{-j} p$  consists of the single preimage  $(y, 0) \cong y$ , and  $w(n)(L_{\Delta}^n v)(p) = w(n)(L_{\Delta}^{n-j} v)(y) \rightarrow A$ .  $\blacksquare$

Let  $\pi : \Delta \rightarrow X$  be the projection  $\pi(y, j) = f^j y$ . Let  $1 \leq p, q \leq \infty$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ . Define  $\pi^* : L^p(X) \rightarrow L^p(\Delta)$ ,  $\pi^* v = v \circ \pi$ , and by duality define  $\hat{\pi} : L^q(\Delta) \rightarrow L^q(X)$ ,  $\int_{\Delta} \hat{\pi} v w d\mu_X = \int_X v \pi^* w d\mu_{\Delta}$ . As usual,  $\|\pi^*\|_p = \|\hat{\pi}\|_q = 1$  and we have the standard properties  $\hat{\pi}1 = 1$ ,  $\hat{\pi}\pi^* = I$ ,  $\hat{\pi}L_{\Delta} = L\hat{\pi}$ ,  $\hat{\pi}(1_{\pi^{-1}E} v) = 1_E \hat{\pi}v$ .

**Proposition 10.2** *Let  $v \in L^1(\Delta)$ ,  $w(n) \in \mathbb{R}$ ,  $A \in \mathbb{R}$ . Suppose that  $w(n)L_{\Delta}^n v \rightarrow A$  almost everywhere on  $\Delta$ . Then  $w(n)L^n \hat{\pi}v \rightarrow A$  almost everywhere on  $X$ .*

**Proof** Suppose for contradiction that  $E \subset X$  is a set of positive finite measure such that everywhere on  $E$ ,  $w(n)L^n\hat{\pi}v$  fails to converge to  $A$ . By assumption,  $w(n)L^n_\Delta v \rightarrow A$  almost everywhere on  $\pi^{-1}E$ . By Egorov's Theorem, there is a subset  $C \subset \pi^{-1}E$  of positive measure such that  $w(n)L^n_\Delta v \rightarrow A$  uniformly on  $C$ . Indeed  $\mu_\Delta(\pi^{-1}E \setminus C)$  is arbitrarily small, and since  $\pi$  has only countably many branches it follows from an  $\epsilon/2^k$  argument that we can choose  $C = \pi^{-1}E'$  where  $E'$  is a positive measure subset of  $E$ .

In particular,  $\|1_{\pi^{-1}E'}(w(n)L^n_\Delta v - A)\|_{L^\infty(\Delta)} \rightarrow 0$  and since  $\hat{\pi} : L^\infty(\Delta) \rightarrow L^\infty(X)$  is bounded,  $\|\hat{\pi}\{1_{\pi^{-1}E'}(w(n)L^n_\Delta v - A)\}\|_{L^\infty(X)} \rightarrow 0$ . But  $\hat{\pi}\{1_{\pi^{-1}E'}(w(n)L^n_\Delta v - A)\} = 1_{E'}\hat{\pi}(w(n)L^n_\Delta v - A) = 1_{E'}(w(n)L^n\hat{\pi}v - A)$  so we conclude that  $w(n)L^n\hat{\pi}v \rightarrow A$  on  $E'$  which is the desired contradiction.  $\blacksquare$

**Corollary 10.3** *If  $\beta \in (\frac{1}{2}, 1]$  and  $v \in \mathcal{B}$ , then  $\lim_{n \rightarrow \infty} m(n)n^{1-\beta}L^n v = d_\beta \int_Y v d\mu$  uniformly on  $Y$  and almost everywhere on  $X$ .*

**Proof** Since  $v$  is supported on  $Y$ , we can regard  $v$  as an observable on  $\Delta$  or on  $X$  and  $\hat{\pi}v = v$ . Also,  $T_n v = 1_Y L^n v = 1_Y L^n_\Delta v$ . Theorem 2.1 immediately implies uniform convergence on  $Y$ . By Propositions 10.1 and 10.2,  $m(n)n^{1-\beta}L^n_\Delta v$  converges pointwise on  $\Delta$  and  $m(n)n^{1-\beta}L^n v$  converges almost everywhere on  $X$ .  $\blacksquare$

## 10.2 Convergence for general observables

In this subsection, we enlarge the class of observables so that they need not be supported on  $Y$ . Define  $X_k = f^{-k}Y \setminus \bigcup_{j=0}^{k-1} f^{-j}Y$ . Thus  $z \in X_k$  if and only if  $k \geq 0$  is least such that  $f^k z \in Y$ . (In particular,  $X_0 = Y$ .)

Given  $v \in L^\infty(X)$ , define  $v_k = 1_{X_k}v$ . Then  $L^k v_k$  is supported in  $Y$  and  $L^n v_k$  vanishes on  $Y$  for all  $n < k$ . If  $n \geq k$ , we have  $1_Y L^n v_k = T_{n-k} L^k v_k$ .

Write  $v \in \mathcal{B}(X)$  if  $v \in L^1(X)$  and  $L^k v_k \in \mathcal{B}$  for each  $k \geq 0$ .

**Theorem 10.4** *Let  $\beta \in (\frac{1}{2}, 1]$ . Suppose that  $v \in \mathcal{B}(X)$  and moreover that  $\sum_{n=0}^\infty \|L^n v_n\| < \infty$  and either (i)  $\|L^n v_n\| = o(n^{-1})$  or (ii)  $\sum_{k=n}^\infty \|L^k v_k\| = o((m(n)n^{1-\beta})^{-1})$ . Then  $\lim_{n \rightarrow \infty} m(n)n^{1-\beta}L^n v = d_\beta \int_X v d\mu$  uniformly on  $Y$  and pointwise on  $X$ .*

**Proof** Let  $w(n) = d_\beta^{-1}m(n)n^{1-\beta}$ . First we show that

$$w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| \rightarrow 0. \quad (10.1)$$



In case (i),  $\|L^n v_n\| = o(n^{-1})$ , it follows from Karamata that

$$\begin{aligned} w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| &\ll n^{-1} w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} (j \|L^j v_j\|) \\ &\leq n^{-1} w(n) \left( \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \right) \max_{j \geq n/2} j \|L^j v_j\| \ll \max_{j \geq n/2} j \|L^j v_j\| \rightarrow 0. \end{aligned}$$

In case (ii),

$$w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| \ll w(n) \sum_{n/2 \leq j \leq n} \|L^j v_j\| \ll w(n/2) \sum_{j \geq n/2} \|L^j v_j\| \rightarrow 0.$$

Hence (10.1) is verified in both cases.

Next we prove uniform convergence on  $Y$ . Let  $c_{j,n} = \frac{w(n)}{w(n-j)} - 1$ . By Theorem 2.1,  $T_n = w(n)^{-1}P + S_n$  where  $S_n = o(w(n)^{-1})$ . Hence on  $Y$ ,

$$\begin{aligned} w(n)L^n v - \int v &= w(n) \sum_{j=0}^n T_{n-j} L^j v_j - \sum_{j=0}^{\infty} \int v_j \\ &= w(n) \sum_{j=0}^n w(n-j)^{-1} \int L^j v_j - \sum_{j=0}^n \int L^j v_j + w(n) \sum_{j=0}^n S_{n-j} L^j v_j - \sum_{j>n} \int v_j \\ &= \sum_{j=0}^n c_{j,n} \int L^j v_j + w(n) \sum_{j=0}^n S_{n-j} L^j v_j - \sum_{j>n} \int v_j, \end{aligned}$$

and so

$$|w(n)L^n v - \int v| \leq \sum_{j=0}^n |c_{j,n}| \left| \int L^j v_j \right| + w(n) \sum_{j=0}^n \|S_{n-j}\| \|L^j v_j\| + \left| \sum_{j>n} \int v_j \right|.$$

It is immediate that the third term converges to zero. Write  $\|S_n\| = w(n)^{-1} a_n$  where  $a_n = o(1)$ . Then the second term satisfies

$$\begin{aligned} w(n) \sum_{j=0}^n \|S_{n-j}\| \|L^j v_j\| &= w(n) \sum_{0 \leq j < n/2} w(n-j)^{-1} a_{n-j} \|L^j v_j\| \\ &\quad + w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} a_{n-j} \|L^j v_j\| \\ &\ll \sum_{0 \leq j < n/2} a_{n-j} \|L^j v_j\| + w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| \\ &\leq \left( \sum_{k=0}^{\infty} \|L^k v_k\| \right) \max_{j \geq n/2} a_j + w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| \rightarrow 0 \end{aligned}$$

by (10.1), summability of  $\|L^k v_k\|$  and the definition of  $a_n$ .

The first term satisfies

$$\begin{aligned} \sum_{j=0}^n |c_{j,n}| \|f L^j v_j\| &\leq \sum_{0 \leq j < n/2} |c_{j,n}| \|L^j v_j\| + \sum_{n/2 \leq j \leq n} |c_{j,n}| \|L^j v_j\| \\ &\ll \sum_{0 \leq j < n/2} |c_{j,n}| \|L^j v_j\| + w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\|. \end{aligned}$$

Again  $w(n) \sum_{n/2 \leq j \leq n} w(n-j)^{-1} \|L^j v_j\| \rightarrow 0$  by (10.1). Since  $\lim_{n \rightarrow \infty} c_{j,n} = 0$  for each fixed  $j$  and  $\max_{j,n: 0 \leq j < n/2} |c_{j,n}| < \infty$ , it follows from summability of  $L^n v_n$  that  $\sum_{0 \leq j < n/2} |c_{j,n}| \|L^j v_j\| \rightarrow 0$ . This completes the proof of uniform convergence on  $Y$ .

To prove pointwise convergence on  $X$ , define  $u = \pi^* v : \Delta \rightarrow \mathbb{R}$  and note that  $\hat{\pi} u = v$ . Also define  $\Delta_k = f_{\Delta}^{-k} Y \setminus \bigcup_{j=0}^{k-1} f_{\Delta}^{-j} Y$ . Then  $\Delta_0 = Y$  and for  $k \geq 1$ ,  $\Delta_k$  consists of those points  $(y, j) \in \Delta$  with  $j = \varphi(y) - k > 0$ . Note also that  $\pi^{-1} X_k \subset \Delta_k$  (since  $(y, j) \in \pi^{-1} X_k$  if and only if  $f^j y \in X_k$ , that is  $\varphi(y) = j + k$ ). In particular,  $u_k = \pi^* v_k$  is supported in  $\Delta_k$ . Hence  $L_{\Delta}^k u_k$  is supported in  $Y$  and  $L_{\Delta}^k u_k = \hat{\pi} L_{\Delta}^k u_k = L^k \hat{\pi} u_k = L^k v_k$ . In particular,  $L_{\Delta}^k u_k$  inherits the assumptions on  $L^k v_k$ , and the argument above shows that  $w(n) L_{\Delta}^n u \rightarrow \int_{\Delta} u d\mu_{\Delta}$  uniformly on  $Y$ . By Proposition 10.1, pointwise convergence extends to  $\Delta$ . By Proposition 10.2, pointwise convergence for  $u$  drops down to pointwise convergence for  $v$ .  $\blacksquare$

Note that Theorem 10.4 includes the case where  $v$  is supported on  $\bigcup_{j=0}^k X_j$  for some  $k$ , and hence significantly extends Theorem 2.1.

In the next result, we extend Theorem 2.3, and we drop the assumptions on  $L^n v_n$  in Theorem 10.4.

**Proposition 10.5** *Let  $\beta \in (0, 1]$  and  $v \in \mathcal{B}(X)$ .*

(a) *For each  $y \in Y$ , there is a zero density set  $E \subset \mathbb{N}$  such that  $\lim_{n \rightarrow \infty, n \notin E} m(n) n^{1-\beta} (L^n v)(y) = d_{\beta} \int_X v d\mu$ .*

(b) *If  $v \geq 0$ , then  $\liminf_{n \rightarrow \infty} m(n) n^{1-\beta} L^n v = d_{\beta} \int_X v d\mu$  pointwise on  $Y$ .*

**Proof** Let  $w(n) = d_{\beta}^{-1} m(n) n^{1-\beta}$ . By the argument in the proof of Theorem 2.3, it suffices to prove the  $\geq$  inequality in part (b). Let  $v \geq 0$  and define  $v(k) = \sum_{j=0}^k v_j$ . By Theorem 2.3(b),  $\liminf_{n \rightarrow \infty} w(n) 1_Y L^n v \geq \liminf_{n \rightarrow \infty} w(n) 1_Y L^n v(k) = \sum_{j=0}^k \liminf_{n \rightarrow \infty} w(n) 1_Y L^n v_j = \sum_{j=0}^k \int v_j = \int v(k)$ . Since  $k$  is arbitrary,  $\liminf_{n \rightarrow \infty} w(n) 1_Y L^n v \geq \int v$  as required.  $\blacksquare$

### 10.3 Second order asymptotics on $X$

Under the assumptions of Theorem 9.1, we can investigate second order asymptotics in Theorem 10.4. For example, we have the following:

**Theorem 10.6** *Suppose that  $\beta \in (\frac{1}{2}, 1)$  and that  $\mu(\varphi > n) = cn^{-\beta} + O(n^{-2\beta})$  where  $c > 0$ . Suppose further that  $v \in \mathcal{B}(X)$  and that (i)  $\|L^k v_k\| = O(k^{-p})$ , and (ii)  $\int L^k v_k d\mu = O(k^{-q})$ , where  $p > \max\{\frac{3}{2} - \beta, \beta\}$  and  $q > 1$ . Then*

$$n^{1-\beta} L^n v = c^{-1} d_\beta \int_X v d\mu + O(n^{-\gamma} \|v\|) \text{ uniformly on } Y,$$

where  $\gamma = \min\{1 - \beta, \beta - \frac{1}{2}, q - 1\}$  if  $p > 1$ , and  $\gamma = \min\{p - \beta, p + \beta - \frac{3}{2}, q - 1\}$  if  $p > \max\{\frac{3}{2} - \beta, \beta\}$ .

**Proof** We assume without loss that  $q \in (1, 2)$ . In the notation of the proof of Theorem 10.4,

$$cd_\beta^{-1} n^{1-\beta} L^n v - \int v \ll \sum_{j=0}^n |c_{j,n}| |\int L^j v_j| + n^{1-\beta} \sum_{j=0}^n \|S_{n-j}\| \|L^j v_j\| + \left| \sum_{j>n} \int L^j v_j \right|.$$

It is immediate that the third term is  $O(n^{-(q-1)})$ . Since  $c_{j,n} = (n/(n-j))^{1-\beta} - 1 = (1 + (j/(n-j)))^{1-\beta} - 1 \ll j/(n-j)$ , the first term satisfies

$$\begin{aligned} \sum_{j=0}^n |c_{j,n}| |\int L^j v_j| &\leq \sum_{0 \leq j < n/2} |c_{j,n}| |\int L^j v_j| + \sum_{n/2 \leq j \leq n} |c_{j,n}| |\int L^j v_j| \\ &\ll \sum_{0 \leq j < n/2} (j/(n-j)) j^{-q} + n^{-q} n^{1-\beta} \sum_{n/2 \leq j \leq n} (n-j)^{-(1-\beta)} \\ &\leq n^{-1} \sum_{1 \leq j < n/2} j^{1-q} + n^{1-\beta-q} \sum_{j=1}^n j^{-(1-\beta)} \ll n^{-(q-1)}. \end{aligned}$$

By Theorem 9.1,  $\|S_n\| = O(n^{-(1-\beta+\gamma_1)})$  where  $\gamma_1 = \min\{1 - \beta, \beta - \frac{1}{2}\}$ . Hence the second term satisfies

$$n^{1-\beta} \sum_{j=0}^n \|S_{n-j}\| \|L^j v_j\| \ll n^{1-\beta} \{(n^{-(1-\beta+\gamma_1)}) \star (n^{-p})\} \ll \begin{cases} n^{-\gamma_1}, & p > 1 \\ n^{-(\gamma_1+p-1)}, & p < 1 \end{cases}$$

as required. ■

## 11 Examples

In this section, we apply our results to specific examples. In Subsection 11.1, we describe a method for verifying our functional-analytic hypotheses (H1), (H2), that suffices for our purposes. In Subsection 11.2, we consider situations where the first return map  $F : Y \rightarrow Y$  is Gibbs-Markov. This includes the nonuniformly expanding maps studied by Thaler [38] and parabolic rational maps of the complex plane [6]. In Subsection 11.3, we consider the full class of AFN maps [44]. In Subsection 11.4, we specialise to the case of Pomeau-Manneville intermittency maps (1.1).

## 11.1 Verification of hypotheses (H1) and (H2)

In our examples, (H1) can be verified in the process of verifying (H2), so we focus on the latter. The standard approach (cf. [18, Lemma 6.7] and [37, Section 5]) to (H2) proceeds via the following result.

**Proposition 11.1** *Suppose that  $F : Y \rightarrow Y$  is ergodic. Assume that (1)  $R(z) : \mathcal{B} \rightarrow \mathcal{B}$  has essential spectral radius strictly less than 1 for every  $z \in \bar{\mathbb{D}}$ . (2) For each  $\theta \in (0, 2\pi)$ , there are no nontrivial  $L^2$  solutions to the equation  $v \circ F = e^{i\theta\varphi}v$  a.e. Then (H2) is satisfied.*

**Proof** By (1), it suffices to consider generalized eigenfunctions  $v \in L^2$  for the operator  $R(z)$ . Suppose that  $R(z)v = v$  where  $v \in L^2$  is nonzero. Write  $z = \rho e^{i\theta}$ ,  $\rho \in [0, 1]$ ,  $\theta \in [0, 2\pi)$ . Then  $|v|_2 = |R(z)v|_2 = |R(\rho^\varphi e^{i\theta\varphi}v)|_2 = |\rho^\varphi v|_2 \leq |\rho^\varphi|_\infty |v|_2 \leq \rho |v|_2$ , so  $\rho = 1$  and  $R(e^{i\theta})v = v$ . The  $L^2$  adjoint of  $U = R(e^{i\theta})$  is the operator  $U^*v = e^{-i\theta\varphi}v \circ F$  and an elementary calculation shows that  $|U^*v - v|_2^2 = |Uv|_2^2 - |v|_2^2 = 0$ . Hence  $v \circ F = e^{i\theta\varphi}v$ . By (2),  $\theta = 0$ . Hence we have established (H2)(ii).

When  $z = 1$ , the eigenvalue 1 is isolated in the spectrum by (1) and the eigenvalue is simple by ergodicity of  $F$ , so (H2)(i) is valid.  $\blacksquare$

**Definition 11.2** Suppose that  $Y$  is a topological space, that  $(Y, \mu)$  is a probability space, and that  $F : Y \rightarrow Y$  is a measure preserving transformation. Let  $\varphi : Y \rightarrow \mathbb{R}$  be a measurable map. We say that  $(F, \varphi)$  satisfies property (\*) if for every  $\theta \in [0, 2\pi]$  and every measurable solution  $v : Y \rightarrow S^1$  to the equation  $v \circ F = e^{i\theta\varphi}v$  a.e., there exists an open set  $U \subset Y$  such that  $v$  is constant almost everywhere on  $U$ .

**Lemma 11.3** *Suppose that  $f$  is topologically mixing with first return map  $F = f^\varphi : Y \rightarrow Y$ . Assume that  $(F, \varphi)$  satisfies property (\*). Then there are no nontrivial measurable solutions  $v : Y \rightarrow \mathbb{C}$  to the equation  $v \circ F = e^{i\theta\varphi}v$  for all  $\theta \in (0, 2\pi)$ .*

**Proof** If  $v$  is a nontrivial solution, then by (\*) there is an open set  $U \subset Y$  on which  $v$  is almost everywhere constant. Now we follow the second half of the proof of [18, Lemma 6.7]. Since  $f$  is topologically mixing, there exists  $N \geq 1$  such that  $f^n U \cap U \neq \emptyset$  for all  $n \geq N$ . In particular, for each  $n \geq N$  we can choose  $y \in U$  such that  $f^n y \in U$  and  $v(y) = v(f^n y) \neq 0$ .

Let  $0 = k_0 < k_1 < \dots < k_p = n$  be the successive return times of  $y$  to  $U$ . Then  $T^n y = F^p y$  and  $n = \varphi_p(y) = \sum_{j=0}^{p-1} \varphi(F^j y)$ . Hence

$$e^{i\theta n} = \prod_{j=0}^{p-1} e^{i\theta\varphi(F^j y)} = \prod_{j=0}^{p-1} \frac{v(F^{j+1}y)}{v(F^j y)} = \frac{v(f^n y)}{v(y)} = 1.$$

Taking  $n = N$  and  $n = N + 1$ , we deduce that  $e^{i\theta} = 1$  which is a contradiction.  $\blacksquare$

## 11.2 Maps with Gibbs-Markov first return maps

A large class of examples covered by our methods are those with first return maps that are Gibbs-Markov. This includes parabolic rational maps of the complex plane (Aaronson *et al* [6]) and Thaler's class of interval maps with indifferent fixed points [38] (in particular the family (1.1)).

We recall the key definitions [3]. Let  $(X, \mu)$  be a Lebesgue space with countable measurable partition  $\alpha_X$ . Let  $f : X \rightarrow X$  be an ergodic, conservative, measure-preserving, Markov map transforming each partition element bijectively onto a union of partition elements. Recall that  $f$  is *topologically mixing* if for all  $a, b \in \alpha_X$  there exists  $N \geq 1$  such that  $b \subset f^n a$  for all  $n \geq N$ .

Let  $Y$  be a union of partition elements with  $\mu(Y) \in (0, \infty)$ . Define the first return time  $\varphi : Y \rightarrow \mathbb{R}$  and first return map  $F = f^\varphi : Y \rightarrow Y$ . Let  $\alpha$  be the partition of  $Y$  consisting of nonempty cylinders of the form  $a \cap (\bigcap_{j=1}^{n-1} T^{-j} \xi_j) \cap T^{-n} \alpha$  where  $a, \xi_j \in \alpha_X$ , and  $a \subset Y$ ,  $\xi_j \subset X \setminus Y$ . Fix  $\tau \in (0, 1)$  and define  $d_\tau(x, y) = \tau^{s(x, y)}$  where the *separation time*  $s(x, y)$  is the greatest integer  $n \geq 0$  such that  $F^n x$  and  $F^n y$  lie in the same partition element in  $\alpha$ . It is assumed that the partition  $\alpha$  separates orbits of  $F$ , so  $s(x, y)$  is finite for all  $x \neq y$ . Then  $d_\tau$  is a metric. Let  $\text{Lip}(Y)$  be the Banach space of  $d_\tau$ -Lipschitz functions  $v : Y \rightarrow \mathbb{R}$  with norm  $\|v\| = |v|_\infty + \text{Lip}(v)$ .

Define the potential function  $p = \log \frac{d\mu}{d\mu \circ F} : Y \rightarrow \mathbb{R}$ . We require that  $p$  is uniformly piecewise Lipschitz: that is,  $p|_a$  is  $d_\tau$ -Lipschitz for each  $a \in \alpha$  and the Lipschitz constants can be chosen independent of  $a$ . We also require the big images condition  $\inf_a \mu(Fa) > 0$ . A *Gibbs-Markov map* is a Markov map with uniformly piecewise Lipschitz potential and satisfying the big images property.

**Proposition 11.4** *Suppose that  $(X, \mu)$  is a Lebesgue space, that  $f : X \rightarrow X$  is an ergodic, conservative, measure preserving, topologically mixing, Markov map, and that  $Y \subset X$  is a union of partition elements with  $\mu(Y) \in (0, \infty)$ . Suppose further that the first return map  $F = f^\varphi : Y \rightarrow Y$  is Gibbs-Markov. Then the Banach space  $\mathcal{B} = \text{Lip}(Y)$  satisfies hypotheses (H1) and (H2).*

**Proof** Since the details can be found in [18, 37], we only sketch the argument. By equation (8) in the proof of [18, Lemma 6.7], there is a constant  $C > 0$  such that  $\|R(z)^n v\| \leq C(|v|_\infty + \tau^n \|v\|)$  for all  $v \in \mathcal{B}$ ,  $z \in \mathbb{D}$ ,  $n \geq 1$ . Since the unit ball in  $\mathcal{B}$  is compact in  $L^\infty$ , it follows from [23] that the essential spectral radius of  $R(z)$  is at most  $\tau$  establishing property (1) of Proposition 11.1. Property (\*) follows from [4, Theorem 3.1]: measurable solutions  $v$  are constant almost everywhere on each partition element of  $\alpha$  (even  $F\alpha$ ). Hence property (2) of Proposition 11.1 follows from Lemma 11.3. This completes the verification of (H2). (H1) is established during the proof of [18, Lemma 6.7], see [18, Lemma 6.4]. ■

**Corollary 11.5** *In the setting of Proposition 11.4, if in addition  $\mu(X) = \infty$  and  $\mu(y \in Y : \varphi(y) > n)$  is regularly varying with index  $\beta \in (0, 1]$ , then our main results*

(including Theorems 2.1, 2.2, 2.3) apply. ■

**Example 11.6 (Parabolic rational maps of the complex plane)** Let  $f : \bar{\mathbb{C}} \rightarrow \bar{\mathbb{C}}$  be a rational map of the Riemann sphere with spherical metric  $d$ . A period  $k$  point  $z \in \mathbb{C}$  is *rationally indifferent* if  $(f^k)'(z)$  is a root of unity. The map  $f$  is *parabolic* if  $J$  contains no critical points and contains at least one rationally indifferent periodic point [11]

Aaronson *et al.* [6, Section 9] establish a number of properties of parabolic rational maps. Such maps are topologically mixing, conservative and exact with respect to Lebesgue measure and possess a  $\sigma$ -finite invariant measure  $\mu$  equivalent to Lebesgue. Moreover, there is a Gibbs-Markov first return map  $F = f^\varphi : Y \rightarrow Y$  where  $\mu(Y) \in (0, \infty)$ . Criteria are given for  $\mu(X)$  to be finite or infinite, and in the infinite case it is shown that  $\mu(\varphi > n) \sim Cn^{-\beta}$  where  $C > 0$  and  $\beta \in (0, 1]$ . For any  $\eta > 0$ , it is possible to choose  $\tau \in (0, 1)$  so that  $C^\eta(Y) \subset \text{Lip}(Y)$ . By Corollary 11.5, our main results apply to Hölder observables supported on  $Y$ .

**Example 11.7 (Thaler maps)** Thaler [38] considers a class of topologically mixing one-dimensional maps  $f : X \rightarrow X$ ,  $X = [0, 1]$  for which there is a countable measurable partition  $\{B(k) : k \in I\}$  consisting of intervals, and a nonempty finite set  $J \subset I$  such that each  $B(j)$ ,  $j \in J$ , contains an indifferent fixed point  $x_j$  with  $f'(x_j) = 1$ . It is required that

- (1)  $f|_{B(k)}$  is twice differentiable and  $\overline{fB(k)} = [0, 1]$  for all  $k$ .
- (2)  $|f'| \geq \rho(\epsilon) > 1$  on  $\bigcup_{k \in I} B(k) \setminus \bigcup_{j \in J} (x_j - \epsilon, x_j + \epsilon)$  for each  $\epsilon > 0$ .
- (3) For each  $j \in J$  there exists  $\eta > 0$  such that  $f'$  is decreasing on  $(x_j - \eta, x_j) \cap B(j)$  and increasing on  $(x_j, x_j + \eta) \cap B(j)$ .
- (4)  $f''/(f')^2$  is bounded on  $\bigcup_{k \in I} B(k)$ .

Such a map  $f$  is conservative and exact with respect to Lebesgue measure, and admits an infinite  $\sigma$ -finite invariant measure  $\mu$  equivalent to Lebesgue measure  $m$ . The density  $h = d\mu/dm$  is Hölder, bounded below, and bounded above on compact subsets of  $X'$ . As a special case of the construction in Zweimüller [45],  $f$  has a Gibbs-Markov first return map  $F = f^\varphi : Y \rightarrow Y$ , where  $\mu(Y) \in (0, \infty)$ . Furthermore, for every compact set  $C \subset X'$  (the complement of the indifferent fixed points), the first return set  $Y$  can be chosen to contain  $C$ . As shown in [38], a sufficient condition for regularly varying return tail probabilities is that  $f$  has a “good” asymptotic expansion near each indifferent fixed point. For example, it suffices that  $f(x) = x + a_j|x - x_j|^{p_j+1} + o(|x - x_j|^{p_j+1})$  as  $x \rightarrow x_j$  for each  $j \in J$ , where  $a_j \neq 0$ ,  $p_j \geq 1$  and  $p = \max_j p_j > 1$ . In this case  $\mu(\varphi > n) \sim Cn^{-1/p}$  where  $C > 0$ .

To summarise, suppose that  $f$  is a Thaler map and  $\mu(\varphi > n) = \ell(n)n^{-\beta}$  is regularly varying with  $\beta \in (0, 1]$ . Set  $w(n) = d_\beta^{-1}m(n)n^{1-\beta}$ . By Corollary 11.5, our

main results apply to Hölder continuous observables  $v : X \rightarrow \mathbb{R}$  supported on a compact subset of  $X'$ . In particular, if  $\beta \in (\frac{1}{2}, 1]$ , we obtain uniform convergence on compact subsets of  $X'$ .

**Theorem 11.8** *Suppose that  $f : X \rightarrow X$  is a Thaler map with regularly varying tails,  $\beta \in (\frac{1}{2}, 1]$ . Then  $\lim_{n \rightarrow \infty} w(n)L^n v = \int_X v d\mu$  uniformly on compact subsets of  $X'$  for all  $v$  of the form  $v = u/h$  where  $u$  is Riemann integrable on  $X$ .*

**Proof** Suppose first that  $u$  is Hölder. Fix a first return set  $Y$  chosen so that  $f(Y) = X$ . Define the sets  $X_k$  as in Subsection 10.2 and write  $v = \sum v_k$  where  $v_k = v|_{X_k}$ . We claim that  $\|L^k v_k\| \ll \mu(\varphi = k+1)\|u\|_{C^\eta}$ . Hence  $\sum_{k \geq n} \|L^k v_k\| \leq \mu(\varphi > n)\|u\|_{C^\eta} = O(\ell(n)n^{-\beta}) = o(w(n))^{-1}$  since  $\beta > \frac{1}{2}$ , verifying condition (ii) in Theorem 10.4. Also  $u$  is Lebesgue integrable, so  $v \in L^1(X, \mu)$ . It follows from Theorem 10.4 that  $\lim_{n \rightarrow \infty} w(n)L^n v = \int_X v d\mu$  uniformly on  $Y$ .

To prove the claim, let  $\tilde{L}$  and  $\tilde{R}$  denote the (unnormalised) transfer operators for  $f$  and  $F$  respectively corresponding to Lebesgue measure  $m$  (rather than  $\mu$ ). In particular,  $L = h^{-1}\tilde{L}h$ . Every point in  $X_k$  has a preimage in  $Y$ . To simplify notation, suppose that such preimages are unique (otherwise specify one of the preimages and omit the other preimages in the argument below). Write  $(\tilde{L}v)(x) = \sum_{f x' = x} g(x')v(x')$  and  $(\tilde{L}^n v)(x) = \sum_{f^n x' = x} g_n(x')v(x')$  where  $g_n(x) = g(x)g(fx) \cdots g(f^{n-1}x)$ . Similarly, write  $(\tilde{R}v)(y) = \sum_{F y' = y} G(y')v(y')$ . Let  $u_k = u|_{X_k}$ . Then

$$\begin{aligned} (\tilde{L}^k u_k)(x) &= \sum_{f^k x' = x} g_k(x') 1_{X_k}(x') u(x') = \sum_{y \in \{\varphi = k+1\}, f^{k+1} y = x} g(y)^{-1} g_{k+1}(y) u(fy) \\ &= \sum_{F y = x} g(y)^{-1} 1_{\{\varphi = k+1\}}(y) G(y) u(fy) = \sum_a g(y_a)^{-1} G(y_a) u(fy_a), \end{aligned}$$

where the summation is over those  $a$  with  $\varphi|_a = k+1$ , and  $y_a$  is the unique point in  $a$  such that  $F y_a = x$ . For Gibbs-Markov maps it is standard that  $\|1_a G\| \ll \mu(a)$ . For the systems in [38],  $f'$  is bounded and  $f$  is uniformly expanding on  $Y$  so that  $\|1_{\{\varphi = k+1\}}(u \circ f)\| \leq C\|u_k\|_{C^\eta(X_k)}$  and  $\|1_Y g^{-1}\| < \infty$ . Hence  $\|\tilde{L}^k u_k\| \ll \sum_a \mu(a)\|u\|_{C^\eta(X_k)} = \mu(\varphi = k+1)\|u\|_{C^\eta(X_k)}$ , and so  $\|L^k v_k\| = \|h^{-1}\tilde{L}^k u_k\| \ll \|\tilde{L}^k u_k\| \ll \mu(\varphi = k+1)\|u\|_{C^\eta(X_k)}$ . This completes the proof of the claim.

Finally, if  $v = u/h$  where  $u$  is Riemann integrable, then we can approximate  $u$  from above and below by Hölder functions  $u^\pm$ . In particular,  $w(n)\tilde{L}^n u^\pm = w(n)hL^n(u^\pm/h) \rightarrow h \int_X u^\pm dm$  uniformly on compact subsets of  $X'$ . By positivity of the transfer operator,  $\tilde{L}^n u^- \leq \tilde{L}^n u \leq \tilde{L}^n u^+$  and so  $h \int_X u^- dm \leq \liminf w(n)\tilde{L}^n u \leq \limsup w(n)\tilde{L}^n u \leq h \int_X u^+ dm$ . Since  $\int_X (u^+ - u^-) dm$  can be made arbitrarily small,  $w(n)L^n v = w(n)h^{-1}\tilde{L}^n u \rightarrow \int_X u dm = \int_X v d\mu$  uniformly on compact subsets of  $X'$ . ■

**Remark 11.9** Suppose that  $v \in L^1(X, \mu)$  and that  $v$  is Hölder on each  $X_k$ . As is evident from the proof, it suffices that  $\sum_{k \geq n} \mu(\varphi = k) \|1_{X_k}(vh)\|_{C^\eta(X_k)} = o(w(n)^{-1})$  for some  $\eta > 0$ . Alternatively, using condition (i) in Theorem 10.4, it suffices that  $\mu(\varphi = n) \|1_{X_n}(vh)\|_{C^\eta(X_n)}$  is summable and  $o(n^{-1})$ .

### 11.3 AFN maps

Zweimüller [44, 45] studied a class of non-Markovian interval maps  $f : X \rightarrow X$ ,  $X = [0, 1]$ , with indifferent fixed points. It is assumed that there is a measurable partition  $\xi$  of  $X$  into open intervals such that  $f$  is  $C^2$  and strictly monotone on each  $Z \in \xi$ , and such that the following conditions are satisfied:

- (A) *Adler's condition:*  $f''/(f')^2$  is bounded on  $\bigcup_{Z \in \xi} Z$ ,
- (F) *Finite images:*  $\{TZ : Z \in \xi\}$  is finite.
- (N) *Nonuniform expansion:* There is a finite set  $\zeta \subset \xi$  such that each interval  $Z \in \zeta$  has an indifferent fixed point  $x_Z$  at one of its endpoints (so  $fx_Z = x_Z$  and  $f'(x_Z) = 1$ ) such that  $f$  has a  $C^1$  extension to  $Z \cup x_Z$  and  $T'$  is increasing (resp. decreasing) on  $Z$  if  $x_Z$  is the left (resp. right) end point of  $Z$ . Moreover,  $|f'| \geq \rho(\epsilon) > 1$  on  $X \setminus \bigcup_{Z \in \zeta} ((x_Z - \epsilon, x_Z + \epsilon) \cap Z)$  for each  $\epsilon > 0$ .

Such a map is called an *AFN map*. A Thaler map (Example 11.7) is an AFN map with full branches. If condition (N) is replaced by

- (U) *Uniform expansion:*  $|f'| \geq \rho > 1$  on  $\bigcup_{Z \in \xi} Z$ ,

then  $f$  is called an *AFU map*.

By the spectral decomposition theorem in [44], any AFN map decomposes into basic sets that are topologically mixing up to a finite cycle. From now on we suppose that  $f : X \rightarrow X$  is a topologically mixing AFN map. Such a map is conservative and exact with respect to Lebesgue measure  $m$ , and admits an equivalent  $\sigma$ -finite invariant measure  $\mu$ . The measure is infinite if and only if  $X$  includes an indifferent fixed point, and we suppose that this is the case. Let  $X' \subset X$  denote the complement of the indifferent fixed points. The density  $h = d\mu/dm$  is of bounded variation, bounded below, and bounded above on compact subsets of  $X'$ .

**Proposition 11.10** *If  $f : X \rightarrow X$  is a topologically mixing AFN map with  $\mu(X) = \infty$ , and  $C$  is a compact subset of  $X'$ , then there exists a first return set  $Y$  with  $\mu(Y) \in (0, \infty)$  such that  $Y$  contains  $C$ , and such that the first return map  $F = f^\varphi : Y \rightarrow Y$  is AFU. Moreover, the Banach space  $\mathcal{B} = BV(Y)$  consisting of bounded variation functions on  $Y$  satisfies hypotheses (H1) and (H2).*

**Proof** By [45, Lemma 8], the first return map  $F$  is AFU. By [36] and [44, Appendix]),  $\mathcal{B} = BV(Y)$  is a suitable Banach space. In particular,  $R(1) : \mathcal{B} \rightarrow \mathcal{B}$  has essential



spectral radius less than 1. The argument in [36] is extended by [5, Proposition 4] who show that there exist constants  $C > 0$ ,  $\tau < (0, 1)$  such that  $\|R(e^{i\theta})^n v\| \leq C(|v|_1 + \tau^n \|v\|)$  for all  $v \in \mathcal{B}$ ,  $\theta \in \mathbb{R}$ ,  $n \geq 1$ . It is easy to extend this argument to cover  $R(z)$  for all  $z \in \mathbb{D}$ . (It should be noted that in our setting, the proof in [5] is greatly simplified since in [5]  $\varphi$  is not assumed to be locally constant and  $F$  is not required to satisfy Adler's condition or finite images.) Since the unit ball in  $\mathcal{B}$  is compact in  $L^1$ , property (1) of Proposition 11.1 again follows from [23]. Property (\*) follows from [5, Theorems 1 and 2]: measurable solutions  $v$  are constant almost everywhere on "recurrent image sets" and there are plenty of such sets by [5, Theorem 3(4)], yielding property (2) of Proposition 11.1. Again, (H1) can be verified en route to the estimate for  $\|R(z)^n v\|$  (the crucial estimate is stated in [5, p. 57, line 10] and is a simple consequence of the AFU structure).  $\blacksquare$

If in addition,  $\mu(y \in Y : \varphi(y) > n) = \ell(n)n^{-\beta}$  is regularly varying with  $\beta \in (0, 1]$  (which includes the case when  $f$  has good asymptotic expansions near each indifferent fixed point as in Example 11.7), then again our main results apply. In particular, for  $\beta \in (\frac{1}{2}, 1]$  it follows that for every BV observable  $v : X \rightarrow \mathbb{R}$  supported on a compact subset of  $X'$ ,  $\lim_{n \rightarrow \infty} w(n)L^n v = \int_X v d\mu$  uniformly on compact subsets of  $X'$ , where  $w(n) = d_\beta^{-1} m(n)n^{1-\beta}$ . Again, we can consider a larger class of observables as in Theorem 1.1.

**Proof of Theorem 1.1** (a) This is identical to the proof of Theorem 11.8 with Hölder replaced by BV. By Theorem 10.4, it suffices to observe that  $\|L^k v_k\| \ll \mu(\varphi = k+1)\|u\|_{BV}$ .

(b) As in the proof of Proposition 10.5, it suffices to prove the  $\geq$  inequality in the lim inf statement. Again, define  $v(k) = \sum_{j=0}^k 1_{X_j} v$ . Choose  $v^- \in \mathcal{B}(X)$  such that  $v \geq v^-$  on  $\bigcup_{j=0}^k X_j$  and define  $v^-(k) = \sum_{j=0}^k 1_{X_j} v^-$ . By Proposition 10.5, for  $k$  fixed,

$$\begin{aligned} \liminf_{n \rightarrow \infty} m(n)n^{1-\beta} L^n v &\geq \liminf_{n \rightarrow \infty} m(n)n^{1-\beta} L^n v^- \geq \liminf_{n \rightarrow \infty} m(n)n^{1-\beta} L^n v^-(k) \\ &= d_\beta \int_X v^-(k) d\mu. \end{aligned}$$

Since  $v$  is Riemann integrable, it follows that  $\liminf_{n \rightarrow \infty} m(n)n^{1-\beta} L^n v \geq \int_X v(k) d\mu$ . Now let  $k \rightarrow \infty$  to complete the proof.

(c) We have  $1_Y L^n v \ll 1_Y L^n(1/h) = \sum_{k=0}^n T_{n-k} L^k(1_{X_k}/h) \ll \sum_{k=0}^n T_{n-k} \tilde{L}^k 1_{X_k}$ , and hence  $\|1_Y L^n v\| \ll \sum_{k=0}^n \|T_{n-k}\| \|\tilde{L}^k 1_{X_k}\| \ll [\{\|T_n\|\} \star \{\mu(\varphi = n)\}]_n \ll \ell(n)n^{-\beta}$ .  $\blacksquare$

**Remark 11.11** (i) The proof of Theorem 1.1(a) shows that the hypotheses are easily generalised just as in Remark 11.9. Suppose that  $v \in L^1(X, \mu)$  and that  $v$  is BV on each  $X_k$ . Then it suffices that  $\sum_{k \geq n} \mu(\varphi = k) \|1_{X_k}(vh)\|_{BV(X_k)} = o(w(n)^{-1})$ . Alternatively, using condition (i) in Theorem 10.4, it suffices that  $\mu(\varphi = n) \|1_{X_n}(vh)\|_{BV(X_n)}$  is summable and  $o(n^{-1})$ .

(ii) For Thaler maps, which are AFN with Gibbs-Markov first return maps, we can work with Hölder or BV norms.

## 11.4 Pomeau-Manneville maps

In this subsection, we verify that our results on second order asymptotics are applicable to certain Pomeau-Manneville intermittency maps, in particular the family (1.1) studied by Liverani *et al.* [31]. Write the invariant measure as  $d\mu = h dm$  where  $m$  is Lebesgue measure and  $h$  is the density.

**Proposition 11.12** *Suppose that  $f : X \rightarrow X$  is given as in (1.1) with  $\beta = 1/\alpha \in (0, 1]$ . Then  $\mu(\varphi > n) = cn^{-\beta} + O(n^{-2\beta})$  where  $c = \frac{1}{4}\beta^\beta h(\frac{1}{2})$ .*

**Proof** First, let  $Y = [\frac{1}{2}, 1]$  with partition sets  $Y_j = \{\varphi = j\}$ . Let  $x_n \in (0, \frac{1}{2}]$  be the sequence with  $x_0 = \frac{1}{2}$  and  $x_n = fx_{n+1}$ , so  $x_n$  is decreasing and  $x_n \rightarrow 0$ . A standard argument shows that  $x_n \sim \frac{1}{2}\beta^\beta n^{-\beta}$  (cf. [37, Corollary 1]). Then  $x_n - x_{n+1} = fx_{n+1} - x_{n+1} = 2^\alpha x^{\alpha+1} = O(n^{-(\beta+1)})$ . Hence  $x_n = \frac{1}{2}\beta^\beta n^{-\beta} + O(n^{-(\beta+1)})$ .

Write  $Y_n = [y_{n-1}, y_{n-2}]$ . Then  $f([\frac{1}{2}, y_n]) = [0, x_n]$ . In particular  $m(\varphi > n) = \frac{1}{2}m([0, x_{n-1}]) = \frac{1}{2}x_{n-1} = m(\varphi > n) = \frac{1}{4}\beta^\beta n^{-\beta} + O(n^{-(\beta+1)})$ .

The density  $h$  is globally Lipschitz on  $(\epsilon, 1]$  for any  $\epsilon > 0$  (see for example [25] or [31, Lemma 2.1]). Hence  $\mu(\varphi > n) = m(\varphi > n)(h(\frac{1}{2}) + O(n^{-\beta}))$ , and the result for  $Y = [\frac{1}{2}, 1]$  follows. The same estimates are obtained by inducing on the set  $Y = [x_q, 1]$  for any fixed  $q \geq 0$ . ■

The next result is immediate by Theorems 9.1 and 9.6.

**Corollary 11.13** *Suppose that  $f : X \rightarrow X$  is given as in (1.1) with  $\beta \in (\frac{1}{2}, 1]$ . Suppose that  $v : [0, 1] \rightarrow \mathbb{R}$  is Hölder or bounded variation supported on a compact subset of  $(0, 1]$ . Let  $m(n) = c$  if  $\beta \in (\frac{1}{2}, 1)$  and  $m(n) = c \log n$  if  $\beta = 1$ . Let  $\gamma = \min\{1 - \beta, \beta - \frac{1}{2}\}$ . Then*

$$m(n)n^{1-\beta}L^n v = d_\beta \int_X v d\mu + O(m(n)^{-1}n^{-\gamma}) \text{ uniformly on compact subsets of } (0, 1].$$

Moreover, if  $\beta \in (\frac{3}{4}, 1]$ , then  $\lim_{n \rightarrow \infty} m(n)n^{1-\beta}\{m(n)n^{1-\beta}L^n v - d_\beta \int_X v d\mu\} = d_{\beta,1} \int_X v d\mu$ , uniformly on compact subsets of  $(0, 1]$ , where typically  $d_{\beta,1} \neq 0$ . ■

Finally, we mention a result on higher order asymptotics for general observables  $v(x) = x^q$ .

**Theorem 11.14** *Suppose that  $f : X \rightarrow X$  is given as in (1.1) with  $\beta \in (\frac{1}{2}, 1]$ . Let  $v(x) = x^q$  where  $\beta(q+1) > 1$ . Then*

$$\begin{aligned} n^{1-\beta}L^n v &= c^{-1}d_\beta \int_X v d\mu + O(n^{-(1-\beta)}, n^{-(\beta-\frac{1}{2})}, n^{-(\beta(q+1)-1)} \log n), \text{ for } \beta \in (\frac{1}{2}, 1), \\ (\log n)L^n v &= c^{-1} \int_X v d\mu + O((\log n)^{-1}), \text{ for } \beta = 1, \end{aligned}$$

uniformly on compact subsets of  $(0, 1]$ .

**Proof** We give the details for  $\beta \in (\frac{1}{2}, 1)$ . By Theorem 9.1,  $T_n = c^{-1}d_\beta n^{-(1-\beta)}P + O(n^{-2(1-\beta)}P) + O(n^{-\frac{1}{2}})$ . Following the proof of Theorem 10.4 with  $v_j = 1_{X_j}$  and  $X_j = (x_j, x_{j-1}]$ ,

$$cd_\beta^{-1}n^{1-\beta}L^n v - \int v = cd_\beta^{-1}n^{1-\beta} \sum_{j=0}^n T_{n-j} L^j v_j - \int v = A - B + C + D,$$

where

$$A = n^{1-\beta} \sum_{j=0}^n \{(n-j)^{-(1-\beta)} - n^{-(1-\beta)}\} \int v_j, \quad B = \sum_{j>n} \int v_j,$$

$$C = O\left(n^{1-\beta} \sum_{j=0}^n (n-j)^{-2(1-\beta)} \int v_j\right), \quad D = O\left(n^{1-\beta} \sum_{j=0}^n (n-j)^{-\frac{1}{2}} \|L^j v_j\|\right).$$

Now  $h(x) \approx x^{-1/\beta}$ , so  $\int v_j d\mu \ll \int_{X_j} x^{q-1/\beta} dx \ll j^{-(\beta q-1)} \mu(X_j) \ll j^{-\beta(q+1)}$ . Hence  $B = O(n^{-(\beta(q+1)-1)})$  and  $C = O(n^{-(1-\beta)})$ . Using the estimate

$$\begin{aligned} (n-j)^{-(1-\beta)} - n^{-(1-\beta)} &= (n-j)^{-(1-\beta)} n^{-(1-\beta)} \{n^{1-\beta} - (n-j)^{1-\beta}\} \\ &= n^{-(1-\beta)} \{(1 + j/(n-j))^{1-\beta} - 1\} \ll n^{-(1-\beta)} (n-j)^{-1} j, \end{aligned}$$

we obtain that  $A \ll \sum_{j=0}^n (n-j)^{-1} j \int v_j \ll \sum_{j=0}^n (n-j)^{-1} j^{-(\beta(q+1)-1)} \ll n^{-(\beta(q+1)-1)} \log n$ .

Now  $h$  is monotone [31, Section 2] and so  $\|h\|_{BV(X_j)} \approx |1_{X_j} h|_\infty$  for each  $j$ . Similarly for  $v$  and we obtain that  $\|vh\|_{BV(X_j)} \leq \|v\|_{BV(X_j)} \|h\|_{BV(X_j)} \ll |1_{X_j} v|_\infty |1_{X_j} h|_\infty = |1_{X_j} v h|_\infty \ll j^{-(\beta q-1)}$ . Hence  $\|L^j v_j\| \ll \|\tilde{L}^j(v_j h)\| \ll j^{-(\beta+1)} \|v_j h\|_{BV(X_j)} \ll j^{-\beta(q+1)}$ . It follows that  $D = O(n^{-(\beta-\frac{1}{2})})$ .  $\blacksquare$

**Acknowledgements** The research of IM and DT was supported in part by EP-SRC Grant EP/F031807/1. We are very grateful to Sébastien Gouëzel and Roland Zweimüller for helpful discussions and encouragement, and to the referees for helpful suggestions.

## References

- [1] J. Aaronson. The asymptotic distributional behaviour of transformations preserving infinite measures. *J. Analyse Math.* **39** (1981) 203–234.
- [2] J. Aaronson. Random  $f$ -expansions. *Ann. Probab.* **14** (1986) 1037–1057.
- [3] J. Aaronson. *An Introduction to Infinite Ergodic Theory*. Math. Surveys and Monographs **50**, Amer. Math. Soc., 1997.

- [4] J. Aaronson and M. Denker. Local limit theorems for partial sums of stationary sequences generated by Gibbs-Markov maps. *Stoch. Dyn.* **1** (2001) 193–237.
- [5] J. Aaronson, M. Denker, O. Sarig and R. Zweimüller. Aperiodicity of cocycles and conditional local limit theorems. *Stoch. Dyn.* **4** (2004) 31–62.
- [6] J. Aaronson, M. Denker and M. Urbański. Ergodic theory for Markov fibred systems and parabolic rational maps. *Trans. Amer. Math. Soc.* **337** (1993) 495–548.
- [7] N. H. Bingham, C. M. Goldie and J. L. Teugels. *Regular variation*. Encyclopedia of Mathematics and its Applications **27**, Cambridge University Press, Cambridge, 1987.
- [8] N. I. Chernov. Markov approximations and decay of correlations for Anosov flows. *Ann. of Math.* **147** (1998) 269–324.
- [9] K. L. Chung. *Markov chains with stationary transition probabilities*. 2nd edition. Die Grundlehren der mathematischen Wissenschaften **104** Springer, New York 1967.
- [10] D. A. Darling and M. Kac. On occupation times for Markoff processes. *Trans. Amer. Math. Soc.* **84** (1957) 444–458.
- [11] M. Denker and M. Urbański. Hausdorff and conformal measures on Julia sets with a rationally indifferent periodic point. *J. London Math. Soc.* **43** (1991) 107–118.
- [12] D. Dolgopyat. On the decay of correlations in Anosov flows. *Ann. of Math.* **147** (1998) 357–390.
- [13] R. A. Doney. One-sided local large deviation and renewal theorems in the case of infinite mean. *Probab. Theory Related Fields* **107** (1997) 451–465.
- [14] E. B. Dynkin. Some limit theorems for sums of independent random variables with infinite mathematical expectations. *Select. Transl. Math. Statist. and Probability, Vol. 1*, Inst. Math. Statist. and Amer. Math. Soc., Providence, R.I., 1961, pp. 171–189.
- [15] W. Feller. Fluctuation theory of recurrent events. *Trans. Amer. Math. Soc.* **67** (1949) 98–119.
- [16] W. Feller. *An Introduction to Probability Theory and its Applications, II*. Wiley, New York, 1966.
- [17] A. Garsia and J. Lamperti. A discrete renewal theorem with infinite mean. *Comment. Math. Helv.* **37** (1962/1963) 221–234.

- [18] S. Gouëzel. Sharp polynomial estimates for the decay of correlations. *Israel J. Math.* **139** (2004) 29–65.
- [19] S. Gouëzel. Berry-Esseen theorem and local limit theorem for non uniformly expanding maps. *Ann. Inst. H. Poincaré Probab. Statist.* **41** (2005) 997–1024.
- [20] S. Gouëzel. Private communication.
- [21] A. B. Hajian and S. Kakutani. Weakly wandering sets and invariant measures. *Trans. Amer. Math. Soc.* **110** (1964) 136–151.
- [22] G. H. Hardy. Some theorems concerning trigonometrical series of special type. *Proc. London Math. Soc.* **32** (1931) 441–448.
- [23] H. Hennion. Sur un théorème spectral et son application aux noyaux lipchitziens. *Proc. Amer. Math. Soc.* **118** (1993) 627–634.
- [24] G. Herglotz. Über Potenzreihen mit positiven reellem Teil im Einheitskreis. *Ber. Verh. Sächs. Akad. Wiss. Leipzig, Math.-Nat. Kl.* **63** (1911).
- [25] H. Hu. Decay of correlations for piecewise smooth maps with indifferent fixed points. *Ergodic Theory Dynam. Systems* **24** (2004) 495–524.
- [26] W. Hurewicz. Ergodic theorem without invariant measure. *Ann. of Math.* **45** (1944) 192–206.
- [27] J. Karamata. Sur un mode to croissance reguliere. Theoremes fondamentaux. *Bull. Soc. Math. France* **61** (1933) 55–62.
- [28] T. Kato. *Perturbation Theory for Linear Operators*. Grundlehren **132**, Springer, New York, 1976.
- [29] J. Lamperti. Some limit theorems for stochastic processes. *J. Math. Mech.* **7** (1958) 433–448.
- [30] C. Liverani. On contact Anosov flows. *Ann. of Math.* **159** (2004) 1275–1312.
- [31] C. Liverani, B. Saussol and S. Vaienti. A probabilistic approach to intermittency. *Ergodic Theory Dynam. Systems* **19** (1999) 671–685.
- [32] K. Petersen. *Ergodic Theory*. Cambridge Studies in Adv. Math. **2**, Cambridge Univ. Press, 1983.
- [33] Y. Pomeau and P. Manneville. Intermittent transition to turbulence in dissipative dynamical systems. *Comm. Math. Phys.* **74** (1980) 189–197.
- [34] H. S. A. Potter. The mean value of a Dirichlet series, II. *Proc. London. Math. Soc.* **47** (1942) 1–19.

- [35] H. L. Royden. *Real Analysis*. Prentice-Hall, New Jersey, 1963.
- [36] M. Rychlik. Bounded variation and invariant measures. *Studia Math.* **76** (1983) 69–80.
- [37] O. M. Sarig. Subexponential decay of correlations. *Invent. Math.* **150** (2002) 629–653.
- [38] M. Thaler. A limit theorem for the Perron-Frobenius operator of transformations on  $[0, 1]$  with indifferent fixed points. *Israel J. Math.* **91** (1995) 111–127.
- [39] M. Thaler. The Dynkin-Lamperti arc-sine laws for measure preserving transformations. *Trans. Amer. Math. Soc.* **350** (1998) 4593–4607.
- [40] M. Thaler. The asymptotics of the Perron-Frobenius operator of a class of interval maps preserving infinite measures. *Studia Math.* **143** (2000) 103–119.
- [41] M. Thaler and R. Zweimüller. Distributional limit theorems in infinite ergodic theory. *Probab. Theory Related Fields* **135** (2006) 15–52.
- [42] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math.* **147** (1998) 585–650.
- [43] L.-S. Young. Recurrence times and rates of mixing. *Israel J. Math.* **110** (1999) 153–188.
- [44] R. Zweimüller. Ergodic structure and invariant densities of non-Markovian interval maps with indifferent fixed points. *Nonlinearity* **11** (1998) 1263–1276.
- [45] R. Zweimüller. Ergodic properties of infinite measure-preserving interval maps with indifferent fixed points. *Ergodic Theory Dynam. Systems* **20** (2000) 1519–1549.