

Superpolynomial and Polynomial Mixing for Semiflows and Flows

Ian Melbourne*

30 September 2017. Revised 3 September 2018

Abstract

We give a review of results on superpolynomial decay of correlations, and polynomial decay of correlations for nonuniformly expanding semiflows and nonuniformly hyperbolic flows. A self-contained proof is given for semiflows. Results for flows are stated without proof (the proofs are contained in separate joint work with Bálint and Butterley). Applications include intermittent solenoidal flows, suspended Hénon attractors, Lorenz attractors and singular hyperbolic attractors, and various Lorenz gas models including the infinite horizon Lorenz gas.

Contents

1	Introduction	2
1.1	Rapid mixing for nonuniformly hyperbolic flows	4
1.2	Statistical limit laws for time-one maps	6
1.3	Polynomial mixing for nonuniformly hyperbolic flows	7
I	Statement of results	7
2	Preliminaries on suspensions	8
3	Mixing rates for nonuniformly expanding semiflows	9
3.1	Gibbs-Markov semiflows	9
3.1.1	Approximate eigenfunctions	10
3.1.2	Observables on Y^φ	10
3.1.3	Decay of correlations	11
3.2	Nonuniformly expanding semiflows on metric spaces	11
3.3	Dynamically Hölder semiflows and observables	14
4	Mixing rates for nonuniformly hyperbolic flows	15
4.1	Skew product Gibbs-Markov flows	15
4.2	General nonuniformly hyperbolic flows	16

*Mathematics Institute, University of Warwick, Coventry, CV4 7AL, UK

5	Criteria for absence of approximate eigenfunctions	17
5.1	Diophantine condition on periods	18
5.2	Good asymptotics	18
5.3	Temporal distance function	19
II	Proof of theorems for semiflows	22
6	Strategy for obtaining rates of mixing	22
6.1	Laplace transforms and inversion formulas	22
6.2	Formula for $\hat{\rho}_{v,w}$	23
7	Proof of rapid mixing for semiflows	25
7.1	Background on Gibbs-Markov maps	25
7.2	Smoothness of $R\hat{V} : \mathcal{F}_\theta(Y^\varphi) \rightarrow \mathcal{F}_\theta(Y)$	26
7.3	Smoothness of $\hat{R} : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$ and spectral properties	27
7.4	Dolgopyat estimate	29
7.5	Smoothness of $\hat{T} = (I - \hat{R})^{-1} : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$	32
7.6	Proof of Theorem 3.7	33
8	Proof of polynomial mixing for semiflows	33
8.1	Some conventions	34
8.2	Refined estimates for $J_0, w_0, R\hat{V}$ and \hat{R}	35
8.3	Further estimates for dealing with the singularity at zero	40
8.4	Truncation	41
8.5	Small b : Proof of Lemma 8.22	44
8.6	Large b : Proof of Lemma 8.23	46
9	Some open questions	47

1 Introduction

Let (Λ, μ_Λ) be a probability space. Given a measure-preserving flow $T_t : \Lambda \rightarrow \Lambda$ and observables $v, w \in L^2(\Lambda)$, we define the correlation function $\rho_{v,w}(t) = \int_\Lambda v w \circ T_t d\mu_\Lambda - \int_\Lambda v d\mu_\Lambda \int_\Lambda w d\mu_\Lambda$. The flow is *mixing* if $\lim_{t \rightarrow \infty} \rho_{v,w}(t) = 0$ for all $v, w \in L^2(\Lambda)$.

Of interest is the rate of decay of correlations, or rate of mixing, namely the rate at which $\rho_{v,w}$ converges to zero. For nontrivial mixing flows, the decay rate is arbitrarily slow for L^2 observables. Hence the aim is to establish decay rates under regularity hypotheses on the flow T_t , the measure μ_Λ , and the observables v, w .

Consider a mixing uniformly hyperbolic (Axiom A) basic set $\Lambda \subset M$ for a smooth flow $T_t : M \rightarrow M$ with invariant probability measure μ_Λ that is an equilibrium state for a Hölder potential [24]. Bowen & Ruelle [24] asked whether Λ is exponentially mixing ($\rho_{v,w}(t) = O(e^{-ct})$ for some $c > 0$) for sufficiently regular v, w . (In the discrete time case, it is well-known that mixing Axiom A diffeomorphisms are exponentially mixing.) However, Pollicott [73] and Ruelle [78] showed that mixing Axiom A flows need not mix exponentially, and Pollicott [74] showed that the decay rates could be arbitrarily slow.

Building upon results of Chernov [39], Dolgopyat [47] showed that geodesic flows on compact surfaces of negative curvature are exponentially mixing. Liverani [59] extended this result to arbitrary dimensional geodesic flows in negative curvature and more generally to contact Anosov flows. Subsequent developments include [5, 10, 11, 13, 32, 75, 81, 82, 83]. However, exponential mixing remains poorly understood in general. Two major open questions, even for Anosov flows, are

- Is exponential mixing typical in any reasonable sense?
- Does mixing imply exponential mixing rate?

Indeed, it is only recently that the first robust examples of exponentially mixing Axiom A flows [3] and Anosov flows [33] have been obtained.

Dolgopyat [48] introduced the weaker notion of *rapid mixing* (superpolynomial decay of correlations) where $\rho_{v,w}(t) = O(t^{-q})$ for sufficiently regular observables for any fixed $q \geq 1$, and showed that rapid mixing is ‘prevalent’ for Axiom A flows: it suffices that the flow contains two periodic solutions with periods whose ratio is Diophantine. In addition, Dolgopyat [47] showed that for Anosov flows, joint nonintegrability of the stable and unstable foliations (an open and dense condition by Brin [25, 26]) implies rapid mixing. Field *et al.* [50] introduced the notion of *good asymptotics* and used this to prove that amongst C^r Axiom A flows, $r \geq 2$, an open and dense set of flows is rapid mixing.

The topic of this review article is the extension of the rapid mixing method to the nonuniformly hyperbolic setting. In [63, 64], we obtained results on rapid mixing and polynomial mixing for nonuniformly hyperbolic semiflows and flows, combining the rapid mixing method of Dolgopyat [48] with advances by Young [85, 86] in the discrete time setting. The purpose of this review article is twofold. In Part I, we state the results in a way that incorporates applications that have arisen since [63, 64]. (This should simplify the exposition in future applications avoiding complicated referencing of the type in for example [8, Section 4]). In Part II, we provide complete self-contained proofs for semiflows. The proofs for flows are given in a separate joint paper with Péter Bálint & Oliver Butterley [14].

In Subsections 1.1 and 1.3 below, we summarize the applications discussed in [63, 64] as well as subsequent applications to Lorentz gases with cusps and in flowers, higher-dimensional Lorentz gases, Lorenz attractors, and intermittent solenoidal flows.

The treatment here (and in [14]) differs in two main respects from [63, 64]. The differences apply equally to flows and semiflows so we just mention flows here. First, flows are viewed as suspensions over a uniformly hyperbolic map with a possibly unbounded roof function (rather than as suspensions over a uniformly or nonuniformly hyperbolic map with a bounded roof function). This simplifies the analysis since it suffices to consider twisted transfer operators with one complex parameter, whereas two complex parameters were required in [63]. A consequence of this is that the number of periodic orbits in the Diophantine condition in Section 5.1 reduces from 4 to 3. (On the other hand, the methods in [63, 64] can be applied also to toral extensions of nonuniformly expanding maps, see [12, 67], and the results here cannot.) Second, [48, 63, 64] use analyticity properties of the Laplace transform of $\rho_{v,w}$ in the complex plane, whereas we use smoothness on the imaginary axis incorporating ideas developed more recently with Dalia Terhesiu, especially [66]. One advantage is that Theorem 3.7 on rapid mixing works for roof functions that lie in L^p for all $p \geq 1$, whereas [63] requires an exponential tails condition. Another advantage is that the

estimate in Lemma 8.20 is much cleaner than the corresponding estimate in [64]. (Also, the approach here leads to an estimate in Lemma 8.23 that is much better than might have been expected.)

In addition, we resolve two issues that were overlooked in [64]:

- (i) The Dolgopyat argument focuses on controlling high frequencies in the Laplace transform of the correlation function $\rho_{v,w}$ but it is also necessary to deal with the singularity at 0. This requires extra arguments in the slowly mixing cases.
- (ii) The hypotheses in [64] for passing from semiflows to flows are too strong for the intended applications where there is often slow contraction along stable leaves.

Issue (i) is covered in the proofs for semiflows in Part II of this article. Regarding point (ii), similar problems have arisen even for discrete time dynamical systems. Various polynomial mixing results in [40, 42, 61] described in Subsection 1.3 rely on a result of [86] which is formulated only for noninvertible dynamical systems. The extra argument required for passing to invertible systems is due to Gouëzel [54] based on work of [34], and can be found in [65, Appendix B] and [58, Theorem 2.10]. Resolving the corresponding problem for flows turns out to be more subtle (even formulating the correct hypotheses is difficult), and is completed in Bálint *et al.* [14].

In the remainder of the introduction, we give an overview of rapid and polynomial mixing, as well as consequent statistical properties, for various classes of flows.

1.1 Rapid mixing for nonuniformly hyperbolic flows

Young [85] established exponential mixing for a large class of nonuniformly hyperbolic maps modelled by Young towers with exponential tails, including dispersing billiards and Hénon-like maps [23]. In [63], we extended the ideas of Dolgopyat [48] to the corresponding class of nonuniformly hyperbolic flows where there is a suitable Poincaré map modelled by a Young tower with exponential tails. Roughly speaking, one of the main results described in this review article is that

A ‘prevalent’ set of nonuniformly hyperbolic flows with exponential tails are rapid mixing.

Rapid mixing is established for sufficiently regular observables, and prevalence is understood in terms of a Diophantine condition on the periodic data and also in terms of the good asymptotics condition [50] which is open and dense.

Planar Lorentz gases A rich class of examples of three-dimensional nonuniform hyperbolic flows is provided by planar Lorentz gas models [80, 37]. See [38] for a survey of results about these models. From now on, all Lorentz gases are planar unless stated otherwise.

In particular, the periodic Lorentz gas is a billiard flow on $\mathbb{T}^2 \setminus \Omega$ where Ω is a disjoint union of convex regions with C^3 boundaries. (The phase-space of the flow is three-dimensional; planar position and direction.) The flow has a natural global cross-section corresponding to collisions and the Poincaré map is called the *billiard map*. The Lorentz flow satisfies the *finite horizon* condition if the time between collisions is uniformly bounded. For the billiard map associated to the finite horizon Lorentz gas, Bunimovich, Sinaï & Chernov [31] proved stretched exponential mixing rates, and exponential mixing was established

by Young [85]. Chernov [35] extended Young’s method to prove exponential mixing for billiard maps corresponding to (i) periodic Lorentz gases with infinite horizon, (ii) periodic Lorentz gases with external forcing, and (iii) Lorentz gases in bounded domains with sides that are convex inwards and corner points with positive angles. (For the third case, billiards with corners, a technical condition in [35] was removed in [44].) For the corresponding flows, we have:

Finite horizon planar periodic Lorentz flows, and planar Lorentz flows with cusps or corner points, are rapid mixing. A prevalent set of externally forced finite horizon planar periodic Lorentz flows are rapid mixing.

Except for the flows with cusps, this is a consequence of [63]. (When there is no external forcing, the contact structure ensures that the prevalence assumption is unnecessary [64], see Remark 5.7.)

The case of cusps is treated in [17]. Here, the billiard map mixes only at the rate $O(n^{-1})$ so [63] does not apply directly. However, collisions within a cusp occur in quick succession so it is reasonable to expect much quicker, possibly even exponential, mixing in continuous time. As shown in [17], the billiard map on the ‘natural’ cross-section can be replaced by a Poincaré map (with cross-section bounded away from the cusps) that is modelled by a Young tower with exponential tails in such a way that [63] applies. The same idea applies to a class of billiards called Bunimovich flowers [29] (see also [17, Section 3.1] for a precise description). It was shown in [40] that the billiard map mixes at roughly the rate $O(n^{-2})$. By [17],

For Bunimovich flowers, the corresponding Lorentz flow is rapid mixing.

For the finite horizon planar periodic Lorentz flow, Chernov [36] proved stretched exponential mixing, and exponential mixing was finally achieved in the recent paper of Baladi, Demers & Liverani [11]. The methods of [11, 36] rely crucially on the contact structure. It is reasonable to expect that these methods also apply to the Lorentz flows with corners and cusps, as well as the Bunimovich flower examples, though the technical details have not been verified at the time of writing. However, it seems unlikely that the result on prevalent rapid mixing for externally forced Lorentz flows [63] will be improved upon in the near future — the lack of contact structure and the irregularity of the foliations means that the current technology for proving (stretched) exponential mixing is insufficient.

Once again, we emphasize that the rapid mixing results above are proved for observables which are smooth along the flow direction. Unfortunately this excludes certain important physical observables such as position and velocity. In contrast, the results in [11, 36] cover all Hölder (and dynamically Hölder) observables, including position and velocity, but the methods apply less generally.

For higher-dimensional periodic billiards, there is a technical and unresolved problem concerning “growth of complexity” [15]. It is conjectured that growth of complexity is typically bounded, but there are no examples where the growth is known to be subexponential and there are counterexamples where the growth of complexity is exponential [20]. Bálint & Tóth [19] proved that this is the only obstruction, obtaining the following conditional result: For higher-dimensional Lorentz gases, if the growth of complexity is subexponential,

then the billiard map is modelled by a Young tower with exponential tails and in particular is exponentially mixing. Hence by [63],

Higher-dimensional periodic Lorentz gas flows with subexponential growth of complexity are rapid mixing.

Flows near homoclinic tangencies Benedicks & Carleson [21] studied the Hénon map $T_{a,b}(x, y) = (1 - ax^2 + y, bx)$ and proved the existence of a strange attractor for a positive measure set of parameters a, b . The attractor admits an SRB measure [22] and is exponentially mixing by Benedicks & Young [23]. Mora & Viana [71] showed that Hénon-like attractors arise for positive measure sets of parameters in the unfoldings of homoclinic tangencies for surface diffeomorphisms and this was extended to higher dimensions by [84, 46]. By [63],

Positive measure sets of flows near a homoclinic tangency are rapid mixing.

Again, (stretched) exponential mixing is beyond the current technology.

Lorenz attractors and singular hyperbolic attractors The classical Lorenz attractor is exponentially mixing [5], but the proof relies on the smoothness of the stable foliation for the flow [6, 8]. General Lorenz attractors (and singular hyperbolic attractors) have a Hölder stable foliation [6] but this foliation need not be smooth. Building upon [9], it is shown in [7] that an open and dense set of Lorenz attractors (and codimension two singular hyperbolic attractors) are rapid mixing even when the stable foliation is not smooth.

1.2 Statistical limit laws for time-one maps

In situations where we obtain rapid mixing for a nonuniformly hyperbolic flow, certain statistical properties such as the central limit theorem (CLT) and almost sure invariance principle (ASIP) hold for the time-one map of the flow. (Such properties for the flow itself are much simpler since they are inherited from the statistical properties of the Poincaré map and no mixing properties are required [77, 45, 69].)

In particular, given any of the rapidly mixing flows described in Subsection 1.1 and a mean zero observable $v : \Lambda \rightarrow \mathbb{R}$ sufficiently smooth in the flow direction, setting $v_n = \sum_{j=0}^{n-1} v \circ T_j$, the following hold by [8]:

- (a) (CLT) $n^{-1/2}v_n \rightarrow_d N(0, \sigma^2)$ where $\sigma^2 = \lim_{n \rightarrow \infty} n^{-1} \int_{\Lambda} v_n^2 d\mu_{\Lambda} = \sum_{n=-\infty}^{\infty} \int_{\Lambda} v v \circ T_n d\mu_{\Lambda}$.
- (b) (Nondegeneracy) If $\sigma^2 = 0$, then for every periodic orbit q there exists $t_q > 0$ such that $\int_0^{t_q} v(T_t q) dt = 0$. (The value of t_q is independent of v but in general is not directly related to the period of q . A formula for t_q is given in Remark 5.2.)
- (c) (ASIP) Passing to an enriched probability space, there exists a sequence X_0, X_1, \dots of iid normal random variables with mean 0 and variance σ^2 such that $v_n = \sum_{j=0}^{n-1} X_j + O(n^{1/4}(\log n)^{1/2}(\log \log n)^{1/4})$ a.e.

The proof uses [43] to obtain the ASIP for time-one maps of rapidly mixing semiflows and then passes as in [68] from the semiflow to the flow. See also [68] for the CLT. The ASIP has various consequences including the CLT and law of the iterated logarithm as well as their functional versions. For a more complete list of consequences see [72].

1.3 Polynomial mixing for nonuniformly hyperbolic flows

Young [86] established polynomial mixing rates for a large class of nonuniformly expanding maps including intermittent maps of Pomeau-Manneville type [76]. We consider the corresponding class of nonuniformly hyperbolic flows with Poincaré map modelled by a polynomially mixing Young tower. Roughly speaking, the second main result in this review article is that for $q > 0$,

Amongst nonuniformly hyperbolic flows for which the Poincaré map has mixing rate $O(n^{-q})$, a prevalent set have mixing rate $O(t^{-q})$.

Here, prevalent has the same meaning as in Subsection 1.1. This result applies in particular to intermittent solenoidal flows (Example 4.2).

Remark 1.1 The statistical properties described in Subsection 1.2 for rapid mixing flows hold also for flows that decay at rate $O(t^{-q})$ for q large enough. The CLT requires only that $q > 1$. For the ASIP with error term $O(n^{1/4}(\log n)^{1/2}(\log \log n)^{1/4})$, it suffices that $q > 2\sqrt{2}$ (see [8, Theorem 5.2]).

Again, planar Lorentz gases provide a rich source of examples. For the billiard maps, Markarian [61] combined the work of [35, 86] to prove the mixing rate $O(n^{-1})$ for Bunimovich stadia [30], and the method was extended by Chernov & Zhang [40, 42] to obtain the rate $O(n^{-1})$ for semidispersing billiards (rectangular tables with at least one convex obstacle). Our results apply to the corresponding flows (again the contact structure ensures that there is no prevalence restriction).

It has been conjectured (especially for the infinite horizon Lorentz gas flow in [51, 62]) that the decay rate for the flow is $O(t^{-1})$. (An elementary argument in [16] shows that this rate is optimal; see [14, Proposition 9.14].) The conjectured decay rate $O(t^{-1})$ is proved in [14]:

Lorentz flows in Bunimovich stadia, semidispersing Lorentz flows, and infinite horizon planar periodic Lorentz gas flows have mixing rate $O(t^{-1})$.

Notation We use the “big O ” and \ll notation interchangeably, writing $a_n = O(b_n)$ or $a_n \ll b_n$ if there is a constant $C > 0$ such that $a_n \leq Cb_n$ for all $n \geq 1$. As usual, $a_n \sim b_n$ means that $\lim_{n \rightarrow \infty} a_n/b_n = 1$. There are various “universal” constants $C_1, \dots, C_5 \geq 1$ depending only on the flow that do not change throughout the article.

Part I

Statement of results

In this part of the review article, we describe in detail the main results on rapid and polynomial mixing for nonuniformly expanding semiflows and nonuniformly hyperbolic flows. In Section 2, we recall the notion of a suspension (semi)flow, as well as associated notation. Sections 3 and 4 contain the main results for semiflows and flows respectively. These rely on a technical condition, absence of approximate eigenfunctions (Definition 3.4). In Section 5, we discuss criteria ensuring that this condition holds.

2 Preliminaries on suspensions

It is standard to model flows and semiflows as suspensions. Here, we review the basic definitions and notation that will be used throughout the review article. For brevity we speak of semiflows, even when some of them are flows.

Suspension semiflows in their own right Let (Y, μ) be a probability space and let $F : Y \rightarrow Y$ be a measure-preserving transformation. Let $\varphi : Y \rightarrow \mathbb{R}^+$ be an integrable roof function. Define the suspension $Y^\varphi = \{(y, u) \in Y \times [0, \infty) : u \in [0, \varphi(y)]\} / \sim$ where $(y, \varphi(y)) \sim (Fy, 0)$. The suspension semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$ is given by $F_t(y, u) = (y, u + t)$ computed modulo identifications. We obtain an F_t -invariant probability measure on Y^φ given by $\mu^\varphi = \mu \times \text{Lebesgue} / \int_Y \varphi d\mu$.

Suspension semiflows as models for ambient semiflows Let $T_t : \Lambda \rightarrow \Lambda$ be a semiflow defined on an ambient measurable space Λ . (We assume that $(x, t) \mapsto T_t x$ is measurable from $\Lambda \times [0, \infty) \rightarrow \Lambda$.) Let $Y \subset \Lambda$ be a measurable subset with probability measure μ . We suppose that $F : Y \rightarrow Y$ is a measure-preserving transformation and $\varphi : Y \rightarrow \mathbb{R}^+$ is an integrable function such that $T_{\varphi(y)} y = Fy$ for all $y \in Y$. Then we can form the suspension semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$ with invariant probability measure μ^φ .

Define $\pi : Y^\varphi \rightarrow \Lambda$ by $\pi(y, u) = T_u y$. This is well-defined since $\pi(y, \varphi(y)) = T_{\varphi(y)} y = Fy = \pi(Fy, 0)$, and defines a measurable semiconjugacy from F_t to T_t . Hence the measure $\mu_\Lambda = \pi_* \mu^\varphi$ is a T_t -invariant probability measure on Λ .

Given observables $v, w \in L^2(\Lambda)$, define $\tilde{v} = v \circ \pi$, $\tilde{w} = w \circ \pi$. Then $\tilde{v}, \tilde{w} \in L^2(Y^\varphi)$ and

$$\int_\Lambda v w \circ T_t d\mu_\Lambda - \int_\Lambda v d\mu_\Lambda \int_\Lambda w d\mu_\Lambda = \int_{Y^\varphi} \tilde{v} \tilde{w} \circ F_t d\mu^\varphi - \int_{Y^\varphi} \tilde{v} d\mu^\varphi \int_{Y^\varphi} \tilde{w} d\mu^\varphi. \quad (2.1)$$

Hence rates of mixing for the ambient semiflow T_t and observables v, w on (Λ, μ_Λ) reduces to rates of mixing for the suspension semiflow F_t and observables \tilde{v}, \tilde{w} on (Y^φ, μ^φ) .

Remark 2.1 For the systems considered in this article, the measure μ on Y is ergodic by construction, and the form of F ensures ergodicity of μ^φ and hence μ_Λ .

Remark 2.2 Throughout this review article we suppose that $\inf \varphi > 0$, though some of the results, particularly Theorems 3.6 and 3.7 below, do not require this. In any case, this condition is not very restrictive — if $\inf \varphi = 0$, then this can be circumvented either by shrinking the cross-section Y or by waiting for a later return to Y (see for example [17]).

3 Mixing rates for nonuniformly expanding semiflows

In this section, we review results on rapid mixing and polynomial mixing for nonuniformly expanding semiflows. In Subsection 3.1, we define a class of Gibbs-Markov semiflows and state the main results for such semiflows. In Subsection 3.2, we consider nonuniformly expanding Hölder semiflows on an ambient metric space M , with Hölder observables, and show how these reduce to the situation in Subsection 3.1. As an application we consider intermittent semiflows. In Subsection 3.3, we generalise to dynamically Hölder semiflows and observables.

3.1 Gibbs-Markov semiflows

In this subsection, we consider a class of Gibbs-Markov semiflows built as suspensions over Gibbs-Markov maps. Standard references for background material on Gibbs-Markov maps are [1, Chapter 4] and [2].

Suppose that (Y, μ) is a probability space with an at most countable measurable partition $\{Y_j, j \geq 1\}$ and let $F : Y \rightarrow Y$ be a measure-preserving transformation. For $\theta \in (0, 1)$, define $d_\theta(y, y') = \theta^{s(y, y')}$ where the *separation time* $s(y, y')$ is the least integer $n \geq 0$ such that $F^n y$ and $F^n y'$ lie in distinct partition elements in $\{Y_j\}$. It is assumed that the partition $\{Y_j\}$ separates trajectories, so $s(y, y') = \infty$ if and only if $y = y'$. Then d_θ is a metric, called a *symbolic metric*.

A function $v : Y \rightarrow \mathbb{R}$ is d_θ -Lipschitz if $|v|_\theta = \sup_{y \neq y'} |v(y) - v(y')|/d_\theta(y, y')$ is finite. Let $\mathcal{F}_\theta(Y)$ be the Banach space of Lipschitz functions with norm $\|v\|_\theta = |v|_\infty + |v|_\theta$.

More generally (and with a slight abuse of notation), we say that a function $v : Y \rightarrow \mathbb{R}$ is *piecewise d_θ -Lipschitz* if $|1_{Y_j} v|_\theta = \sup_{y, y' \in Y_j, y \neq y'} |v(y) - v(y')|/d_\theta(y, y')$ is finite for all j . If in addition, $\sup_j |1_{Y_j} v|_\theta < \infty$ then we say that v is *uniformly piecewise d_θ -Lipschitz*. Note that such a function v is bounded on partition elements but need not be bounded on Y .

Definition 3.1 A measure-preserving transformation map $F : Y \rightarrow Y$ is called a (*full branch*) *Gibbs-Markov map* if

- $F|_{Y_j} : Y_j \rightarrow Y$ is a measurable bijection for each $j \geq 1$, and
- The potential function $p = \log \frac{d\mu}{d\mu \circ F} : Y \rightarrow \mathbb{R}$ is uniformly piecewise d_θ -Lipschitz for some $\theta \in (0, 1)$.

Definition 3.2 A suspension semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$ is called a *Gibbs-Markov semiflow* if there exist constants $C_1 \geq 1$, $\theta \in (0, 1)$ such that $F : Y \rightarrow Y$ is a Gibbs-Markov map, $\varphi : Y \rightarrow \mathbb{R}^+$ is an integrable roof function with $\inf \varphi > 0$, and

$$|1_{Y_j} \varphi|_\theta \leq C_1 \inf_{Y_j} \varphi \quad \text{for all } j \geq 1. \quad (3.1)$$

It follows that $\sup_{Y_j} \varphi \leq 2C_1 \inf_{Y_j} \varphi$ for all $j \geq 1$. As mentioned in Remark 2.2 the assumption $\inf \varphi > 0$ is not really needed, though condition (3.1) would need to be changed to $|1_{Y_j} \varphi|_\theta \leq C_1 (\inf_{Y_j} \varphi + 1)$ to avoid being too restrictive (similarly, in the definition of $\mathcal{F}_\theta(Y^\varphi)$ below), and this would result in more complicated formulas throughout.

3.1.1 Approximate eigenfunctions

For $b \in \mathbb{R}$, we define the operators

$$M_b : L^\infty(Y) \rightarrow L^\infty(Y), \quad M_b v = e^{ib\varphi} v \circ F.$$

Evidently, $L^\infty(Y)$ here denotes complex-valued essentially bounded measurable functions on Y . As is standard, we often pass to complexifications of various Banach spaces without comment, for example when discussing spectral properties. In particular, the functions in $F_\theta(Y)$ in Definition 3.4 below are complex-valued.

Definition 3.3 A subset $Z_0 \subset Y$ is a *finite subsystem* of Y if $Z_0 = \bigcap_{n \geq 0} F^{-n} Z$ where Z is the union of finitely many elements from the partition $\{Y_j\}$. (Note that $F|_{Z_0} : Z_0 \rightarrow Z_0$ is a full one-sided shift on finitely many symbols.)

Definition 3.4 We say that M_b has *approximate eigenfunctions* on a subset $Z_0 \subset Y$ if for any $\alpha_0 > 0$, there exist constants $\alpha, \xi > \alpha_0$ and $C > 0$, and sequences $|b_k| \rightarrow \infty$, $\psi_k \in [0, 2\pi)$, $u_k \in \mathcal{F}_\theta(Y)$ with $|u_k| \equiv 1$ and $|u_k|_\theta \leq C|b_k|$, such that setting $n_k = \lceil \xi \ln |b_k| \rceil$,

$$|(M_{b_k}^{n_k} u_k)(y) - e^{i\psi_k} u_k(y)| \leq C|b_k|^{-\alpha} \quad \text{for all } y \in Z_0, k \geq 1. \quad (3.2)$$

Remark 3.5 For brevity, the statement ‘‘Assume absence of approximate eigenfunctions’’ is the assumption that there exists at least one finite subsystem Z_0 such that M_b does not have approximate eigenfunctions on Z_0 .

3.1.2 Observables on Y^φ

Given $v : Y^\varphi \rightarrow \mathbb{R}$ and $\theta \in (0, 1)$, we define

$$|v|_\theta = \sup_{(y,u),(y',u) \in Y^\varphi, y \neq y'} \frac{|v(y,u) - v(y',u)|}{\varphi(y)d_\theta(y,y')}, \quad \|v\|_\theta = |v|_\infty + |v|_\theta.$$

Let $\mathcal{F}_\theta(Y^\varphi)$ be the space of observables $v : Y^\varphi \rightarrow \mathbb{R}$ with $\|v\|_\theta < \infty$.

Also, for $\eta \in (0, 1]$, we define

$$|v|_{\infty,\eta} = \sup_{(y,u),(y,u') \in Y^\varphi, u \neq u'} \frac{|v(y,u) - v(y,u')|}{|u - u'|^\eta}, \quad \|v\|_{\theta,\eta} = \|v\|_\theta + |v|_{\infty,\eta}.$$

(Here and elsewhere, $|u - u'|$ denotes absolute value, with u, u' regarded as elements of $[0, \infty)$.) Let $\mathcal{F}_{\theta,\eta}(Y^\varphi)$ be the space of observables $v : Y^\varphi \rightarrow \mathbb{R}$ with $\|v\|_{\theta,\eta} < \infty$.

We say that $w : Y^\varphi \rightarrow \mathbb{R}$ is *differentiable in the flow direction* if the limit $\partial_t w = \lim_{t \rightarrow 0} (w \circ F_t - w)/t$ exists pointwise. Note that $\partial_t w = \frac{\partial w}{\partial u}$ on the set $\{(y,u) : y \in Y, 0 < u < \varphi(y)\}$. For $m \geq 0$, let w be m -times differentiable in the flow direction and define $|w|_{\infty,m} = \sum_{j=0}^m |\partial_t^j w|_\infty$. Let $L^{\infty,m}(Y^\varphi)$ be the space of observables $w : Y^\varphi \rightarrow \mathbb{R}$ with $|w|_{\infty,m} < \infty$.

3.1.3 Decay of correlations

We can now state the main results for Gibbs-Markov semiflows.

Theorem 3.6 *Let $F_t : Y^\varphi \rightarrow Y^\varphi$ be a Gibbs-Markov semiflow such that $\mu(\varphi > t) = O(t^{-\beta})$ for some $\beta > 1$. Assume absence of approximate eigenfunctions.*

Then for any $\eta \in (0, 1]$, there exists $m \geq 1$, $\theta \in (0, 1)$ and $C > 0$ such that

$$|\rho_{v,w}(t)| \leq C \|v\|_{\theta,\eta} \|w\|_{\infty,m} t^{-(\beta-1)} \quad \text{for all } v \in \mathcal{F}_{\theta,\eta}(Y^\varphi), w \in L^{\infty,m}(Y^\varphi), t > 1.$$

For rapid mixing, we obtain a slightly stronger result where v lies in $\mathcal{F}_\theta(Y^\varphi)$ (instead of $\mathcal{F}_{\theta,\eta}(Y^\varphi)$). For convenience, we state the result on rapid mixing separately.

Theorem 3.7 *Let $F_t : Y^\varphi \rightarrow Y^\varphi$ be a Gibbs-Markov semiflow such that $\varphi \in L^q(Y)$ for all $q \geq 1$. Assume absence of approximate eigenfunctions.*

Then F_t is rapid mixing: for any $q \in \mathbb{N}$, there exists $m \geq 1$, $\theta \in (0, 1)$ and $C > 0$ such that

$$|\rho_{v,w}(t)| \leq C \|v\|_{\theta} \|w\|_{\infty,m} t^{-q} \quad \text{for all } v \in \mathcal{F}_\theta(Y^\varphi), w \in L^{\infty,m}(Y^\varphi), t > 1.$$

The proofs of Theorems 3.6 and 3.7 are given in Sections 8 and 7 respectively.

Theorem 3.6 is sharp. Precise asymptotics and error rates hold for “nicely” supported observables:

Theorem 3.8 ([66, Theorem 2.4(a)]) *Assume the setup of Theorem 3.6 and suppose further that $\text{supp } v, \text{supp } w \in Y \times [0, \inf \varphi]$. For $\beta \in (1, 2)$, set $s = 2(\beta - 1)$. For $\beta \geq 2$, choose any $s < \beta$. Then*

$$\rho_{v,w}(t) = |\varphi|_1^{-1} \int_{Y^\varphi} v d\mu^\varphi \int_{Y^\varphi} w d\mu^\varphi \int_t^\infty \mu(\varphi > t') dt' + O(\|v\|_{\theta} \|w\|_{\infty,m} t^{-s}).$$

In particular, if $\int_t^\infty \mu(\varphi > t') dt' \sim ct^{-(\beta-1)}$ for some $c > 0$, then

$$\rho_{v,w}(t) \sim c |\varphi|_1^{-1} \int_{Y^\varphi} v d\mu^\varphi \int_{Y^\varphi} w d\mu^\varphi t^{-(\beta-1)} \quad \text{as } t \rightarrow \infty. \quad \blacksquare$$

We also mention a result which is the continuous time analogue of [53, Theorem 1.3, last statement].

Theorem 3.9 ([66, Theorem 2.4(b)]) *Assume the setup of Theorem 3.8. If in addition $\int_{Y^\varphi} v d\mu^\varphi = 0$ or $\int_{Y^\varphi} w d\mu^\varphi = 0$, then $|\rho_{v,w}(t)| \leq C \|v\|_{\theta} \|w\|_{\infty,m} t^{-(\beta-\epsilon)}$ for all $\epsilon > 0$. \blacksquare*

3.2 Nonuniformly expanding semiflows on metric spaces

Let $T_t : M \rightarrow M$ be a semiflow defined on a metric space (M, d) with $\text{diam } M \leq 1$. Fix $\eta \in (0, 1]$.

Given $v : M \rightarrow \mathbb{R}$, define $|v|_{C^\eta} = \sup_{x \neq x'} |v(x) - v(x')|/d(x, x')^\eta$ and $\|v\|_{C^\eta} = |v|_\infty + |v|_{C^\eta}$. Let $C^\eta(M) = \{v : M \rightarrow \mathbb{R} : \|v\|_{C^\eta} < \infty\}$. Also, define $|v|_{C^{0,\eta}} = \sup_{x \in M, t > 0} |v(T_t x) - v(x)|/t^\eta$ and let $C^{0,\eta}(M) = \{v : M \rightarrow \mathbb{R} : |v|_\infty + |v|_{C^{0,\eta}} < \infty\}$. (Such observables are Hölder in the flow direction.)

We say that $w : M \rightarrow \mathbb{R}$ is differentiable in the flow direction if the limit $\partial_t w = \lim_{t \rightarrow 0} (w \circ T_t - w)/t$ exists pointwise. Define $|w|_{\infty, m} = \sum_{j=0}^m |\partial_t^j w|_{\infty}$ and let $L^{\infty, m}(M) = \{w : M \rightarrow \mathbb{R} : |w|_{\infty, m} < \infty\}$.

Let $X \subset M$ be a Borel subset and define $C^\eta(X)$ using the metric d restricted to X . We suppose that $T_{h(x)}x \in X$ for all $x \in X$, where $h : X \rightarrow \mathbb{R}^+$ lies in $C^\eta(X)$ and $\inf h > 0$. In addition, we suppose that there exists $C > 0$ such that

$$d(T_t x, T_t x') \leq C d(x, x')^\eta \quad \text{for all } t \in [0, |h|_{\infty}], x, x' \in M. \quad (3.3)$$

Define $f : X \rightarrow X$ by $fx = T_{h(x)}x$. We assume that f is modelled by a Young tower with polynomial tails [86]. This means that there is a Borel subset $Y \subset X$ with finite Borel measure μ_0 and a return time $\tau : Y \rightarrow \mathbb{Z}^+$ with $\mu_0(\tau > n) = O(n^{-\beta})$ for some $\beta > 1$ such that $Fy = f^{\tau(y)}y \in Y$ for almost all $y \in Y$. Moreover, there is an at most countable measurable partition $\{Y_j\}$ such that τ is constant on partition elements, and constants $\lambda > 1, C > 0$, such that for all $j \geq 1$,

- (1) F restricts to a (measure-theoretic) bijection from Y_j onto Y .
- (2) $d(Fy, Fy') \geq \lambda d(y, y')$ for all $y, y' \in Y_j$.
- (3) $\zeta_0 = \frac{d\mu_0}{d\mu_0 \circ F}$ satisfies $|\log \zeta_0(y) - \log \zeta_0(y')| \leq C d(Fy, Fy')^\eta$ for all $y, y' \in Y_j$.
- (4) $d(f^\ell y, f^\ell y') \leq C d(Fy, Fy')$ for all $0 \leq \ell \leq \tau(y), y, y' \in Y_j$.

A standard consequence of conditions (1)–(3) is that there is a unique ergodic F -invariant absolutely continuous probability measure μ on Y . Moreover, $d\mu/d\mu_0 \in L^\infty(Y)$ and $F : Y \rightarrow Y$ is a Gibbs-Markov map.

Define the induced roof function $\varphi : Y \rightarrow \mathbb{R}^+$ by $\varphi(y) = \sum_{\ell=0}^{\tau(y)-1} h(f^\ell y)$. Since $\varphi \leq |h|_{\infty} \tau$, it follows that $\mu(\varphi > t) = O(t^{-\beta})$ and in particular φ is integrable. We can now define the suspension semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$ and ergodic F_t -invariant and T_t -invariant probability measures μ^φ and $\mu_M = \pi_* \mu^\varphi$ as in Section 2.

Proposition 3.10 *Let $T_t : M \rightarrow M$ be a semiflow satisfying conditions (1)–(4) and (3.3) with h Hölder and $\inf h > 0$. Set $\theta = \lambda^{-\eta^2}$. Then*

- (a) $F_t : Y^\varphi \rightarrow Y^\varphi$ is a Gibbs-Markov semiflow.
- (b) Observables $v \in C^\eta(M) \cap C^{0, \eta}(M)$ lift to observables $\tilde{v} = v \circ \pi : Y^\varphi \rightarrow \mathbb{R}$ that lie in $\mathcal{F}_{\theta, \eta}(Y^\varphi)$. Moreover, there is a constant $C > 0$ such that $\|\tilde{v}\|_{\theta, \eta} \leq C(\|v\|_{C^\eta} + \|v\|_{C^{0, \eta}})$.

Proof Set $\theta_1 = \lambda^{-\eta}$. Let $y, y' \in Y$ with separation time $n = s(y, y')$. By (2),

$$1 \geq \text{diam } Y \geq d(F^n y, F^n y') \geq \lambda^n d(y, y'),$$

so $d(y, y')^\eta \leq (\lambda^{-n})^\eta = \theta_1^n = d_{\theta_1}(y, y')$.

Define $h_\ell = \sum_{j=0}^{\ell-1} h \circ f^j$. By (4), for $y, y' \in Y_j, \ell = 0, \dots, \tau(y) - 1$,

$$\begin{aligned} |h_\ell(y) - h_\ell(y')| &\leq \sum_{j=0}^{\tau(y)-1} |h(f^j y) - h(f^j y')| \leq |h|_\eta \sum_{j=0}^{\tau(y)-1} d(f^j y, f^j y')^\eta \ll \tau(y) d(Fy, Fy')^\eta \\ &\leq \tau(y) d_{\theta_1}(Fy, Fy') \leq \theta_1^{-1} (\inf h)^{-1} \inf_{Y_j} \varphi d_{\theta_1}(y, y') \ll \inf_{Y_j} \varphi d_{\theta_1}(y, y'). \end{aligned} \quad (3.4)$$

Taking $\ell = \tau(y)$ in (3.4), we obtain $|\varphi(y) - \varphi(y')| \ll \inf_{Y_j} \varphi d_\theta(y, y')$ proving part (a).

Next, let $(y, u), (y', u) \in Y^\varphi$ with $s(y, y') \geq 1$. There exists $\ell, \ell' \in \{0, \dots, \tau(y) - 2\}$ such that

$$u \in [h_\ell(y), h_{\ell+1}(y)] \cap [h_{\ell'}(y'), h_{\ell'+1}(y')].$$

Suppose without loss that $\ell \leq \ell'$. Then

$$u = h_\ell(y) + r = h_\ell(y') + r',$$

where $r \in [0, |h|_\infty]$ and $r' = u - h_\ell(y') \geq u - h_{\ell'}(y') \geq 0$. Note that $T_u y = T_r T_{h_\ell(y)} y = T_r f^\ell y$ and similarly $T_u y' = T_{r'} f^{\ell'} y'$. Hence $\tilde{v}(y, u) - \tilde{v}(y', u) = v(T_r f^\ell y) - v(T_{r'} f^{\ell'} y')$. By conditions (4) and (3.3),

$$\begin{aligned} |v(T_r f^\ell y) - v(T_{r'} f^{\ell'} y')| &\leq |v|_{C^\eta} d(T_r f^\ell y, T_{r'} f^{\ell'} y')^\eta \ll |v|_{C^\eta} d(f^\ell y, f^{\ell'} y')^{\eta^2} \\ &\ll |v|_{C^\eta} d(Fy, Fy')^{\eta^2} \leq \theta^{-1} |v|_{C^\eta} d_\theta(y, y'), \end{aligned}$$

and by (3.4),

$$|v(T_r f^\ell y) - v(T_{r'} f^{\ell'} y')| \leq |v|_{C^{0,\eta}} |r - r'|^\eta = |v|_{C^{0,\eta}} |h_\ell(y) - h_\ell(y')|^\eta \ll |v|_{C^{0,\eta}} \varphi(y) d_\theta(y, y').$$

Hence $|\tilde{v}(y, u) - \tilde{v}(y', u)| \ll (|v|_{C^\eta} + |v|_{C^{0,\eta}}) \varphi(y) d_\theta(y, y')$, whenever $s(y, y') \geq 1$. For $s(y, y') = 0$, we have the estimate $|\tilde{v}(y, u) - \tilde{v}(y', u)| \leq 2|v|_\infty = 2|v|_\infty d_\theta(y, y') \ll |v|_\infty \varphi(y) d_\theta(y, y')$, so in all cases we obtain $|\tilde{v}(y, u) - \tilde{v}(y', u)| \ll (|v|_{C^\eta} + |v|_{C^{0,\eta}}) \varphi(y) d_\theta(y, y')$. Also,

$$|\tilde{v}(y, u) - \tilde{v}(y, u')| = |v(T_u y) - v(T_{u'} y)| \leq |v|_{C^{0,\eta}} |u - u'|^\eta.$$

Hence $\tilde{v} \in \mathcal{F}_{\theta,\eta}(Y^\varphi)$ completing the proof of part (b). \blacksquare

It follows that the main results in Subsection 3.1 go over to semiflows $T_t : M \rightarrow M$ satisfying (1)–(4):

Theorem 3.11 *Let $T_t : M \rightarrow M$ be a semiflow satisfying conditions (1)–(4) and (3.3) with h Hölder and $\inf h > 0$. Suppose that $\mu_0(\tau > t) = O(t^{-\beta})$ for some $\beta > 1$. Assume absence of approximate eigenfunctions for the corresponding Gibbs-Markov semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$.*

Then there exists $m \geq 1$ and $C > 0$ such that

$$|\rho_{v,w}(t)| \leq C(\|v\|_{C^\eta} + |v|_{C^{0,\eta}}) |w|_{\infty,m} t^{-(\beta-1)},$$

for all $v \in C^\eta(M) \cap C^{\eta,0}(M)$, $w \in L^{\infty,m}(M)$, $t > 1$.

Proof By Proposition 3.10(a), F_t is a Gibbs-Markov semiflow with $\mu(\varphi > t) = O(t^{-\beta})$. By Proposition 3.10(b), $\tilde{v} = v \circ \pi \in \mathcal{F}_{\theta,\eta}(Y^\varphi)$. Also, $\tilde{w} = w \circ \pi \in L^{\infty,m}(Y^\varphi)$. Hence by (2.1) and Theorem 3.6, $|\rho_{v,w}(t)| = |\rho_{\tilde{v},\tilde{w}}(t)| \ll \|\tilde{v}\|_{\theta,\eta} |\tilde{w}|_{\infty,m} t^{-(\beta-1)} \ll (\|v\|_{C^\eta} + |v|_{C^{0,\eta}}) |w|_{\infty,m} t^{-(\beta-1)}$ for m sufficiently large. \blacksquare

If h is the first return to X and τ is the first return to Y , then observables supported on $\bigcup_{t \in [0, \inf h]} T_t Y$ lift to observables supported on $Y \times [0, \inf \varphi]$ and we can apply Theorem 3.8 to obtain lower bounds for such observables.

Example 3.12 (Intermittent semiflows) We consider a class of two-dimensional Hölder semiflows $T_t : M \rightarrow M$, with an intermittent Poincaré map of Pomeau-Manneville type [76]. Let $X \subset M$ be the interval $[0, 1]$ and suppose as above that $h : X \rightarrow \mathbb{R}^+$ is Hölder with $\inf h > 0$ such that $f(x) = T_{h(x)}x$ defines a map $f : X \rightarrow X$. We continue to assume condition (3.3).

Suppose that $f : X \rightarrow X$ is given by $f(x) = \begin{cases} x(1 + 2^\gamma x^\gamma) & x \in [0, \frac{1}{2}) \\ 2x - 1 & x \in [\frac{1}{2}, 1] \end{cases}$ where $\gamma \in (0, 1)$

is fixed. The maps f were studied by [56, 60, 86] and decay of correlations for Hölder observables is $O(n^{-(\beta-1)})$ where $\beta = 1/\gamma$. This rate is sharp [79, 53].

Here we obtain the analogous results for semiflows. Let $Y = [\frac{1}{2}, 1]$. The first return map $F : Y \rightarrow Y$ satisfies conditions (1)–(4) and the first return time $\tau : Y \rightarrow \mathbb{Z}^+$ satisfies $\mu(\tau > n) \sim c_0 n^{-\beta}$ for some $c_0 > 0$. Hence we obtain decay of correlations $O(\|v\|_{C^\eta} \|w\|_{\infty, m} t^{-(\beta-1)})$ for the semiflow by Theorem 3.11. If in addition h is the first return to X , then a standard calculation (see for example [52, Theorem 1.3] or [66, Proposition 2.17]) shows that $\varphi(y) \sim h(0)\tau(y)$ as $y \rightarrow \frac{1}{2}^+$, $\mu(\varphi > t) \sim c_1 t^{-\beta}$ where $c_1 = h(0)^\beta c_0$, and $\int_t^\infty \mu(\varphi > t') dt' \sim ct^{-(\beta-1)}$ where $c = c_1/(\beta - 1)$. Hence for Hölder observables supported on $\bigcup_{t \in [0, \inf h]} T_t Y$ we obtain the asymptotic $\rho_{v,w}(t) \sim c|\varphi|_1^{-1} \int_M v d\mu_M \int_M w d\mu_M t^{-(\beta-1)}$.

The Markovian nature of the map f simplified the analysis above but is not necessary. In the nonMarkov case, the first return map F is certainly not Gibbs-Markov, but there still exists a induced map $F : Y \rightarrow Y$ (not the first return map) on a suitable set Y such that F is Gibbs-Markov and the return time τ has the same tails as the first return time [86, Section 7]. Hence we still obtain the upper bound $O(t^{-(\beta-1)})$ for mixing rates of Hölder observables by Theorem 3.11. For sharpness of this bound we refer to forthcoming work of [28] which extends [66] to a general functional analytic framework including nonMarkov intermittent (semi)flows.

3.3 Dynamically Hölder semiflows and observables

It is standard that the assumptions on T_t , h and v in Theorem 3.11 can be relaxed from Hölder to dynamically Hölder, as we now describe.

We continue to assume that $v \in C^{0,\eta}(M)$ and that $\inf h > 0$. Condition (3.3) on the semiflow is removed, as are the assumptions that h is Hölder and $v \in C^\eta(M)$. Instead, we require that there are constants $C > 0$, $\gamma \in (0, 1)$ such that for all $y, y' \in Y_j$, $j \geq 1$,

$$\begin{aligned} |h(f^\ell y) - h(f^\ell y')| &\leq C\gamma^{s(y,y')} \quad \text{for } 0 \leq \ell \leq \tau(y) - 1, \\ |v(T_u y) - v(T_u y')| &\leq C\varphi(y)\gamma^{s(y,y')} \quad \text{for } u \in [0, \varphi(y)] \cap [0, \varphi(y')]. \end{aligned}$$

Here, $s(y, y')$ is the separation time for the Gibbs-Markov map F .

It is easily verified that the proof of Proposition 3.10, and hence Theorem 3.11, goes through under these more relaxed assumptions on T_t , h and v .

Remark 3.13 The notion of dynamically Hölder can be relaxed even further, see [14, Section 7.3]. This turns out to be crucial for Lorentz gas examples [14, Remark 9.2].

4 Mixing rates for nonuniformly hyperbolic flows

In this section, we review results on rapid mixing and polynomial mixing for nonuniformly hyperbolic flows. In Subsection 4.1, we define a class of Gibbs-Markov flows with skew product structure (the roof function φ is constant along stable leaves) and state our result on rates of mixing for such flows. In Subsection 4.2, we discuss several situations where the skew product assumption can be relaxed; these include all the examples in the introduction.

4.1 Skew product Gibbs-Markov flows

Let (Y, d) be a metric space with $\text{diam } Y \leq 1$, and let $F : Y \rightarrow Y$ be a piecewise continuous map with ergodic F -invariant probability measure μ . Let \mathcal{W}^s be a cover of Y by disjoint measurable subsets called *stable leaves*. For each $y \in Y$, let $W^s(y)$ denote the stable leaf containing y . We require that $F(W^s(y)) \subset W^s(Fy)$ for all $y \in Y$.

Let \bar{Y} denote the space obtained from Y after quotienting by \mathcal{W}^s , with natural projection $\bar{\pi} : Y \rightarrow \bar{Y}$. We assume that the quotient map $\bar{F} : \bar{Y} \rightarrow \bar{Y}$ is a Gibbs-Markov map as in Definition 3.1, with partition $\{\bar{Y}_j\}$, separation time $s(y, y')$, and ergodic invariant probability measure $\bar{\mu} = \bar{\pi}_*\mu$.

Let $Y_j = \bar{\pi}^{-1}\bar{Y}_j$; these form a partition of Y and each Y_j is a union of stable leaves. The separation time extends to Y , setting $s(y, y') = s(\bar{\pi}y, \bar{\pi}y')$ for $y, y' \in Y$.

Next, we require that there is a measurable subset $\tilde{Y} \subset Y$ such that for every $y \in Y$ there is a unique $\tilde{y} \in \tilde{Y} \cap W^s(y)$. Let $\pi : Y \rightarrow \tilde{Y}$ define the associated projection $\pi y = \tilde{y}$. (Note that \tilde{Y} can be identified with \bar{Y} , but in general $\pi_*\mu \neq \bar{\mu}$.)

We assume that there are constants $C > 0$, $\gamma \in (0, 1)$ such that

- (a) $d(F^n y, F^n y') \leq C\gamma^n$ for all $n \geq 0$ and all $y, y' \in Y$ with $y' \in W^s(y)$.
- (b) $d(F^n y, F^n y') \leq C\gamma^{s(y, y') - n}$ for all $n \geq 0$ and all $y, y' \in \tilde{Y}$.

Let $\varphi : Y \rightarrow \mathbb{R}^+$ be an integrable roof function with $\inf \varphi > 0$, and define the suspension flow $F_t : Y^\varphi \rightarrow Y^\varphi$ with ergodic invariant probability measure μ^φ (see Section 2).¹

In this subsection, we suppose that φ is constant along stable leaves and hence projects to a well-defined roof function $\varphi : \bar{Y} \rightarrow \mathbb{R}^+$. It follows that the suspension flow F_t projects to a suspension semiflow $\bar{F}_t : \bar{Y}^\varphi \rightarrow \bar{Y}^\varphi$. We assume that $\varphi : \bar{Y} \rightarrow \mathbb{R}^+$ satisfies condition (3.1), so \bar{F}_t is a Gibbs-Markov semiflow. We call F_t a *skew product Gibbs-Markov flow*, and we say that F_t has *approximate eigenfunctions* if \bar{F}_t has approximate eigenfunctions (Definition 3.4).

Fix $\eta \in (0, 1]$ and let $\mathcal{H}_{\gamma, \eta}(Y^\varphi)$ be the space of observables $v : Y^\varphi \rightarrow \mathbb{R}$ with $\|v\|_{\gamma, \eta} < \infty$, where

$$|v|_{\gamma, \eta} = \sup_{\substack{(y, u), (y', u') \in Y^\varphi \\ (y, u) \neq (y', u')}} \frac{|v(y, u) - v(y', u')|}{\varphi(y)\{d(y, y') + \gamma^{s(y, y')}\} + |u - u'|^\eta}, \quad \|v\|_{\gamma, \eta} = |v|_\infty + |v|_{\gamma, \eta}.$$

We also define $\mathcal{H}_{\gamma, 0, m}(Y^\varphi)$ to consist of observables that lie in $\mathcal{H}_\gamma(Y^\varphi)$ and are m -times differentiable in the flow direction with derivatives in $\mathcal{H}_\gamma(Y^\varphi)$, with norm $\|w\|_{\gamma, 0, m} = \sum_{j=0}^m \|\partial_t^j w\|_\gamma$.

¹Strictly speaking, F_t is not always a flow since F need not be invertible. However, F_t is used as a model for various flows, and it is then a flow when φ is the first return to Y , so it is convenient to call it a flow.

In [14], we prove:

Theorem 4.1 *Suppose that $F_t : Y^\varphi \rightarrow Y^\varphi$ is a skew product Gibbs-Markov flow such that $\mu(\varphi > t) = O(t^{-\beta})$ for some $\beta > 1$. Assume absence of approximate eigenfunctions. Then there exists $m \geq 1$ and $C > 0$ such that*

$$|\rho_{v,w}(t)| \leq C \|v\|_{\gamma,\eta} \|w\|_{\gamma,0,m} t^{-(\beta-1)} \quad \text{for all } v \in \mathcal{H}_{\gamma,\eta}(Y^\varphi), w \in \mathcal{H}_{\gamma,0,m}(Y^\varphi), t > 1.$$

4.2 General nonuniformly hyperbolic flows

In this subsection, we drop the assumption that φ is constant along stable leaves, and mention various classes of nonuniformly hyperbolic flows that do not possess a skew product structure to which our methods apply. For details we refer to [14] who introduce a class of *Gibbs-Markov flows* that are conjugate by a Hölder conjugacy to a skew product Gibbs-Markov flow. As shown in [14], our results on rapid and polynomial mixing go over to Gibbs-Markov flows and to nonuniformly hyperbolic flows that are modelled by Gibbs-Markov flows. This includes the following situations.

(i) Flows with exponential contraction along stable leaves Condition (a) in Subsection 4.1 asserts exponential contraction along stable leaves for the uniformly hyperbolic map $F : Y \rightarrow Y$. This means that exponential contraction is assumed only on returns to the inducing set Y . Note that such a return occurs at the flow time $\varphi_n = \sum_{j=0}^{n-1} \varphi \circ F^j$, so an alternative stronger condition is that $d(F^n y, F^n y') \leq C \gamma^{\varphi_n(y)}$ for all $n \geq 0$ and all $y, y' \in Y$ with $y' \in W^s(y)$. This condition was assumed in [63, 64] and incorporates all of the rapidly mixing examples in Section 1.1. Indeed, it is automatic for flows modelled by Young towers with exponential tails. However for flows modelled by Young towers with polynomial tails, the condition is very restrictive and excludes the slowly mixing billiard examples.

(ii) Roof functions with bounded Hölder constants Condition (3.1) reflects the fact that the variation of φ on partition elements is likely to be as large as the size of φ . The argument in Section 3.2 indicates that this is the “correct” condition in general. A stronger condition is that $|\varphi(y) - \varphi(y')| \leq C_1 \theta^{s(y,y')}$ along unstable leaves. This includes [14] the slowly mixing Lorentz gas examples in Section 1.3.

(iii) Flows with Hölder stable foliation Under the assumption that T_t has a Hölder stable foliation \mathcal{W}^{ss} (in a neighborhood of the attractor Λ) we can reduce to the skew product case by using a cross-section Y comprising leaves in \mathcal{W}^{ss} . (For flows with a DT_t -invariant dominated splitting $T_\Lambda M = E^{ss} \oplus E^{cu}$, results on the existence of a Hölder stable foliation \mathcal{W}^{ss} can be found in [6, Section 4] and [7, Theorem 6.2].)

Example 4.2 (Intermittent solenoidal flows) The classical Smale-Williams solenoid construction can be adapted (see for example [4, Section 5] and [70, Example 4.2]) to construct intermittent transformations $f : X \rightarrow X$ that are the invertible analogue of the intermittent maps in Example 3.12. These have polynomial decay rates $O(n^{-(\beta-1)})$ for any specified $\beta > 1$. Hence we can construct intermittent flows $T_t : M \rightarrow M$ with $f : X \rightarrow X$ as a Poincaré map and Hölder return time function $h : X \rightarrow \mathbb{R}^+$ with $\inf h > 0$.

The examples in [4] and some of the examples in [70] have exponential contraction along stable leaves and hence fall within scenario (i). However even the examples in [70] with slow contraction along stable leaves are covered by scenario (iii). Hence we obtain polynomial mixing $O(t^{-(\beta-1)})$ for general intermittent solenoidal flows. Again these results are sharp by [66] in the Markovian case and [28] in the general case. Optimal lower bounds, asymptotics and error rates are achieved by observables that are supported away from the neutral periodic orbit and constant along stable leaves.

5 Criteria for absence of approximate eigenfunctions

The approximate eigenfunction condition in Definition 3.4 is somewhat technical. In this subsection, we discuss three sufficient conditions to rule out the existence of approximate eigenfunctions. We suppose throughout that $F_t : Y^\varphi \rightarrow Y^\varphi$ is a Gibbs-Markov semiflow or skew product Gibbs-Markov flow, but it is immediate from the definitions in [14] that the conditions apply to the situations mentioned in Subsection 4.2.

In Subsection 5.1, we show that a Diophantine condition on the periods of three periodic solutions ensures absence of approximate eigenfunctions. This condition is satisfied with probability one but is not robust. In Subsection 5.2, we use good asymptotics of periodic data to give an open and dense condition. In Subsection 5.3, we define the temporal distortion function and give a condition involving the dimension of its range.

In preparation for Subsections 5.1 and 5.2, we recall the relationship between periodic data and approximate eigenfunctions. Define $\varphi_n = \sum_{j=0}^{n-1} \varphi \circ F^j$. If y is a periodic point of period p for F (that is, $F^p y = y$), then y is periodic of period $\mathcal{L} = \varphi_p(y)$ for F_t (that is, $F_{\mathcal{L}} y = y$).

Proposition 5.1 *Suppose that there exist approximate eigenfunctions on $Z_0 \subset Y$. Let α, C, b_k, n_k be as in Definition 3.4. If $y \in Z_0$ is a periodic point with $F^p y = y$ and $F_{\mathcal{L}} y = y$ where $\mathcal{L} = \varphi_p(y)$, then*

$$\text{dist}(b_k n_k \mathcal{L} - p\psi_k, 2\pi\mathbb{Z}) \leq C(\inf \varphi)^{-1} \mathcal{L} |b_k|^{-\alpha} \quad \text{for all } k \geq 1. \quad (5.1)$$

Proof For $n, p \geq 1$,

$$M_b^{np} u - e^{ip\psi} u = M_b^n (M_b^{n(p-1)} u - e^{i(p-1)\psi} u) + e^{i(p-1)\psi} (M_b^n u - e^{i\psi} u),$$

so $|M_b^{np} u - e^{ip\psi} u| \leq |M_b^{n(p-1)} u - e^{i(p-1)\psi} u| + |M_b^n u - e^{i\psi} u|$. Inductively, $|M_b^{np} u - e^{ip\psi} u| \leq p|M_b^n u - e^{i\psi} u|$ and hence by (3.2),

$$|(M_b^{pn_k} u_k)(y) - e^{ip\psi_k} u_k(y)| \leq Cp |b_k|^{-\alpha} \quad \text{for all } y \in Z_0, k \geq 1, p \geq 1. \quad (5.2)$$

Next, $(M_b^p v)(y) = e^{ib\varphi_p(y)} v(F^p y) = e^{ib\mathcal{L}} v(y)$. Hence substituting y into (5.2), we obtain $|e^{ib_k n_k \mathcal{L}} - e^{ip\psi_k}| \leq Cp |b_k|^{-\alpha}$. Also $\mathcal{L} = \varphi_p(y) \geq p \inf \varphi$. \blacksquare

Remark 5.2 We now have the notation required to provide the formula for t_q promised in Section 1.2. Recall that q is a periodic point for the flow. Let $y \in Y$ be a point lying on the periodic orbit through q with $F^p y = y$. By [8, Corollary 6.2], we can choose $t_q = \varphi_p(y)$, so t_q coincides with the period of q for the suspension flow F_t . (If φ is the first return time to Y then t_q is also the period of q under T_t .)

5.1 Diophantine condition on periods

Proposition 5.3 *Let $y_1, y_2, y_3 \in \bigcup Y_j$ be fixed points for F , and let $\mathcal{L}_i = \varphi(y_i)$, $i = 1, 2, 3$, be the corresponding periods for F_t . Let Z_0 be the finite subsystem corresponding to the three partition elements containing y_1, y_2, y_3 .*

If $(\mathcal{L}_1 - \mathcal{L}_3)/(\mathcal{L}_2 - \mathcal{L}_3)$ is Diophantine, then there do not exist approximate eigenfunctions on Z_0 .

Proof We give the proof when F_t is a Gibbs-Markov semiflow. It is immediate from the definitions that the result is the same for skew product Gibbs-Markov flows.

Define $\mathcal{L}_{i3} = \mathcal{L}_i - \mathcal{L}_3$, $i = 1, 2$. For $\mathcal{L}_{13}/\mathcal{L}_{23}$ Diophantine, there exists $\alpha' > 2$ such that $|\mathcal{L}_{13}/\mathcal{L}_{23} - m_1/m_2| \leq C|m_2|^{-\alpha'}$ has only finitely many integer solutions m_1, m_2 for each $C > 0$.

Arguing as in [48, Section 13], we choose $\alpha > \alpha'$. Suppose for contradiction that there exist approximate eigenfunctions on Z_0 . Substituting $y = y_i$ in (5.1), we obtain the estimates

$$\text{dist}(b_k n_k \mathcal{L}_i - \psi_k, 2\pi\mathbb{Z}) = O(|b_k|^{-\alpha}), \quad i = 1, 2, 3,$$

where $|b_k| \rightarrow \infty$ and $n_k = O(\ln |b_k|)$ as $k \rightarrow \infty$. Eliminating ψ_k , we obtain

$$\text{dist}(b_k n_k \mathcal{L}_{i3}, 2\pi\mathbb{Z}) = O(|b_k|^{-\alpha}), \quad i = 1, 2.$$

Hence, for each k we have integers m_1, m_2 such that

$$b_k n_k \mathcal{L}_{i3} = 2\pi m_i + O(|b_k|^{-\alpha}),$$

where in particular $m_2 = O(b_k n_k) = O(b_k \ln |b_k|)$. Also, $m_1/m_2 \sim \mathcal{L}_{13}/\mathcal{L}_{23}$ so $m_1 = O(m_2)$. Hence

$$b_k n_k \mathcal{L}_{i3} = 2\pi m_i + O(|m_2|^{-\alpha'}), \quad i = 1, 2.$$

It follows that $\frac{\mathcal{L}_{13}}{\mathcal{L}_{23}} = \frac{m_1}{m_2} + O(|m_2|^{-\alpha'})$, which is the desired contradiction. \blacksquare

5.2 Good asymptotics

We recall the following definition from [50]:

Definition 5.4 Let $y_0 \in Y$ be a fixed point for F with period $\mathcal{L}_0 = \varphi(y_0)$ for the flow. A sequence of periodic points $y_N \in Y$, $N \geq 1$, with $F^N y_N = y_N$ has *good asymptotics* if their periods $\mathcal{L}_N = \varphi_N(y_N)$ for the flow satisfy

$$\mathcal{L}_N = N\mathcal{L}_0 + \kappa + E_N \gamma^N \cos(N\omega + \omega_N) + o(\gamma^N),$$

where $\kappa \in \mathbb{R}$, $\gamma \in (0, 1)$ are constants, $E_N \in \mathbb{R}$ is a bounded sequence with $\liminf_{N \rightarrow \infty} |E_N| > 0$, and either (i) $\omega = 0$ and $\omega_N \equiv 0$, or (ii) $\omega \in (0, \pi)$ and $\omega_N \in (\omega_0 - \pi/12, \omega_0 + \pi/12)$ for some ω_0 .

Proposition 5.5 *If there exists a sequence of periodic points with good asymptotics in a finite subsystem Z_0 , then there do not exist approximate eigenfunctions on Z_0 .*

Proof We argue as in the proof of [50, Theorem 1.6(a)]. The proof works for any fixed $\alpha > 0$.

Note that $\mathcal{L}_N = O(N + 1)$. Suppose for contradiction that there exist approximate eigenfunctions on Z_0 . Substituting y_N into (5.1) we obtain the estimates

$$\text{dist}(b_k n_k \mathcal{L}_N - N \psi_k, 2\pi\mathbb{Z}) = O(N|b_k|^{-\alpha}), \quad \text{dist}(b_k n_k N \mathcal{L}_0 - N \psi_k, 2\pi\mathbb{Z}) = O(N|b_k|^{-\alpha}),$$

for all $k \geq 1, N \geq 1$. Hence we can eliminate ψ_k and \mathcal{L}_0 simultaneously, yielding

$$\text{dist}(b_k n_k (\kappa + E_N \gamma^N \cos(N\omega + \omega_N) + o(\gamma^N)), 2\pi\mathbb{Z}) = O(N|b_k|^{-\alpha}), \quad (5.3)$$

for all $k \geq 1, N \geq 1$.

Momentarily, choose $N = N(k) = [\rho \ln |b_k|]$ in (5.3). For $\rho > 0$ large enough but fixed, we have $|b_k| n_k \gamma^{N(k)} = O(|b_k|^{-\alpha})$. It follows that $\text{dist}(b_k n_k \kappa, 2\pi\mathbb{Z}) = O(|b_k|^{-\alpha} \ln |b_k|)$ for all $k \geq 1$. Combining this with (5.3) (which still holds for all $N \geq 1$), we obtain for any $\alpha' < \alpha$ that

$$\text{dist}(b_k n_k (E_N \gamma^N \cos(N\omega + \omega_N) + o(\gamma^N)), 2\pi\mathbb{Z}) = O(N|b_k|^{-\alpha'}), \quad (5.4)$$

for all $k \geq 1, N \geq 1$.

Let $S = \sup_N |E_N|$ and set $M(k) = [\ln(|b_k| n_k S) / (-\ln \gamma)] + 1$. Then $|b_k| n_k S \gamma^{M(k)} \in [\gamma, 1)$. Taking $N = M(k) + j$ in (5.4), we obtain

$$\lim_{k \rightarrow \infty} b_k n_k E_{M(k)+j} \gamma^{M(k)} \cos((M(k) + j)\omega + \omega_{M(k)+j}) = 0 \quad \text{for all } j \in \mathbb{Z}.$$

Since $|b_k n_k \gamma^{M(k)}| \geq \gamma/S$ and $\liminf_{N \rightarrow \infty} |E_N| > 0$, it follows that

$$\lim_{k \rightarrow 0} \{(M(k) + j)\omega + \omega_{M(k)+j}\} = \frac{\pi}{2} \bmod \pi \quad \text{for all } j \in \mathbb{Z}. \quad (5.5)$$

This is clearly impossible when $\omega = \omega_N = 0$ for all N , so it remains to consider the case $\omega \in (0, \pi)$ and $|\omega_N - \omega_0| < \pi/12$. Taking differences of (5.5) for different values of j , we obtain that $\ell\omega \in [-\pi/6, \pi/6] \bmod \pi$ for all ℓ , which is impossible, providing the desired contradiction. \blacksquare

As shown in [50], for any finite subsystem Z_0 , the existence of periodic points with good asymptotics in Z_0 is a C^2 -open and C^∞ -dense condition. (Although [50] is set in the uniformly hyperbolic setting, the construction of periodic points with good asymptotics uses only the existence of a transverse homoclinic point y_0 .)

5.3 Temporal distance function

In this subsection, following [64, Section 5.3] and [8, Section 3], we extend an argument of Dolgopyat [48, Appendix] for Axiom A flows to the nonuniformly hyperbolic setting. We assume throughout that we are in the setup of Section 4.1. In particular, φ is constant along stable leaves in the cover \mathcal{W}^s .

First, we recall the notion of product structure on Y , taking only the parts from [85] that are needed here. Assume that there is a second cover \mathcal{W}^u of Y by disjoint measurable subsets (called *unstable leaves*) such that each stable leaf intersects each unstable leaf in

precisely one point. For each $y \in Y$ let $W^u(y)$ denote the unstable leaf containing y . We require that $F(W^u(y) \cap Y_j) \supset W^u(Fy)$ for all $y \in Y_j$, $j \geq 1$.

Also we assume that there are constants $C > 0$, $\gamma \in (0, 1)$ such that

$$|\varphi(y) - \varphi(y')| \leq C\gamma^{s(y,y')} \quad \text{for all } y, y' \in Y \text{ with } s(y, y') \geq 1 \text{ and } y' \in W^u(y).$$

Given $y, y' \in Y$, $y' \in W^u(y)$, we define compatible inverse branches $F^n y$, $F^n y'$ for $n \leq -1$ as follows. First, set $z_0 = y$, $z'_0 = y'$. By transitivity of F , there exists $z_1 \in Y$ such that $Fz_1 = z_0$. Let Y_{j_1} be the partition element containing z_1 . Since $F(W^u(z_1) \cap Y_{j_1}) \supset W^u(z_0)$, there exists $z'_1 \in W^u(z_1) \cap Y_{j_1}$ such that $Fz'_1 = z'_0$. Inductively, we obtain a sequence of partition elements Y_{j_n} and pairs of points $z_n, z'_n \in Y_{j_n}$ with $z'_n \in W^u(z_n)$ for all $n \geq 1$. Note that $|\varphi(z_n) - \varphi(z'_n)| \leq C\gamma^{s(z_n, z'_n)} \leq C\gamma^n$ for all $n \geq 1$, so we obtain a well-defined function

$$D_0(y, y') = \sum_{n=1}^{\infty} (\varphi(z_n) - \varphi(z'_n)) = \sum_{n=-\infty}^{-1} (\varphi(F^n y) - \varphi(F^n y')).$$

Now, let $y_1, y_4 \in Y$ and set $y_2 = W^s(y_1) \cap W^u(y_4)$, $y_3 = W^u(y_1) \cap W^s(y_4)$. For $n \leq -1$ choose compatible inverse branches $F^n y_1$ and $F^n y_3$ as done in the definition of D_0 , and similarly for $F^n y_4$ and $F^n y_2$. Define the *temporal distance function* $D : Y \times Y \rightarrow \mathbb{R}$,

$$\begin{aligned} D(y_1, y_4) &= \sum_{n=-\infty}^{\infty} \left(\varphi(F^n y_1) - \varphi(F^n y_2) - \varphi(F^n y_3) + \varphi(F^n y_4) \right) \\ &= \sum_{n=-\infty}^{-1} \left(\varphi(F^n y_1) - \varphi(F^n y_2) - \varphi(F^n y_3) + \varphi(F^n y_4) \right), \end{aligned}$$

where the last equality follows from the assumption that φ is constant along stable leaves. This coincides with $D_0(y_1, y_3) - D_0(y_4, y_2)$ and hence is well-defined.

Theorem 5.6 *Let $Z_0 = \bigcap_{n=0}^{\infty} F^{-n}Z$ where Z is a union of finitely many elements of the partition $\{Y_j\}$. Let \bar{Z}_0 denote the corresponding finite subsystem of \bar{Y} . If the lower box dimension of $D(Z_0 \times Z_0)$ is positive, then there do not exist approximate eigenfunctions on \bar{Z}_0 .*

Proof We suppose that there exist approximate eigenfunctions on \bar{Z}_0 and show that $\text{BD}(D(Z_0 \times Z_0)) = 0$.

First, note that $D(y_1, y_4) = D_m(y_1, y_4) + O(\gamma^m)$ where

$$\begin{aligned} D_m(y_1, y_4) &= \sum_{n=-m}^{-1} \left(\varphi(F^n y_1) - \varphi(F^n y_2) - \varphi(F^n y_3) + \varphi(F^n y_4) \right) \\ &= \varphi_m(F^{-m} y_1) - \varphi_m(F^{-m} y_2) - \varphi_m(F^{-m} y_3) + \varphi_m(F^{-m} y_4). \end{aligned}$$

Hence

$$\exp\{ibD(y_1, y_4)\} = \frac{\exp\{ib\varphi_m(F^{-m} y_1)\} \exp\{ib\varphi_m(F^{-m} y_4)\}}{\exp\{ib\varphi_m(F^{-m} y_2)\} \exp\{ib\varphi_m(F^{-m} y_3)\}} + O(|b|\gamma^m). \quad (5.6)$$

By Definition 3.4, there are sequences such that

$$|\exp\{ib_k\bar{\varphi}_{n_k}\}u_k \circ \bar{F}^{n_k} - e^{i\psi_k}u_k| \ll |b_k|^{-\alpha} \quad \text{on } \bar{Z}_0.$$

Recall from Definition 3.4 that $|u_k| \equiv 1$, so

$$\exp\{ib_k\varphi_{n_k}\} = \exp\{ib_k\bar{\varphi}_{n_k}\} \circ \bar{\pi} = \frac{e^{i\psi_k}u_k \circ \bar{\pi}}{u_k \circ \bar{\pi} \circ F^{n_k}} + O(|b_k|^{-\alpha}) \quad \text{on } Z_0.$$

Substituting into (5.6) and using that $\bar{\pi}y_1 = \bar{\pi}y_2$, $\bar{\pi}y_3 = \bar{\pi}y_4$, we obtain

$$\exp\{ib_k D(y_1, y_4)\} = \frac{u_k \circ \bar{\pi}(F^{-n_k}y_1)u_k \circ \bar{\pi}(F^{-n_k}y_4)}{u_k \circ \bar{\pi}(F^{-n_k}y_2)u_k \circ \bar{\pi}(F^{-n_k}y_3)} + O(|b_k|^{-\alpha}) + O(|b_k|\gamma^{n_k}).$$

Recall from Definition 3.4 that $|u_k|_\theta \ll |b_k|$. Also, α can be fixed arbitrarily large and $n_k = \lceil \xi \ln |b_k| \rceil$ where ξ can be fixed arbitrarily large. We choose ξ so that $\xi \ln \theta + 1 \leq -\alpha$ and $\xi \ln \gamma + 1 \leq -\alpha$. Then

$$|b_k|\theta^{n_k} \leq \theta^{-1}|b_k|\theta^{\xi \ln |b_k|} = \theta^{-1}|b_k|^{\xi \ln \theta + 1} \leq \theta^{-1}|b_k|^{-\alpha},$$

and similarly $|b_k|\gamma^{n_k} \leq \theta^{-1}|b_k|^{-\alpha}$. Hence

$$\begin{aligned} \left| \frac{u_k \circ \bar{\pi}(F^{-n_k}y_1)}{u_k \circ \bar{\pi}(F^{-n_k}y_3)} - 1 \right| &= |u_k \circ \bar{\pi}(F^{-n_k}y_1) - u_k \circ \bar{\pi}(F^{-n_k}y_3)| \\ &\leq |u_k|_\theta d_\theta(\bar{\pi}(F^{-n_k}y_1), \bar{\pi}(F^{-n_k}y_3)) \ll |b_k|\theta^{s(\bar{\pi}(F^{-n_k}y_1), \bar{\pi}(F^{-n_k}y_3))} \leq |b_k|\theta^{n_k} \leq \theta^{-1}|b_k|^{-\alpha}. \end{aligned}$$

The same estimate holds with y_1 and y_3 replaced by y_2 and y_4 , so

$$|\exp\{ib_k D(y_1, y_4)\} - 1| = O(|b_k|^{-\alpha}) \quad \text{for all } y_1, y_4 \in Z_0, k \geq 1.$$

This means that there is a constant $C > 0$ such that

$$D(Z_0 \times Z_0) \subset \bigcup_{j \in \mathbb{Z}} \left(\frac{2\pi j}{b_k} - \frac{C}{|b_k|^{\alpha+1}}, \frac{2\pi j}{b_k} + \frac{C}{|b_k|^{\alpha+1}} \right) \quad \text{for all } k \geq 1.$$

Hence $\underline{\text{BD}}(D(Z_0 \times Z_0)) \leq 1/(\alpha + 1)$. The result follows since α is arbitrarily large. \blacksquare

Remark 5.7 For Axiom A attractors, Z_0 can be taken to be connected and D is continuous, so absence of approximate eigenfunctions is ensured whenever D is not identically zero [47, Section 9]. For nonuniformly hyperbolic flows, where the partition $\{Y_j\}$ is countably infinite, Z_0 is a Cantor set of positive Hausdorff dimension [64, Example 5.7]. In general it is not clear how to use this property since D is generally at best Hölder. However for flows with a contact structure, a formula for D in [57, Lemma 3.2] can be exploited, see [64, Example 5.7]. Hence when there is a contact structure (which includes the Lorentz gas flows in the introduction when there is no external forcing) absence of approximate eigenfunctions is automatic.

Part II

Proof of theorems for semiflows

In this part of the review article, we give a self-contained proof of the main results on rapid and polynomial mixing for semiflows. In Section 6, we recall how to use the Laplace transform of the correlation function to deduce rates of mixing. Theorems 3.7 (rapid mixing) and Theorem 3.6 (polynomial mixing) are proved in Sections 7 and 8 respectively.

We note that the proof of Theorem 3.7 plays a crucial role in the proof of Theorem 3.6, justifying the movement of certain contours of integration to the imaginary axis in Section 8.4. Theorem 3.7 also illustrates many of the main ideas while avoiding numerous technical issues needed for Theorem 3.6.

6 Strategy for obtaining rates of mixing

In this section, we recall how to use smoothness of the Laplace transform $\hat{\rho}_{v,w}$ to deduce decay rates for $\rho_{v,w}$. Basic facts about inversion of Laplace transforms and decay rates are recalled in Subsection 6.1. In Subsection 6.2, we prove a version of Pollicott's formula [74] for $\hat{\rho}_{v,w}$.

6.1 Laplace transforms and inversion formulas

Define $\mathbb{H} = \{s \in \mathbb{C} : \operatorname{Re} s > 0\}$ and $\bar{\mathbb{H}} = \{s \in \mathbb{C} : \operatorname{Re} s \geq 0\}$.

The Laplace transform $\hat{\rho}_{v,w}(s) = \int_0^\infty e^{-st} \rho_{v,w}(t) dt$ is analytic on \mathbb{H} . We deduce decay of $\rho_{v,w}$ from smoothness of $\hat{\rho}_{v,w}$. It is convenient to make a C^∞ modification so that $\rho_{v,w}$ is unchanged for $t \geq 1$ but vanishes near zero. (Such a modification does not affect the asymptotics of $\rho_{v,w}$ nor the smoothness of $\hat{\rho}_{v,w}$.) Abusing notation, we still write $\rho_{v,w}$ and $\hat{\rho}_{v,w}$ for these modified functions.

Let \mathcal{A} be a subspace of $L^\infty(Y^\varphi)$ with norm $\| \cdot \|_{\mathcal{A}} \geq | \cdot |_\infty$. For $m \geq 0$, define $\mathcal{A}_m = \{w \in \mathcal{A} : \|w\|_{\mathcal{A},m} < \infty\}$ to be the subspace of observables that are m -times differentiable in the flow direction, where $\|w\|_{\mathcal{A},m} = \sum_{j=0}^m \|\partial_t^j w\|_{\mathcal{A}}$. Note that $\mathcal{A}_m \subset L^{\infty,m}(Y^\varphi) = (L^\infty(Y^\varphi))_m$.

If $v \in L^1(Y^\varphi)$ and $w \in L^{\infty,m}(Y^\varphi)$, then $\rho_{v,w}$ is m -times differentiable and $\rho_{v,w}^{(m)} = \rho_{v,\partial_t^m w}$. Hence, performing integration by parts m times, we obtain

$$\hat{\rho}_{v,w}(s) = s^{-m} \hat{\rho}_{v,\partial_t^m w}(s) \quad \text{for } s \in \mathbb{H}. \quad (6.1)$$

Corollary 6.1 *Let $v \in L^1(Y^\varphi)$, $w \in L^{\infty,2}(Y^\varphi)$. Then*

$$\rho_{v,w}(t) = \int_{-\infty}^\infty e^{(\epsilon+ib)t} \hat{\rho}_{v,w}(\epsilon+ib) db \quad \text{for any } \epsilon > 0.$$

Proof Note that $|\hat{\rho}_{v,w}(a+ib)| \leq a^{-1} |\rho_{v,w}|_\infty \leq 2a^{-1} |v|_1 |w|_\infty$ for all $s = a+ib \in \mathbb{H}$. Also, by (6.1), $|\hat{\rho}_{v,w}(a+ib)| = |s|^{-2} |\hat{\rho}_{v,\partial_t^2 w}(a+ib)| \leq 2a^{-1} |s|^{-2} |v|_1 |\partial_t^2 w|_\infty$. Combining these,

$$|\hat{\rho}_{v,w}(s)| \leq 4a^{-1} (|s|^2 + 1)^{-1} |v|_1 |w|_{\infty,2} \quad \text{for all } s = a+ib \in \mathbb{H}.$$

Hence, we can apply the classical inverse Laplace transform formula. ■

Lemma 6.2 *Let $v \in L^1(Y^\varphi)$, $\epsilon > 0$, $q \geq 1$. Suppose that*

(i) $s \mapsto \hat{\rho}_{v,w}(s)$ *is continuous on $\{\operatorname{Re} s \in [0, \epsilon]\}$ and $b \mapsto \hat{\rho}_{v,w}(ib)$ is C^q on \mathbb{R} for all $w \in \mathcal{A}$.*

(ii) *There exist constants $C, \alpha > 0$ such that*

$$|\hat{\rho}_{v,w}(s)| \leq C(|b| + 1)^\alpha \|w\|_\infty \quad \text{and} \quad |\hat{\rho}_{v,w}^{(j)}(ib)| \leq C(|b| + 1)^\alpha \|w\|_{\mathcal{A}},$$

for all $w \in \mathcal{A}$, $j \leq q$, and all $s = a + ib \in \mathbb{C}$ with $a \in [0, \epsilon]$.

Let $m = \lceil \alpha \rceil + 2$. Then there exists a constant $C' > 0$ depending only on C, q, α , such that

$$|\rho_{v,w}(t)| \leq C' \|w\|_{\mathcal{A},m} t^{-q} \quad \text{for all } w \in \mathcal{A}_m, t > 1.$$

Proof By Corollary 6.1, $\rho_{v,w}(t) = \int_{-\infty}^{\infty} e^{(\epsilon+ib)t} \hat{\rho}_{v,w}(\epsilon + ib) db$. By (6.1) and condition (ii), $\hat{\rho}_{v,w}(a + ib) = O(|b|^{-2})$ uniformly in $a \in [0, \epsilon]$ as $b \rightarrow \infty$. Using this together with the continuity property in (i) and a standard extension of the Cauchy integral theorem, we can move the contour of integration to the imaginary axis, so

$$\rho_{v,w}(t) = \int_{-\infty}^{\infty} e^{ibt} \hat{\rho}_{v,w}(ib) db.$$

Again, by (6.1) and conditions (i,ii), $\hat{\rho}_{v,w}^{(j)}(ib)$ is integrable for all $j \leq q$, so the result follows by standard Fourier analysis. (See also Proposition 8.2 below.) \blacksquare

6.2 Formula for $\hat{\rho}_{v,w}$

We use the following variant of Pollicott's formula [74]. Since $\rho_{v,w}(t) = \int_{Y^\varphi} (v - \int_{Y^\varphi} v d\mu) w \circ F_t d\mu^\varphi$, we may suppose throughout that $\int_{Y^\varphi} v d\mu^\varphi = 0$.

Given observables $v, w : Y^\varphi \rightarrow \mathbb{R}$, define

$$\begin{aligned} V(t)(y) &= v(y, \varphi(y) - t) = 1_{\{0 \leq t \leq \varphi(y)\}} v(y, \varphi(y) - t), \\ w(t)(y) &= w(y, t) = 1_{\{0 \leq t \leq \varphi(y)\}} w(y, t). \end{aligned}$$

The corresponding Laplace transforms $\hat{V}(s), \hat{w}(s) : Y \rightarrow \mathbb{C}$, $s \in \mathbb{H}$, are given by

$$\hat{V}(s)(y) = \int_0^{\varphi(y)} e^{-s(\varphi(y)-u)} v(y, u) du, \quad \hat{w}(s)(y) = \int_0^{\varphi(y)} e^{-su} w(y, u) du.$$

Also, define $v_s(y) = \int_0^{\varphi(y)} e^{su} v(y, u) du$.

Let $R : L^1(Y) \rightarrow L^1(Y)$ denote the transfer operator corresponding to $F : Y \rightarrow Y$. So $\int_Y v w \circ F d\mu = \int_Y Rv w d\mu$ for all $v \in L^1(Y)$ and $w \in L^\infty(Y)$.

For $s \in \mathbb{H}$, define the *twisted transfer operators*

$$\hat{R}(s) : L^1(Y) \rightarrow L^1(Y), \quad \hat{R}(s)v = R(e^{-s\varphi}v).$$

Proposition 6.3 *Let $v, w \in L^\infty(Y^\varphi)$ with $\int_{Y^\varphi} v d\mu^\varphi = 0$. Then $\rho_{v,w} = \sum_{n=0}^\infty J_n$ where $|J_0(t)| \leq |v|_\infty |w|_\infty |\varphi|_1^{-1} \int_Y 1_{\{\varphi > t\}} \varphi d\mu$ for all $t > 0$ and*

$$\widehat{J}_n(s) = |\varphi|_1^{-1} \int_Y e^{-s\varphi_n} v_s \widehat{w}(s) \circ F^n d\mu \quad \text{for all } s \in \overline{\mathbb{H}}, n \geq 1.$$

Proof Write

$$\begin{aligned} \rho(t) &= \int_{Y^\varphi} 1_{\{t+u < \varphi(y)\}} v(y, u) w \circ F_t(y, u) d\mu^\varphi \\ &\quad + \sum_{n=1}^\infty \int_{Y^\varphi} 1_{\{\varphi_n(y) < t+u < \varphi_{n+1}(y)\}} v(y, u) w \circ F_t(y, u) d\mu^\varphi = \sum_{n=0}^\infty J_n(t), \end{aligned}$$

where

$$J_n(t) = \begin{cases} \int_{Y^\varphi} 1_{\{t+u < \varphi(y)\}} v(y, u) w(y, t+u) d\mu^\varphi, & n = 0 \\ \int_{Y^\varphi} 1_{\{\varphi_n(y) < t+u < \varphi_{n+1}(y)\}} v(y, u) w(F^n y, t+u - \varphi_n(y)) d\mu^\varphi, & n \geq 1 \end{cases}.$$

In particular,

$$|J_0(t)| \leq |v|_\infty |w|_\infty \int_{Y^\varphi} 1_{\{\varphi > t\}} d\mu^\varphi = |v|_\infty |w|_\infty |\varphi|_1^{-1} \int_Y 1_{\{\varphi > t\}} \varphi d\mu.$$

For $n \geq 1$,

$$\begin{aligned} \widehat{J}_n(s) &= \int_0^\infty e^{-st} J_n(t) dt \\ &= |\varphi|_1^{-1} \int_Y \int_0^{\varphi(y)} \int_{\varphi_n(y)-u}^{\varphi_{n+1}(y)-u} e^{-st} v(y, u) w(F^n y, t+u - \varphi_n(y)) dt du d\mu. \end{aligned}$$

Making the substitution $u' = t + u - \varphi_n(y)$,

$$\begin{aligned} \widehat{J}_n(s) &= |\varphi|_1^{-1} \int_Y \left(\int_0^{\varphi(y)} e^{su} v(y, u) du \right) \left(\int_0^{\varphi(F^n y)} e^{-su'} w(F^n y, u') du' \right) e^{-s\varphi_n(y)} d\mu \\ &= |\varphi|_1^{-1} \int_Y e^{-s\varphi_n} v_s \widehat{w}(s) \circ F^n d\mu, \end{aligned}$$

as required. ■

Corollary 6.4 *Let $v \in L^1(Y^\varphi)$ and $w \in L^\infty(Y^\varphi)$ with $\int_{Y^\varphi} v d\mu^\varphi = 0$. Then*

$$\widehat{\rho}_{v,w}(s) = \widehat{J}_0(s) + |\varphi|_1^{-1} \int_Y (I - \widehat{R}(s))^{-1} R \widehat{V}(s) \widehat{w}(s) d\mu \quad \text{for all } s \in \mathbb{H}.$$

Proof Note that $e^{-s\varphi} v_s = \widehat{V}(s)$. Hence

$$R^n(e^{-s\varphi_n} v_s) = R^{n-1} R(e^{-s\varphi_{n-1} \circ F} e^{-s\varphi} v_s) = R^{n-1}(e^{-s\varphi_{n-1}} R(e^{-s\varphi} v_s)) = \widehat{R}(s)^{n-1} R \widehat{V}(s),$$

and so

$$\begin{aligned}\widehat{J}_n(s) &= |\varphi|_1^{-1} \int_Y e^{-s\varphi_n} v_s \widehat{w}(s) \circ F^n d\mu \\ &= |\varphi|_1^{-1} \int_Y R^n(e^{-s\varphi_n} v_s) \widehat{w}(s) d\mu = |\varphi|_1^{-1} \int_Y \widehat{R}(s)^{n-1} R\widehat{V}(s) \widehat{w}(s) d\mu.\end{aligned}$$

The result now follows from Proposition 6.3. ■

Remark 6.5 In contrast with the formula in [74], we do not write $\widehat{v}(s)(y) = \int_0^{\varphi(y)} e^{su} v(y, u) du$ since such a function is not analytic on \mathbb{H} . We reserve the hat notation for functions that are analytic on \mathbb{H} , such as \widehat{w} and \widehat{V} .

The following estimate is immediate from the definition of $w(t)$.

Proposition 6.6 $|w(t)|_1 \leq |w|_\infty \mu(\varphi > t)$ for all $w \in L^\infty(Y^\varphi)$, $t > 0$. ■

7 Proof of rapid mixing for semiflows

In this section, we consider Gibbs-Markov semiflows $F_t : Y^\varphi \rightarrow Y^\varphi$ for which the roof function $\varphi : Y \rightarrow \mathbb{R}^+$ lies in $L^q(Y)$ for all $q \geq 1$. For such semiflows, we prove Theorem 3.7, namely that absence of approximate eigenfunctions is a sufficient condition for rapid mixing.

Subsection 7.1 contains background material on Gibbs-Markov maps. The only results we assume without proof are Propositions 7.1 and 7.4 below. Subsections 7.2 and 7.3 establish smoothness of the expressions $R\widehat{V}$ and \widehat{R} that arose in Subsection 6.2. The key estimate, Dolgopyat's estimate, is established in Subsection 7.4 and is used to establish the smoothness of $\widehat{T} = (I - \widehat{R})^{-1}$ in Subsection 7.5 and thereby $\widehat{\rho}_{v,w}$ in Subsection 7.6, completing the proof of Theorem 3.7.

Throughout this section, $q \in \mathbb{N} = \{0, 1, 2, \dots\}$. Also we take $\mathcal{A} = L^\infty(Y^\varphi)$ in Lemma 6.2. If \mathcal{A}_1 and \mathcal{A}_2 are normed vector spaces, we denote by $\mathcal{B}(\mathcal{A}_1, \mathcal{A}_2)$ the space of bounded linear operators from \mathcal{A}_1 to \mathcal{A}_2 , and we write $\mathcal{B}(\mathcal{A}_1)$ instead of $\mathcal{B}(\mathcal{A}_1, \mathcal{A}_1)$.

7.1 Background on Gibbs-Markov maps

Let $F : Y \rightarrow Y$ be a (full branch) Gibbs-Markov map as in Section 3.1 with partition $\{Y_j\}$. For each $n \geq 1$, let \mathcal{D}_n denote the partition of Y into n -cylinders $d = \bigcap_{i=1, \dots, n} F^{-i} Y_{j_i}$, where j_1, \dots, j_n range over positive integers. Define $p_n = \sum_{i=0}^{n-1} p \circ F^i$.

Proposition 7.1 *There is a constant $C_2 \geq 1$ such that*

$$C_2^{-1} \mu(d) \leq e^{p_n(y)} \leq C_2 \mu(d), \quad |e^{p_n(y)} - e^{p_n(y')}| \leq C_2 \mu(d) d_\theta(F^n y, F^n y'), \quad (7.1)$$

for all $y, y' \in d$, $d \in \mathcal{D}_n$, $n \geq 1$.

Proof This is standard, see for example [1, 2]. ■

The transfer operator corresponding to a Gibbs-Markov map F has the pointwise expression $(Rv)(y) = \sum_{j \geq 1} e^{p(y_j)} v(y_j)$, where y_j is the unique preimage of y in Y_j . More

generally, $(R^n v)(y) = \sum_{d \in \mathcal{D}_n} e^{p_n(y_d)} v(y_d)$, $n \geq 1$, where y_d is the unique F^n -preimage of y in d .

Corollary 7.2 *Let $v \in \mathcal{F}_\theta(Y)$. Then*

$$\|R^n v\|_\theta \leq C_2 \sum_{d \in \mathcal{D}_n} \mu(d) (2|1_d v|_\infty + \theta^n |1_d v|_\theta) \quad \text{for } n \geq 1.$$

In particular $\|Rv\|_\theta \leq 2C_2 \sum_{j \geq 1} \mu(Y_j) \|1_{Y_j} v\|_\theta$.

Proof For $y, y' \in Y$, it follows from (7.1) that

$$\begin{aligned} |(R^n v)(y) - (R^n v)(y')| &\leq \sum_d |e^{p_n(y_d)} - e^{p_n(y'_d)}| |v(y_d)| + \sum_d e^{p_n(y'_d)} |v(y_d) - v(y'_d)| \\ &\leq C_2 \sum_d \mu(d) d_\theta(F^n y_d, F^n y'_d) |1_d v|_\infty + C_2 \sum_d \mu(d) |1_d v|_\theta d_\theta(y_d, y'_d) \\ &= C_2 \sum_d \mu(d) d_\theta(y, y') |1_d v|_\infty + C_2 \sum_d \mu(d) |1_d v|_\theta \theta^n d_\theta(y, y'). \end{aligned}$$

Hence $|R^n v|_\theta \leq C_2 \sum_d \mu(d) (|1_d v|_\infty + \theta^n |1_d v|_\theta)$. A simpler argument shows that $|R^n v|_\infty \leq C_2 \sum_d \mu(d) |1_d v|_\infty$. \blacksquare

Remark 7.3 Often we consider operators $S : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$ of the form $Sv = R(gv)$ for some fixed $g \in \mathcal{F}_\theta(Y)$. Since $\|gv\|_\theta \leq \|g\|_\theta \|v\|_\theta$ it follows from Corollary 7.2 that $\|S\|_\theta \leq 2C_2 \sum_{j \geq 1} \mu(Y_j) \|1_{Y_j} g\|_\theta$.

Proposition 7.4 *The operator $R : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$ has spectral radius 1 and essential spectral radius θ . There is a simple eigenvalue at 1 with eigenfunction 1, and no other eigenvalues on the unit circle. Moreover, there are constants $C_3 \geq 1$, $\gamma_1 \in (0, 1)$ such that*

$$\|R^n v - \int_Y v d\mu\|_\theta \leq C_3 \gamma_1^n \|v\|_\theta \quad \text{for all } n \geq 1, v \in \mathcal{F}_\theta(Y).$$

Proof See for example [1, Section 4.7], [2]. \blacksquare

7.2 Smoothness of $R\widehat{V} : \mathcal{F}_\theta(Y^\varphi) \rightarrow \mathcal{F}_\theta(Y)$

Recall that $\widehat{V}(s)(y) = \int_0^{\varphi(y)} e^{-s(\varphi(y)-u)} v(y, u) du$. In this subsection, we obtain estimates for $R\widehat{V}$ as a function of s and the observable $v \in \mathcal{F}_\theta(Y^\varphi)$. For convenience of notation, the dependence on v is kept implicit.

Proposition 7.5 *$R\widehat{V} : \overline{\mathbb{H}} \rightarrow \mathcal{B}(\mathcal{F}_\theta(Y^\varphi), \mathcal{F}_\theta(Y))$ is C^∞ and ²*

$$\|R\widehat{V}^{(q)}(s)\|_\theta \leq (2C_1)^{q+4} C_2 (|s| + q + 1) \int_Y \varphi^{q+2} d\mu \|v\|_\theta,$$

for all $q \in \mathbb{N}$, $s \in \overline{\mathbb{H}}$, $v \in \mathcal{F}_\theta(Y^\varphi)$.

²Throughout, when we estimate derivatives on $\overline{\mathbb{H}}$, it is to be understood that the estimates for $q = 0$ hold on $\overline{\mathbb{H}}$ and the estimates for $q > 0$ hold on the imaginary axis as required in Lemma 6.2.

Proof We have $\widehat{V}^{(q)}(s)(y) = (-1)^q \int_0^{\varphi(y)} e^{-s(\varphi(y)-u)} (\varphi(y) - u)^q v(y, u) du$. In particular, $|1_{Y_j} \widehat{V}^{(q)}(s)|_\infty \leq |1_{Y_j} \varphi|_\infty^{q+1} |v|_\infty$.

Next, let $y, y' \in Y_j$ with $\varphi(y) \geq \varphi(y')$. Then

$$\widehat{V}^{(q)}(s)(y) - \widehat{V}^{(q)}(s)(y') = (-1)^q (I_1 + I_2 + I_3 + I_4),$$

where

$$\begin{aligned} I_1 &= \int_{\varphi(y')}^{\varphi(y)} e^{-s(\varphi(y)-u)} (\varphi(y) - u)^q v(y, u) du, \\ I_2 &= \int_0^{\varphi(y')} \{e^{-s\varphi(y)} - e^{-s\varphi(y')}\} e^{-su} (\varphi(y) - u)^q v(y, u) du, \\ I_3 &= \int_0^{\varphi(y')} e^{-s(\varphi(y')-u)} \{(\varphi(y) - u)^q - (\varphi(y') - u)^q\} v(y, u) du, \\ I_4 &= \int_0^{\varphi(y')} e^{-s(\varphi(y')-u)} (\varphi(y') - u)^q \{v(y, u) - v(y', u)\} du. \end{aligned}$$

Since $v \in \mathcal{F}_\theta(Y^\varphi)$, we obtain

$$\begin{aligned} |I_1| &\leq (\varphi(y) - \varphi(y')) |1_{Y_j} \varphi|_\infty^q |v|_\infty, & |I_2| &\leq |s| (\varphi(y) - \varphi(y')) |1_{Y_j} \varphi|_\infty^{q+1} |v|_\infty, \\ |I_3| &\leq q (\varphi(y) - \varphi(y')) |1_{Y_j} \varphi|_\infty^q |v|_\infty, & |I_4| &\leq |1_{Y_j} \varphi|_\infty^{q+1} \sup_u |v(y, u) - v(y', u)|. \end{aligned}$$

Also, we have the estimates $|\varphi(y) - \varphi(y')| \leq |1_{Y_j} \varphi|_\theta d_\theta(y, y') \leq C_1 |1_{Y_j} \varphi|_\infty d_\theta(y, y')$ and $|v(y, u) - v(y', u)| \leq |1_{Y_j} \varphi|_\infty |v|_\theta d_\theta(y, y')$, so

$$|1_{Y_j} \widehat{V}^{(q)}(s)|_\theta \leq C_1 (|s| + q + 1) |1_{Y_j} \varphi|_\infty^{q+2} \|v\|_\theta.$$

Hence

$$\|1_{Y_j} \widehat{V}^{(q)}(s)\|_\theta \leq C_1 (|s| + q + 2) |1_{Y_j} \varphi|_\infty^{q+2} \|v\|_\theta \leq (2C_1)^{q+3} (|s| + q + 1) \inf_{Y_j} \varphi^{q+2} \|v\|_\theta.$$

The result follows from Corollary 7.2. ■

7.3 Smoothness of $\widehat{R} : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$ and spectral properties

Proposition 7.6 $\widehat{R} : \overline{\mathbb{H}} \rightarrow \mathcal{B}(\mathcal{F}_\theta(Y))$ is C^∞ and

$$\|\widehat{R}^{(q)}(s)\|_\theta \leq C_2 (2C_1)^{q+3} (|s| + q + 1) \int_Y \varphi^{q+1} d\mu \quad \text{for all } q \in \mathbb{N}, s \in \overline{\mathbb{H}}.$$

Proof First note that $\widehat{R}^{(q)}(s)v = (-1)^q R(f_q(s)v)$, where $f_q(s) = e^{-s\varphi} \varphi^q$.

It is immediate that $|1_{Y_j} f_q(s)|_\infty \leq |1_{Y_j} \varphi|_\infty^q$. Next, let $y, y' \in Y_j$. Then $f_q(s)(y) - f_q(s)(y') = I_1 + I_2$ where

$$I_1 = \{e^{-s\varphi(y)} - e^{-s\varphi(y')}\} \varphi(y)^q, \quad I_2 = e^{-s\varphi(y')} \{\varphi(y)^q - \varphi(y')^q\}.$$

These terms contribute

$$|I_1| \leq |s| C_1 |1_{Y_j} \varphi|_\infty^{q+1} d_\theta(y, y'), \quad |I_2| \leq q C_1 |1_{Y_j} \varphi|_\infty^q d_\theta(y, y').$$

It follows that $|1_{Y_j} f_q(s)|_\theta \leq C_1(|s| + q)|1_{Y_j} \varphi|_\infty^{q+1}$ and so

$$\|1_{Y_j} f_q(s)\|_\theta \leq C_1(|s| + q + 1)|1_{Y_j} \varphi|_\infty^{q+1} \leq (2C_1)^{q+2}(|s| + q + 1)\inf_{Y_j} \varphi^{q+1}.$$

By Remark 7.3, $\|\widehat{R}^{(q)}(s)\|_\theta \leq 2C_2 \sum \mu(Y_j) \|1_{Y_j} f_q(s)\|_\theta \leq C_2(2C_1)^{q+3}(|s| + q + 1) \sum \mu(Y_j) \inf_{Y_j} \varphi^{q+1} \leq C_2(2C_1)^{q+3}(|s| + q + 1) \int_Y \varphi^{q+1} d\mu.$ ■

Proposition 7.7 (a) $|\widehat{R}(s)v|_p \leq |v|_p$ for all $s \in \overline{\mathbb{H}}$, $v \in L^p(Y)$, $1 \leq p \leq \infty$.

(b) $|\widehat{R}(s)^n v|_\theta \leq C_4\{(|s| + 1)|v|_\infty + \theta^n |v|_\theta\}$ for all $s \in \overline{\mathbb{H}}$, $n \geq 1$, $v \in \mathcal{F}_\theta(Y)$.

Proof Part (a) is immediate. For part (b), see [27, Corollary 4.3(a)] where it is shown that $|\widehat{R}(ib)^n v|_\theta \leq \{C_2^2 + |b|C_2\theta(1-\theta)^{-1} \sum_{j \geq 1} \mu(Y_j) |1_{Y_j} \varphi|_\theta\} |v|_\infty + C_2\theta^n |v|_\theta$. The proof given there extends immediately to $s \in \overline{\mathbb{H}}$ as follows.

Write $\widehat{R}(s)^n v = R^n(f_n(s)v)$ where $f_n(s) = e^{-s\varphi^n}$. Fix an n -cylinder $d \in \mathcal{D}_n$ and let $y, y' \in d$. Then

$$\begin{aligned} |f_n(s)(y) - f_n(s)(y')| &\leq |s| \sum_{j=0}^{n-1} |\varphi(F^j y) - \varphi(F^j y')| \leq |s| \sum_{j=0}^{n-1} |1_{F^j d} \varphi|_\theta d_\theta(F^j y, F^j y') \\ &\leq C_1 |s| \sum_{j=0}^{n-1} (\inf_{F^j d} \varphi) \theta^{-j} d_\theta(y, y'). \end{aligned}$$

Hence $|1_d f_n(s)|_\infty \leq 1$ and $|1_d f_n(s)|_\theta \leq C_1 |s| \sum_{j=0}^{n-1} (\inf_{F^j d} \varphi) \theta^{-j}$. It follows that $|1_d f_n(s)v|_\infty \leq |v|_\infty$ and

$$|1_d f_n(s)v|_\theta \leq |1_d f_n(s)|_\theta |v|_\infty + |1_d f_n(s)|_\infty |v|_\theta \leq C_1 |s| \sum_{j=0}^{n-1} (\inf_{F^j d} \varphi) \theta^{-j} |v|_\infty + |v|_\theta.$$

Hence by Corollary 7.2,

$$\begin{aligned} \|\widehat{R}(s)^n v\|_\theta &\leq C_2 \sum_d \mu(d) (2|1_d f_n(s)v|_\infty + \theta^n |1_d f_n(s)v|_\theta) \\ &\leq C_2 \sum_d \mu(d) (2|v|_\infty + \theta^n |v|_\theta) + C_2 C_1 |s| |v|_\infty K = (2C_2 + C_2 C_1 |s| K) |v|_\infty + C_2 \theta^n |v|_\theta, \end{aligned}$$

where

$$\begin{aligned} K &= \sum_{d \in \mathcal{D}_n} \mu(d) \sum_{j=0}^{n-1} \theta^{n-j} \inf_{F^j d} \varphi = \sum_{d \in \mathcal{D}_n} \mu(d) \sum_{j=0}^{n-1} \theta^{n-j} \inf_d \varphi \circ F^j \\ &\leq \int_Y \sum_{j=0}^{n-1} \theta^{n-j} \varphi \circ F^j d\mu = \sum_{j=0}^{n-1} \theta^{n-j} \int_Y \varphi d\mu \leq \theta(1-\theta)^{-1} \int_Y \varphi d\mu. \end{aligned}$$

This completes the proof of part (b). ■

Proposition 7.8 (a) $\widehat{R}(ib)$ has spectral radius at most 1 and essential spectral radius at most θ for all $b \in \mathbb{R}$.

(b) The spectral radius of $\widehat{R}(s)$ is less than 1 for all $s \in \mathbb{H}$.

(c) $\widehat{R}(ib)v = \lambda v$ for some $\lambda \in \mathbb{C}$ with $|\lambda| = 1$, $v \in L^2(Y)$, $b \in \mathbb{R}$ if and only if $e^{ib\varphi}v \circ F = \bar{\lambda}v$.

Proof (a) Since $\mathcal{F}_\theta(Y)$ is compactly embedded in $L^\infty(Y)$, the estimate on the essential spectral radius follows from Proposition 7.7(a,b), see for example [55]. Now apply Proposition 7.7(a).

(b) Again the essential spectral radius is strictly less than 1 by Proposition 7.7(a,b). Also, if $\widehat{R}(s)v = \lambda v$ for some $v \in \mathcal{F}_\theta(Y)$, $\lambda \in \mathbb{C}$, $|\lambda| = 1$, then $\int_Y |v| d\mu = \int_Y |R(e^{-s\varphi}v)| d\mu \leq \int_Y e^{-a\varphi}|v| d\mu$ where $a = \operatorname{Re} s > 0$. But $\varphi > 0$ so $v = 0$.

(c) Recall that $M_b v = e^{ib\varphi}v \circ F$. Note that $\widehat{R}(ib)M_b = I$ and that M_b and $\widehat{R}(ib)$ are L^2 adjoints, i.e. $\langle \widehat{R}(ib)v, w \rangle = \int_Y \widehat{R}(ib)v \bar{w} d\mu = \int_Y v \overline{M_b w} d\mu = \langle v, M_b w \rangle$. If $M_b v = \bar{\lambda}v$, then $v = \widehat{R}(ib)M_b v = \bar{\lambda}\widehat{R}(ib)v$ so $\widehat{R}(ib)v = \lambda v$. Conversely, if $\widehat{R}(ib)v = \lambda v$, then

$$\begin{aligned} \langle M_b v - \bar{\lambda}v, M_b v - \bar{\lambda}v \rangle &= \langle M_b v, M_b v \rangle - \langle M_b v, \bar{\lambda}v \rangle - \langle \bar{\lambda}v, M_b v \rangle + \langle \bar{\lambda}v, \bar{\lambda}v \rangle \\ &= \langle v, v \rangle - \langle v, v \rangle - \langle v, v \rangle + \langle v, v \rangle = 0, \end{aligned}$$

so $M_b v = \bar{\lambda}v$. ■

Corollary 7.9 *There exists $\delta > 0$ and a C^∞ family of simple eigenvalues $\lambda : \overline{\mathbb{H}} \cap B_\delta(0) \rightarrow \mathbb{C}$ such that $\lambda(0) = 1$ and $\lambda(s)$ is isolated in $\operatorname{spec} \widehat{R}(s) : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$. Moreover, $\lambda'(0) = -|\varphi|_1$. The corresponding family of spectral projections $P(s)$ is C^∞ with $P(0)v = \int_Y v d\mu$ for all $v \in \mathcal{F}_\theta(Y)$.*

Proof The existence of the C^∞ families $\lambda(s)$ and $P(s)$ follows from Propositions 7.4 and 7.6. Differentiating $\widehat{R}P = \lambda P$ and applying P to both sides,

$$P\widehat{R}'P + P\widehat{R}P' = \lambda PP' + \lambda'P.$$

Since $P\widehat{R} = \widehat{R}P = \lambda P$, this reduces to $P\widehat{R}'P = \lambda'P$. In particular, $\lambda'(0) = P(0)\widehat{R}'(0)1 = \int_Y \widehat{R}'(0)1 d\mu = -\int_Y R\varphi d\mu = -|\varphi|_1$. ■

7.4 Dolgopyat estimate

In this subsection, we prove the following key estimate.

Theorem 7.10 *Assume absence of approximate eigenfunctions. Then*

(a) The spectral radius of $\widehat{R}(s)$ is less than 1 for all $s \in \overline{\mathbb{H}} \setminus \{0\}$.

(b) For any $\delta > 0$, there exist $\alpha, C > 0$ such that

$$\|(I - \widehat{R}(s))^{-1}\|_\theta \leq C|b|^\alpha \quad \text{for all } s = a + ib \in \mathbb{C} \text{ with } 0 \leq a \leq 1, |b| \geq \delta.$$

Define the scale of equivalent norms

$$\|v\|_b = \max \left\{ |v|_\infty, \frac{|v|_\theta}{2C_4(|b| + 2)} \right\}, \quad b \in \mathbb{R}.$$

By Proposition 7.7(a,b), for all $n \geq 1$,

$$\|\widehat{R}(s)^n\|_b \leq C_4 + \frac{1}{2} \quad \text{for all } s = a + ib \in \mathbb{C} \text{ with } a \in [0, 1].$$

For each b , define the unit ball $\mathcal{F}_\theta(Y)_b = \{v \in \mathcal{F}_\theta : \|v\|_b \leq 1\}$. Let $C_5 = 2C_4(C_4 + \frac{1}{2})$. Then for all $v \in \mathcal{F}_\theta(Y)_b$, $n \geq 1$,

$$|\widehat{R}(s)^n v|_\infty \leq 1 \quad \text{and} \quad |\widehat{R}(s)^n v|_\theta \leq C_5(|b| + 2) \quad \text{for } s = a + ib \in \mathbb{C} \text{ with } a \in [0, 1].$$

Let Z denote a fixed subset of Y consisting of a finite union of partition elements of Y , with finite subsystem $Z_0 = \bigcap_{j \geq 0} F^{-j}Z$. Note that p is uniformly bounded on Z_0 and moreover $|p_n(y)| \leq n|1_{Z_0}p|_\infty$ for all $y \in Z_0$ and $n \geq 1$.

Lemma 7.11 *Fix $\alpha_2 > 0$. Then there exists $\alpha_1 > 0$ and there exists $\xi > 0$ arbitrarily large such that the following is true for each fixed $s = a + ib \in \mathbb{C}$ with $a \in [0, 1]$, setting $n(b) = \lceil \xi \ln(|b| + 2) \rceil$:*

Suppose that there exists $v_0 \in \mathcal{F}_\theta(Y)_b$ such that for all $y \in Z_0$ and all $j = 0, 1, 2$,

$$|(\widehat{R}(s)^{jn(b)} v_0)(y)| \geq 1 - (|b| + 2)^{-\alpha_1}.$$

Then there exists $w \in \mathcal{F}_\theta(Y)$ with $|w(y)| \equiv 1$ and $|w|_\theta \leq 8C_5(|b| + 2)$, and there exists $\psi \in [0, 2\pi)$ such that for all $y \in Z_0$,

$$|(M_b^{n(b)} w)(y) - e^{i\psi} w(y)| \leq 8(|b| + 2)^{-\alpha_2}.$$

Proof We write $\hat{b} = |b| + 2$ and $n = n(b)$. Choose ξ such that $8C_5 \hat{b} \theta^n = 8C_5 \hat{b} \theta^{\lceil \xi \ln \hat{b} \rceil} \leq \hat{b}^{-\alpha_2}$ for all b , and set

$$\alpha_1 = \max\{1, 2\alpha_2 + \xi|1_{Z_0}p|_\infty\}.$$

Write $v_j = \widehat{R}(s)^{jn} v_0$ and $v_j = r_j w_j$, where $|w_j(y)| \equiv 1$ and $1 - \hat{b}^{-\alpha_1} \leq r_j(y) \leq 1$ for $y \in Z_0$. Note that $|r_j|_\theta \leq |v_j|_\theta \leq C_5 \hat{b}$, so

$$|w_j|_\theta = |r_j^{-1} v_j|_\theta \leq 2C_5 \hat{b} (1 - \hat{b}^{-\alpha_1})^{-2} \leq 8C_5 \hat{b}.$$

Rearrange $v_1 = \widehat{R}(s)^n v_0$ to obtain $w_1^{-1} \widehat{R}(s)^n (v_0) = r_1 \geq 1 - \hat{b}^{-\alpha_1}$. Hence $1 - w_1^{-1} \widehat{R}(s)^n (v_0) \leq \hat{b}^{-\alpha_1}$. It follows that

$$\begin{aligned} R^n(1 - \operatorname{Re}\{e^{-ib\varphi_n} w_0 w_1^{-1} \circ F^n\}) &\leq 1 - R^n(r_0 e^{-a\varphi_n} \operatorname{Re}\{e^{-ib\varphi_n} w_0 w_1^{-1} \circ F^n\}) \\ &= 1 - \operatorname{Re} R^n(e^{-s\varphi_n} v_0 w_1^{-1} \circ F^n) = \operatorname{Re}\{1 - w_1^{-1} \widehat{R}(s)^n (v_0)\} \leq \hat{b}^{-\alpha_1}. \end{aligned}$$

Hence

$$e^{pn(y)} [1 - \operatorname{Re}(e^{-ib\varphi_n(y)} w_0(y) w_1^{-1}(F^n y))] \leq \hat{b}^{-\alpha_1}$$

for all $y \in Y$ with $F^n y \in Z_0$. It follows that

$$|e^{-ib\varphi_n(y)} w_0(y) - w_1(F^n y)| \leq 2(e^{-pn(y)} \hat{b}^{-\alpha_1})^{1/2}.$$

Similarly, with w_0 and w_1 replaced by w_1 and w_2 . Restricting to $y \in Z_0$, we have $e^{-p_n(y)}\hat{b}^{-\alpha_1} \leq \hat{b}^{-2\alpha_2}$ and hence

$$|e^{-ib\varphi_n(y)}w_0(y) - w_1(F^n y)| \leq 2\hat{b}^{-\alpha_2}, \quad |e^{-ib\varphi_n(y)}w_1(y) - w_2(F^n y)| \leq 2\hat{b}^{-\alpha_2}, \quad (7.2)$$

for all $y \in Z_0$. Fix $z \in Z_0$ and choose $\psi_0, \psi_1 \in \mathbb{R}$ such that $w_j(z) = e^{i\psi_j}$ for $j = 0, 1$ and such that $\psi = \psi_0 - \psi_1 \in [0, 2\pi)$. To each y , we associate the point $y^* \in Z_0$ with symbol sequence $y^* = z_0 \cdots z_{n-1}y_n y_{n+1} \cdots$ (recall from Definition 3.3 that $F|_{Z_0}$ is a full one-sided shift). Then y^* is within distance θ^n of z and $F^n y^* = F^n y$. We obtain

$$\begin{aligned} |e^{-ib\varphi_n(y^*)}e^{i\psi_0} - w_1(F^n y)| &\leq 2\hat{b}^{-\alpha_2} + 8C_5\hat{b}\theta^n \leq 3\hat{b}^{-\alpha_2} \\ |e^{-ib\varphi_n(y^*)}e^{i\psi_1} - w_2(F^n y)| &\leq 2\hat{b}^{-\alpha_2} + 8C_5\hat{b}\theta^n \leq 3\hat{b}^{-\alpha_2}, \end{aligned}$$

by the choice of ξ . Hence $|e^{-i\psi}w_1(F^n y) - w_2(F^n y)| \leq 6\hat{b}^{-\alpha_2}$. Substituting into (7.2) yields the required approximate eigenfunction $w = w_1$. \blacksquare

Lemma 7.12 *Let $\alpha_1, \xi_1 > 0$. Suppose that for any $v \in \mathcal{F}_\theta(Y)_b$ there exists $y_0 \in Z_0$ and $j \leq [\xi_1 \ln(|b| + 2)]$ such that*

$$|\widehat{R}(s)^j v(y_0)| \leq 1 - (|b| + 2)^{-\alpha_1} \quad \text{for all } s = a + ib \in \mathbb{C} \text{ with } a \in [0, 1].$$

Then there exists $\alpha, \xi, \epsilon > 0, C > 0$ such that

- (a) $\|\widehat{R}(s)^{[\xi \ln(|b| + 2)]}\|_b \leq 1 - \epsilon(|b| + 2)^{-\alpha}$ for all $s = a + ib \in \mathbb{C}$ with $a \in [0, 1]$.
- (b) $\|(I - \widehat{R}(s))^{-1}\|_\theta \leq C(|b| + 2)^\alpha$ for all $s = a + ib \in \mathbb{C}$ with $a \in [0, 1]$.

Proof We use the pointwise estimate on iterates of $\widehat{R}(s)$ to obtain estimates on the L^1 , L^∞ and $\|\cdot\|_b$ norms. Set $\hat{b} = |b| + 2$.

Let $\hat{u} = \widehat{R}(s)^j v$ and note that $|\hat{u}|_\infty \leq 1$, $|\hat{u}|_\theta \leq C_5\hat{b}$. Hence, $|\hat{u}(y)| \leq 1 - 1/(2\hat{b}^{\alpha_1})$ for all y within distance $1/(2C_5\hat{b}^{\alpha_1+1})$ of y_0 . Call this subset U . If $d \in \mathcal{D}_k$ is a k -cylinder, then $\text{diam } d = \theta^k$, so provided $\theta^k < 1/(2C_5\hat{b}^{\alpha_1+1})$, the k -cylinder containing y_0 lies inside U . It suffices to take $k \approx (\alpha_1 + 1) \ln \hat{b}/(-\ln \theta)$. By (7.1),

$$\mu(U) \geq \mu(d) \geq C_2^{-1}e^{-p_k(y_0)} \geq C_2^{-1}e^{-k|1_{Z_0}p|_\infty} \geq \epsilon_1\hat{b}^{-(\alpha_1+1)\alpha_2},$$

where $\alpha_2 = |1_{Z_0}p|_\infty/(-\ln \theta)$.

Next, let $u = \widehat{R}(s)^{n_1(b)}v$ where $n_1(b) = [\xi_1 \ln \hat{b}]$. Again, $|u|_\infty \leq 1$, $|u|_\theta \leq C_5\hat{b}$. Also, $n_1(b) \geq j$ so $|u|_1 \leq |\hat{u}|_1$ by Proposition 7.7(a). Breaking up Y into U and $Y \setminus U$,

$$|u|_1 \leq |\hat{u}|_1 \leq (1 - 1/(2\hat{b}^{\alpha_1}))\mu(U) + 1 - \mu(U) = 1 - \mu(U)/(2\hat{b}^{\alpha_1}) \leq 1 - \epsilon_2\hat{b}^{-\alpha},$$

where $\alpha = \alpha_1 + \alpha_2 + \alpha_1\alpha_2$. By Proposition 7.4, and using that $\| |u| \|_\theta \leq \|u\|_\theta$,

$$\begin{aligned} |\widehat{R}(s)^n u|_\infty &\leq |R^n(|u|)|_\infty \leq |R^n(|u|) - \int_Y |u| d\mu|_\infty + |u|_1 \leq C_3\gamma_1^n \|u\|_\theta + |u|_1 \\ &\leq (1 + C_5\hat{b})C_3\gamma_1^n + 1 - \epsilon_2\hat{b}^{-\alpha}. \end{aligned}$$

Choosing $n = n_2(b) = \lceil \xi_2 \ln \hat{b} \rceil$ where $\xi_2 \gg 1$ ensures that

$$|\widehat{R}(s)^{n_1(b)+n_2(b)}v|_\infty = |\widehat{R}(s)^{n_2(b)}u|_\infty \leq 1 - \epsilon \hat{b}^{-\alpha}.$$

Setting $n(b) = \lceil \xi \ln \hat{b} \rceil$ where $\xi = \xi_1 + \xi_2$,

$$|\widehat{R}(s)^{n(b)}v|_\infty \leq 1 - \epsilon \hat{b}^{-\alpha}.$$

By Proposition 7.7(a,b), $|\widehat{R}(s)^{n(b)+n}v|_\infty \leq 1 - \epsilon \hat{b}^{-\alpha}$ for all $n \geq 0$, and

$$|\widehat{R}(s)^{n(b)+n}v|_\theta / (2C_4 \hat{b}) \leq C_4 \{\hat{b} + \theta^n C_5 \hat{b}\} / (2C_4 \hat{b}) \leq \frac{1}{2} + \frac{1}{2} C_5 \theta^n \leq \frac{3}{4},$$

for n sufficiently large (independent of b). Increasing ξ slightly, $\|\widehat{R}(s)^{n(b)}v\|_b \leq 1 - \epsilon \hat{b}^{-\alpha}$ proving part (a).

It follows that $\|(I - \widehat{R}(s)^{n(b)})^{-1}\|_b \leq \epsilon^{-1} \hat{b}^\alpha$. Using the identity $(I - A)^{-1} = (I + A + \dots + A^{m-1})(I - A^m)^{-1}$,

$$\|(I - \widehat{R}(s))^{-1}\|_b \leq \sum_{j=0}^{n(b)-1} \|\widehat{R}(s)^j\|_b \|(I - \widehat{R}(s)^{n(b)})^{-1}\|_b \leq \xi \ln \hat{b} (C_4 + \frac{1}{2}) \epsilon^{-1} \hat{b}^\alpha = O(\hat{b}^{\alpha+1}).$$

Hence $\|(I - \widehat{R}(s))^{-1}\|_\theta = O(\hat{b}^{\alpha+2})$. Increasing α , we obtain part (b). \blacksquare

Proof of Theorem 7.10 By Lemmas 7.11 and 7.12(b), there exists $b_0 > 0$ such that that part (b) of the theorem holds for all $|b| \geq b_0$. Moreover, by Proposition 7.8(b) and Lemma 7.12(a), the spectral radius of $\widehat{R}(s)$ is less than one for all $s \in \mathbb{H}$ and all $s = ib$ with $|b| \geq b_0$.

Suppose that the spectral radius of $\widehat{R}(ib)$ is 1 for some $b \in \mathbb{R} \setminus \{0\}$. Then $\widehat{R}(ib)v = \lambda v$ for some $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ and some nonzero $v \in \mathcal{F}_\theta(Y)$. By Proposition 7.8(c), $e^{ib\varphi}v \circ F = \bar{\lambda}v$, so $e^{imb\varphi}v^m \circ F = \bar{\lambda}^m v^m$ for all $m \geq 1$. Again by Proposition 7.8(c), $\widehat{R}(imb)v^m = \lambda^m v$ so the spectral radius of $\widehat{R}(imb)$ is equal to 1 for all $m \geq 1$. Choosing m so that $m|b| \geq b_0$, we obtain a contradiction, hence proving part (a).

Finally, by part (a) and continuity of \widehat{R} (Proposition 7.6), part (b) holds for $|b| \in [\delta, b_0]$. \blacksquare

7.5 Smoothness of $\widehat{T} = (I - \widehat{R})^{-1} : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$

Proposition 7.13 *Assume absence of approximate eigenfunctions. Then $\widehat{T} : \overline{\mathbb{H}} \setminus \{0\} \rightarrow \mathcal{B}(\mathcal{F}_\theta(Y))$ is C^∞ . Moreover, for each $q \in \mathbb{N}$, $\delta > 0$, there exists $\alpha, C > 0$ such that $\|\widehat{T}^{(q)}(s)\|_\theta \leq C|b|^\alpha$ for all $s = a + ib \in \overline{\mathbb{H}}$ with $0 \leq a \leq 1$, $|b| \geq \delta$.*

Proof By Proposition 7.8(b) and Theorem 7.10(a), $1 \notin \text{spec } \widehat{R}(s)$ for $s \in \overline{\mathbb{H}} \setminus \{0\}$. Hence $\widehat{T}(s)$ is a bounded operator on $\mathcal{F}_\theta(Y)$ for all $s \in \overline{\mathbb{H}} \setminus \{0\}$. By Proposition 7.6, $s \mapsto \widehat{R}(s)$ is C^∞ on $\overline{\mathbb{H}}$ so $s \mapsto \widehat{T}(s)$ is C^∞ on $\overline{\mathbb{H}} \setminus \{0\}$.

By induction, $\widehat{T}^{(q)}$ is a finite linear combination of finite products of factors of the form \widehat{T} and $\widehat{R}^{(j)}$, $1 \leq j \leq q$. Each of these is $O(|b|^\alpha)$ for some $\alpha > 0$ by Proposition 7.6 and Theorem 7.10. \blacksquare

Let $\mathcal{F}_\theta(Y^\varphi)^0 = \{v \in \mathcal{F}_\theta(Y^\varphi) : \int_{Y^\varphi} v d\mu^\varphi = 0\}$. As shown in the next result, $\widehat{T}\widehat{R}\widehat{V}$ extends smoothly from $\overline{\mathbb{H}} \setminus \{0\}$ to $\overline{\mathbb{H}}$ when restricted to $\mathcal{F}_\theta(Y^\varphi)^0$.

Corollary 7.14 *Assume absence of approximate eigenfunctions. Then $\widehat{TRV} : \mathbb{H} \rightarrow \mathcal{B}(\mathcal{F}_\theta(Y^\varphi)^0, \mathcal{F}_\theta(Y))$ is C^∞ .*

Proof By Propositions 7.5 and 7.13, it suffices to work on $\mathbb{H} \cap B_\delta(0)$ for some $\delta > 0$. By Corollary 7.9, we can choose $\delta > 0$ so that

$$\widehat{TRV} = (1 - \lambda)^{-1}PR\widehat{V} + Q_1,$$

where $s \mapsto Q_1(s) : \mathcal{F}_\theta(Y^\varphi) \rightarrow \mathcal{F}_\theta(Y)$ is C^∞ . Also $(1 - \lambda(s))^{-1} = s^{-1}|\varphi|_1^{-1}(1 + s\tilde{\lambda}(s))$, $P(s) = P(0) + s\tilde{P}(s)$, $\widehat{V}(s) = \widehat{V}(0) + s\tilde{V}(s)$ where $\tilde{\lambda} \in \mathbb{R}$, $\tilde{P} : \mathcal{F}_\theta(Y) \rightarrow \mathcal{F}_\theta(Y)$, $\tilde{V} : \mathcal{F}_\theta(Y^\varphi) \rightarrow \mathcal{F}_\theta(Y)$ are C^∞ functions of s . Hence

$$\widehat{T}(s)R\widehat{V}(s) = s^{-1}|\varphi|_1^{-1}P(0)R\widehat{V}(0) + Q_2,$$

where $s \mapsto Q_2(s) : \mathcal{F}_\theta(Y^\varphi) \rightarrow \mathcal{F}_\theta(Y)$ is C^∞ .

Finally, restricting to $\mathcal{F}_\theta(Y^\varphi)^0$,

$$P(0)R\widehat{V}(0) = \int_Y R\widehat{V}(0) d\mu = \int_Y \widehat{V}(0) d\mu = \int_Y \int_0^{\varphi(y)} v(y, u) du d\mu = |\varphi|_1 \int_{Y^\varphi} v d\mu^\varphi = 0.$$

This completes the proof. ■

7.6 Proof of Theorem 3.7

By Corollary 6.4, $\hat{\rho}_{v,w} = \widehat{J}_0 + |\varphi|_1^{-1} \int_Y \widehat{TRV} \widehat{w} d\mu$.

Since $\varphi \in L^p(Y)$ for all $p \geq 1$, it follows from the Cauchy-Schwarz and Markov inequalities that $\int_Y 1_{\{\varphi > t\}} \varphi d\mu \leq (\mu(\varphi > t))^{1/2} |\varphi|_2 \leq (|\varphi|_p t^{-p})^{1/2} |\varphi|_2 \ll t^{-p/2}$ for all $p \geq 1$. By Propositions 6.3 and 6.6, for each $q \in \mathbb{N}$ there exists $C > 0$ such that

$$|\widehat{J}_0^{(q)}(s)| \leq C|v|_\infty|w|_\infty \leq C\|v\|_\theta\|w\|_\infty \quad \text{and} \quad |\widehat{w}^{(q)}(s)|_1 \leq C|w|_\infty,$$

for all $v \in \mathcal{F}_\theta(Y^\varphi)$, $w \in L^\infty(Y^\varphi)$, $s \in \mathbb{H}$.

By Propositions 7.5 and 7.13 and Corollary 7.14, for each $q \in \mathbb{N}$, there exist $C, \alpha > 0$ such that

$$|(\widehat{TRV})^{(q)}(s)|_\infty \leq \|(\widehat{TRV})^{(q)}(s)\|_\theta \leq C(|b| + 1)^\alpha \|v\|_\theta,$$

and so

$$|\hat{\rho}_{v,w}^{(q)}(s)| \leq C(|b| + 1)^\alpha \|v\|_\theta \|w\|_\infty,$$

for all $v \in \mathcal{F}_\theta(Y^\varphi)^0$, $w \in L^\infty(Y^\varphi)$, $s = a + ib \in \mathbb{C}$ with $a \in [0, 1]$. Hence we have verified the hypotheses of Lemma 6.2. Consequently $|\rho_{v,w}(t)| = O(\|v\|_\theta \|w\|_{\infty, m} t^{-q})$ for all $q \in \mathbb{N}$, completing the proof of Theorem 3.7.

8 Proof of polynomial mixing for semiflows

In this section, we consider Gibbs-Markov semiflows $F_t : Y^\varphi \rightarrow Y^\varphi$ for which the roof function $\varphi : Y \rightarrow \mathbb{R}^+$ satisfies $\mu(\varphi > t) = O(t^{-\beta})$ for some $\beta > 1$. For such semiflows, we prove Theorem 3.6, namely that absence of approximate eigenfunctions is a sufficient condition to obtain the mixing rate $O(t^{-(\beta-1)})$.

The assumption on φ implies that $\varphi \in L^q(Y)$ for all $q < \beta$ but in general $\varphi \notin L^\beta(Y)$. Nonintegrability of $\rho_{v,w}$ (*a priori* for $\beta > 2$ and even *a posteriori* for $\beta \in (1, 2]$) makes inversion of the Laplace transform problematic.

To circumvent this, we use a truncation idea from [64]. The truncated semiflows are rapid mixing by Section 7 and all components of the Laplace transform are C^∞ . The approach in [64] allows for control of the errors that come from truncation.

In Subsection 8.1, we introduce some notation and recall some elementary properties of Fourier transforms and convolutions. Subsections 8.2 and 8.3 contain various refinements of the estimates in Section 7. In Subsection 8.4, we reduce to truncated semiflows. The Laplace transform for truncated semiflows is studied for small b and large b in Subsections 8.5 and 8.6 respectively.

8.1 Some conventions

From now on, we allow q to take noninteger values. (Eventually, we require $q \in (1, \beta)$ with $q \geq \beta - 1$. Hence simplified proofs are available for $\beta > 2$, though certain estimates for intermediate results become less sharp.)

As usual, a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be C^q if f is $C^{[q]}$ and $f^{([q])}$ is $(q - [q])$ -Hölder. Moreover, we write $|f^{(q)}| \leq g$ for some function $g : \mathbb{R} \rightarrow [0, \infty)$ if for all $b, b' \in \mathbb{R}$,

$$|f^{(k)}(b)| \leq g(b), \quad k = 0, 1, \dots, [q], \quad \text{and} \quad |f^{([q]}(b) - f^{([q]}(b'))| \leq (g(b) + g(b'))|b - b'|^{q-[q]}.$$

Definition 8.1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be integrable. We write $f \in \mathcal{R}(a(t))$ if the inverse Fourier transform of f is $O(a(t))$. We use the same notation for Banach space valued functions.

Proposition 8.2 Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be an integrable function such that $g(b) \rightarrow 0$ as $b \rightarrow \pm\infty$. If $|f^{(q)}| \leq g$, then $f \in \mathcal{R}(t^{-q})$.

Proof Let S denote the inverse Fourier transform of f . Write $q = k + r$ where $k = [q]$ and $r \in [0, 1)$. Up to a multiplicative constant, $S(t) = \int_{-\infty}^{\infty} e^{ibt} f(b) db$. Integrating by parts,

$$S(t) = (it)^{-1} e^{ibt} f(b) \Big|_{b=-\infty}^{b=\infty} - (it)^{-1} \int_{-\infty}^{\infty} e^{ibt} f'(b) db = (-it)^{-1} \int_{-\infty}^{\infty} e^{ibt} f'(b) db.$$

Inductively, $S(t) = (-it)^{-k} \int_{-\infty}^{\infty} e^{ibt} f^{[k]}(b) db$. In particular, $|S(t)| \leq t^{-k} \int g \ll t^{-k}$ proving the result when $q = k$ is an integer.

Next, $S(t) = (-it)^{-k} \int_{-\infty}^{\infty} -e^{ibt} f^{[k]}(b + \pi/t) db$ so $S(t) = \frac{1}{2}(-it)^{-k} \int_{-\infty}^{\infty} e^{ibt} \{f^{[k]}(b) - f^{[k]}(b + \pi/t)\} db$. Hence

$$|S(t)| \leq \frac{1}{2} t^{-k} \int_{-\infty}^{\infty} \left(g(b) + g\left(b + \frac{\pi}{t}\right) \right) \left(\frac{\pi}{t} \right)^r db \leq \pi^r t^{-q} \int g \ll t^{-q},$$

as required. ■

Definition 8.3 Let $f, g : [0, \infty) \rightarrow \mathbb{R}$ be integrable. The *convolution* $f \star g$ is defined to be $(f \star g)(t) = \int_0^t f(x)g(t-x) dx$.

Proposition 8.4 Fix $b > a > 0$ with $b > 1$. Suppose that $f, g : [0, \infty) \rightarrow \mathbb{R}$ are integrable and there exists a constant $C > 0$ such that $|f(t)| \leq C(1+t)^{-a}$ and $|g(t)| \leq C(1+t)^{-b}$ for $t \geq 0$. Then there exists a constant $K > 0$ depending only on a and b such that

$$|(f \star g)(t)| \leq C^2 K (1+t)^{-a} \quad \text{for } t \geq 0.$$

Proof Write the convolution as a sum of two integrals

$$I_1(t) = \int_0^{t/2} f(x)g(t-x) dx, \quad I_2(t) = \int_{t/2}^t f(x)g(t-x) dx.$$

Note that $|I_1(t)| \leq C^2(1+t/2)^{-b} \int_0^{t/2} (1+x)^{-a} dx$ and $|I_2(t)| \leq C^2(1+t/2)^{-a} \int_{t/2}^t (1+x)^{-b} dx$. Clearly we can restrict attention to $t \geq 1$. Since $b > 1$, it follows that $|I_2(t)| \ll C^2 t^{-a}$ and $|I_1(t)| \ll C^2 t^{-b}(1+t^{1-a}) \leq 2C^2 t^{-a}$. ■

8.2 Refined estimates for $J_0, w_0, R\widehat{V}$ and \widehat{R} .

Throughout the remainder of this section, we fix

$$\max\{1, \beta - 1\} < q < \beta.$$

Let $\eta \in (0, 1]$ be as in the statement of Theorem 3.6. Shrinking η if needed, we may suppose without loss that

$$q + 2\eta < \beta.$$

Let $\theta_1 = \theta^\eta$. Since $\theta_1 \geq \theta$, condition (3.1) holds also with θ replaced by θ_1 . Hence results from Section 7, for example Theorem 7.10, hold also in $\mathcal{F}_{\theta_1}(Y)$.

We begin by giving improved estimates for $J_0(t)$ and $w(t)$.

Proposition 8.5 Let $\xi \in (0, \beta)$. There exists a constant $C > 0$ such that $\int_Y 1_{\{\varphi > t\}} \varphi^\xi d\mu \leq Ct^{-(\beta-\xi)}$ for all $t \geq 0$.

Proof Let $G(x) = \mu(\varphi \leq x)$. Then

$$\begin{aligned} \int_Y 1_{\{\varphi > t\}} \varphi^\xi d\mu &= \int_t^\infty x^\xi dG(x) = - \int_t^\infty x^\xi d(1 - G(x)) \\ &= -x^\xi(1 - G(x)) \Big|_{x=t}^{x=\infty} + \xi \int_t^\infty x^{\xi-1}(1 - G(x)) dx \ll t^{-(\beta-\xi)} + \int_t^\infty x^{-\beta-1+\xi} dx \ll t^{-(\beta-\xi)}, \end{aligned}$$

as required. ■

Corollary 8.6 $|J_0(t)| = O(|v|_\infty |w|_\infty t^{-(\beta-1)})$ and $|w(t)|_1 = O(|w|_\infty t^{-\beta})$ for all $v, w \in L^\infty(Y^\varphi)$.

Proof This is immediate from Proposition 8.5 together with Propositions 6.3 and 6.6. ■

Next, we mention an improvement to Proposition 7.6.

Proposition 8.7 $\widehat{R} : i\mathbb{R} \rightarrow \mathcal{B}(\mathcal{F}_{\theta_1}(Y))$ is C^q . Indeed,

$$\|\widehat{R}^{(q)}(ib)\|_{\theta_1} \leq C_2(2C_1)^{q+4}(|b| + q + 1) \int_Y \varphi^{q+\eta} d\mu \quad \text{for all } b \in \mathbb{R}.$$

Proof The proof is a refinement of that for Proposition 7.6. Write $\widehat{R}(ib)v = R(f(b)v)$, where $f(b) = e^{-ib\varphi}$. We claim that

$$|1_{Y_j} f^{(q)}(b)|_\infty \leq 2|1_{Y_j} \varphi|_\infty^q, \quad |1_{Y_j} f^{(q)}(b)|_{\theta_1} \leq 2C_1(2|b| + q + 1)|1_{Y_j} \varphi|_\infty^{q+\eta}. \quad (8.1)$$

It then follows that $\|1_{Y_j} f^{(q)}(b)\|_{\theta_1} \leq 2C_1(2|b| + q + 2)|1_{Y_j} \varphi|_\infty^{q+\eta} \leq (2C_1)^{q+3}(|b| + q + 1)\inf_{Y_j} \varphi^{q+\eta}$. Now apply Remark 7.3.

It remains to prove the claim. Write $q = k + r$ where $k = [q]$ and $r \in [0, 1)$. Then $|1_{Y_j} f^{(k)}(b)|_\infty \leq |1_{Y_j} \varphi|_\infty^k$. Hence $|1_{Y_j} \{f^{(k)}(b+h) - f^{(k)}(b)\}|_\infty \leq 2|1_{Y_j} \varphi|_\infty^k$. Also $|1_{Y_j} f^{(k+1)}(b)|_\infty \leq |1_{Y_j} \varphi|_\infty^{k+1}$ so it follows from the mean value theorem that $|1_{Y_j} \{f^{(k)}(b+h) - f^{(k)}(b)\}|_\infty \leq |1_{Y_j} \varphi|_\infty^{k+1}|h|$. Combining these two estimates and using the inequality $\min\{1, x\} \leq x^r$ which holds for all $x \geq 0$, $r \in [0, 1]$,

$$|1_{Y_j} \{f^{(k)}(b+h) - f^{(k)}(b)\}|_\infty \leq 2|1_{Y_j} \varphi|_\infty^k \min\{1, |1_{Y_j} \varphi|_\infty |h|\} \leq 2|1_{Y_j} \varphi|_\infty^{k+r} |h|^r,$$

yielding the first part of (8.1).

Next, for $y, y' \in Y_j$, we have $f^{(k)}(b)(y) - f^{(k)}(b)(y') = (-i)^k (I_1 + I_2)$ where

$$I_1 = \{e^{-ib\varphi(y)} - e^{-ib\varphi(y')}\} \varphi(y)^k, \quad I_2 = e^{-ib\varphi(y')} \{\varphi(y)^k - \varphi(y')^k\}.$$

We have $|I_1| \leq |b|C_1|1_{Y_j} \varphi|_\infty^{k+1} d_\theta(y, y')$ and $|I_2| \leq kC_1|1_{Y_j} \varphi|_\infty^k d_\theta(y, y')$. Also, $|I_1| \leq 2|1_{Y_j} \varphi|_\infty^k$, so

$$\begin{aligned} |I_1| &\leq 2|b|C_1|1_{Y_j} \varphi|_\infty^k \min\{1, |1_{Y_j} \varphi|_\infty d_\theta(y, y')\} \\ &\leq 2|b|C_1|1_{Y_j} \varphi|_\infty^{k+\eta} d_\theta(y, y')^\eta = 2|b|C_1|1_{Y_j} \varphi|_\infty^{k+\eta} d_{\theta_1}(y, y'). \end{aligned}$$

Hence $g^{(k)}(b) = 1_{Y_j} \{f^{(k)}(b)(y) - f^{(k)}(b)(y')\}$ satisfies

$$|g^{(k)}(b)| \leq C_1(2|b| + k)|1_{Y_j} \varphi|_\infty^{k+\eta} d_{\theta_1}(y, y').$$

Now we repeat the mean value theorem argument above to obtain

$$\begin{aligned} |g^{(k)}(b+h) - g^{(k)}(b)| &\leq 2C_1(2|b| + k + 1)|1_{Y_j} \varphi|_\infty^{k+\eta} d_{\theta_1}(y, y') \min\{1, |1_{Y_j} \varphi|_\infty |h|\} \\ &\leq 2C_1(2|b| + k + 1)|1_{Y_j} \varphi|_\infty^{k+r+\eta} d_{\theta_1}(y, y') |h|^r. \end{aligned}$$

Hence $|1_{Y_j} \{f^{(q)}(b)(y) - f^{(q)}(b)(y')\}| \leq 2C_1(2|b| + q + 1)|1_{Y_j} \varphi|_\infty^{q+\eta} d_{\theta_1}(y, y')$, yielding the second part of (8.1). \blacksquare

Remark 8.8 Clearly, the estimate for $\widehat{R}^{(q)}$ holds equally for $\widehat{R}^{(q')}$ for all $q' < q$. We use this observation without comment throughout.

Remark 8.9 During this section, we obtain many estimates of the form $|f^{(k)}(b)| \ll \varphi^{k+\ell}$ for all $k \in \mathbb{N}$. By the mean value theorem argument used in the proof of Proposition 8.7, it follows that $|f^{(q)}(b)| \ll \varphi^{q+\ell}$ for all $q \in [0, \infty)$. From now on, we write ‘‘by the MVT argument’’ and omit the details.

Corollary 8.10 *Assume absence of approximate eigenfunctions. Then $\widehat{T} = (I - \widehat{R})^{-1} : i\mathbb{R} \setminus \{0\} \rightarrow \mathcal{B}(\mathcal{F}_{\theta_1}(Y))$ is C^q . Moreover, for all $\delta > 0$, there exists $\alpha, C > 0$ such that $\|\widehat{T}^{(q)}(ib)\|_{\theta_1} \leq C|b|^\alpha$ for all $b \in \mathbb{R}$ with $|b| \geq \delta$.*

Proof The proof of the Dolgopyat estimate in Theorem 7.10 is completely unchanged (Proposition 7.7(b) used only the integrability of φ). Hence \widehat{T} is C^q by Proposition 8.7.

Let $k \in \mathbb{N}$ with $k < \beta$. By induction, $\widehat{T}^{(k)}$ is a finite linear combination of finite products of factors of the form \widehat{T} and $\widehat{R}^{(j)}$, $1 \leq j \leq k$. Each of these is $O(|b|^\alpha)$ for some $\alpha > 0$ by Theorem 7.10 and Proposition 8.7. Hence there exist constants $C, \alpha > 0$ such that $\|\widehat{T}^{(k)}(ib)\|_{\theta_1} \leq C|b|^\alpha$ for each $k = 0, \dots, [q]$.

Next, write $q = k + r$ where $k = [q]$ and $r \in [0, 1)$. By the resolvent identity together with Theorem 7.10 and Proposition 8.7,

$$\|\widehat{T}(ib) - \widehat{T}(ib')\|_{\theta_1} \leq \|\widehat{T}(ib)\|_{\theta_1} \|\widehat{R}(ib) - \widehat{R}(ib')\|_{\theta_1} \|\widehat{T}(ib')\|_{\theta_1} \leq C^3 |b|^{3\alpha} |b - b'|^r,$$

so $\|\widehat{T}^{(r)}(ib)\|_{\theta_1} \leq C^3 |b|^{3\alpha}$.

Finally, $\widehat{T}^{(q)}$ is a finite linear combination of finite products of factors of the form \widehat{T} , $\widehat{T}^{(r)}$ and $\widehat{R}^{(p)}$, $p \leq q$, each of which is now covered. \blacksquare

In the last part of this subsection, we refine the estimate for $R\widehat{V}$. First, we recall a basic calculus estimate from [66].

Proposition 8.11 *Let $g(x) = (e^{ix} - 1)/x$. For any $k \geq 0$, there exists a constant $C > 0$ such that $|g^{(k)}(x)| \leq C$ and $|g^{(k)}(x)| \leq C/|x|$ for all $x \in \mathbb{R}$.*

Proof This is [66, Proposition 13.2]. We give the proof for completeness.

Define the analytic functions $q_k, r_k : \mathbb{C} \rightarrow \mathbb{C}$ for $k \geq 1$,

$$q_k(z) = e^z - \sum_{m=0}^{k-1} \frac{z^m}{m!}, \quad r_k(z) = \frac{q_k(z)}{z^k}.$$

By Taylor's theorem, there exists ξ between 0 and z such that

$$q_k(z) = \sum_{m=0}^{k-1} q_k^{(m)}(0) z^m / m! + q_k^{(k)}(\xi) z^k / k! = e^\xi z^k / k!,$$

so that $|q_k(ix)| \leq |x|^k / k!$. Similarly,

$$q_k(z) = \sum_{m=0}^{k-2} q_k^{(m)}(0) z^m / m! + q_k^{(k-1)}(\xi) z^{k-1} / (k-1)! = (e^\xi - 1) z^{k-1} / (k-1)!,$$

so that $|q_k(ix)| \leq |x|^{k-1} / (k-1)!$

Next, note by induction that $r_1^{(k)} \in \mathbb{R}\{e^z/z, e^z/z^2, \dots, e^z/z^k, (e^z - 1)/z^{k+1}\}$. But $e^z/z^j - r_j \in \mathbb{R}\{1/z, \dots, 1/z^j\}$. Hence there exist constants a_1, \dots, a_{k+1} and a polynomial p of degree at most k such that

$$r_1^{(k)}(z) = \sum_{j=1}^{k+1} a_j r_j(z) + p(z)/z^{k+1}.$$

Since all terms in this identity are analytic with the possible exception of the last one, we deduce that $p \equiv 0$. Hence

$$r_1^{(k)}(z) = \sum_{j=1}^{k+1} a_j r_j(z) = \sum_{j=1}^{k+1} a_j q_j(z)/z^j.$$

Since $g(x) = ir_1(ix)$, the result follows by substituting in the estimates for q_j . \blacksquare

Proposition 8.12 *Let $v \in \mathcal{F}_\theta(Y^\varphi)$. Then $|v(y, u) - v(y', u)| \leq 4C_1 \|v\|_\theta \inf_{Y_j} \varphi^\eta d_{\theta_1}(y, y')$ for all $(y, u), (y', u) \in Y^\varphi$ with $y, y' \in Y_j, j \geq 1$.*

Proof We have $|v(y) - v(y')| \leq \min\{2\|v\|_\infty, |v|_\theta \varphi(y) d_\theta(y, y')\} \leq 2\|v\|_\theta |1_{Y_j} \varphi|_\infty^\eta d_\theta(y, y')^\eta \leq 4C_1 \|v\|_\theta \inf_{Y_j} \varphi^\eta d_{\theta_1}(y, y')$. \blacksquare

It is convenient to split v into a part independent of u and a part that vanishes at $u = 0$.

Proposition 8.13 $R\widehat{V}^{(q)} : i\mathbb{R} \rightarrow \mathcal{B}(\mathcal{F}_\theta(Y^\varphi), \mathcal{F}_{\theta_1}(Y))$ is C^q for v independent of u . Moreover, there exists $C > 0$ such that

$$\|R\widehat{V}^{(q)}(ib)\|_{\theta_1} \leq C \|v\|_\theta \int_Y \varphi^{q+2\eta} d\mu |b|^{-(1-\eta)},$$

for all $b \in \mathbb{R} \setminus \{0\}$ and $v \in \mathcal{F}_\theta(Y^\varphi)$ such that v is independent of u .

Proof Write $R\widehat{V}^{(q)}(ib) = iR(f(b)v)$ where

$$f(b) = b^{-1}(e^{-ib\varphi} - 1) = \varphi g(b\varphi), \quad g(x) = x^{-1}(e^{-ix} - 1).$$

By Proposition 8.11, for any $k \in \mathbb{N}$, there exists a constant $C > 0$ such that for all $x \in \mathbb{R}$,

$$|g^{(k)}(x)| \leq C \min\{1, |x|^{-1}\} = C|x|^{-1} \min\{1, |x|\} \leq C|x|^{-(1-\eta)}. \quad (8.2)$$

Hence

$$|1_{Y_j} f^{(k)}(b)|_\infty \ll |1_{Y_j} \varphi|_\infty^{k+1} (|b| |1_{Y_j} \varphi|_\infty)^{-(1-\eta)} = |1_{Y_j} \varphi|_\infty^{k+\eta} |b|^{-(1-\eta)}.$$

By the MVT argument

$$|1_{Y_j} f^{(q)}(b)|_\infty \ll |1_{Y_j} \varphi|_\infty^{q+\eta} |b|^{-(1-\eta)} \leq (2C_1)^{q+1} \inf_{Y_j} \varphi^{q+\eta} |b|^{-(1-\eta)}.$$

Next, let $y, y' \in Y_j$. Using the identity $xg(bx) - x'g(bx') = e^{ibx'}(x - x')g(b(x - x'))$,

$$f(b)(y) - f(b)(y') = e^{ib\varphi(y')}(\varphi(y) - \varphi(y'))g(b(\varphi(y) - \varphi(y'))).$$

By (8.2),

$$\begin{aligned} |f^{(k)}(b)(y) - f^{(k)}(b)(y')| &\ll \sum_{j=0}^k \varphi(y')^{k-j} |\varphi(y) - \varphi(y')|^{1+j} |g^{(j)}(b(\varphi(y) - \varphi(y')))| \\ &\ll \sum_{j=0}^k \varphi(y')^{k-j} |\varphi(y) - \varphi(y')|^{j+\eta} |b|^{-(1-\eta)} \\ &\leq (k+1)(2C_1)^{k+1} \inf_{Y_j} \varphi^{k+\eta} d_{\theta_1}(y, y') |b|^{-(1-\eta)}. \end{aligned}$$

By the MVT argument

$$|f^{(q)}(b)(y) - f^{(q)}(b)(y')| \ll \inf_{Y_j} \varphi^{q+\eta} d_{\theta_1}(y, y') |b|^{-(1-\eta)}.$$

It follows that $\|1_{Y_j} f^{(q)}(b)\|_{\theta_1} \ll \inf_{Y_j} \varphi^{q+\eta} |b|^{-(1-\eta)}$. By Proposition 8.12, $\|1_{Y_j} f^{(q)}(b)v\|_{\theta_1} \ll \|v\|_{\theta} \inf_{Y_j} \varphi^{q+2\eta} |b|^{-(1-\eta)}$. Now apply Corollary 7.2. \blacksquare

In the remainder of this subsection, we work with the function spaces $\mathcal{F}_{\theta, \eta}(Y^\varphi)$ with norm $\|\cdot\|_{\theta, \eta} = |\cdot|_{\theta} + |\cdot|_{\infty, \eta}$ as defined in Section 3.1.2.

Proposition 8.14 *There is a constant $C > 0$ such that $\|RV(t)\|_{\theta_1} \leq C\|v\|_{\theta, \eta} t^{-q}$ for all $v \in \mathcal{F}_{\theta, \eta}(Y^\varphi)$ with $v(y, 0) \equiv 0$, and all $t > 1$.*

Proof Recall that $V(t)(y) = 1_{\{\varphi(y) > t\}} v(y, \varphi(y) - t)$. By (3.1),

$$|1_{Y_j} V(t)|_{\infty} \leq |v|_{\infty} 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} \leq |v|_{\infty} 1_{\{\inf_{Y_j} \varphi > t/(2C_1)\}}.$$

Also, for $y, y' \in Y_j$, $j \geq 1$, with $\varphi(y) \geq \varphi(y')$,

$$V(t)(y) - V(t)(y') = \begin{cases} v(y, \varphi(y) - t) - v(y', \varphi(y') - t), & \varphi(y') > t \\ v(y, \varphi(y) - t), & \varphi(y) > t \geq \varphi(y') \\ 0, & \varphi(y) \leq t \end{cases}.$$

If $\varphi(y') > t$, then using Proposition 8.12,

$$\begin{aligned} |V(t)(y) - V(t)(y')| &\leq 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} \{|v(y, \varphi(y) - t) - v(y', \varphi(y) - t)| \\ &\quad + |v(y', \varphi(y) - t) - v(y', \varphi(y') - t)|\} \\ &\leq 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} \{4C_1 \inf_{Y_j} \varphi^\eta |v|_{\theta} d_{\theta_1}(y, y') + |v|_{\infty, \eta} |\varphi(y) - \varphi(y')|^\eta\} \\ &\leq 5C_1 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} \|v\|_{\theta, \eta} \inf_{Y_j} \varphi^\eta d_{\theta_1}(y, y'). \end{aligned}$$

If $\varphi(y) > t \geq \varphi(y')$, then

$$\begin{aligned} |V(t)(y) - V(t)(y')| &= 1_{\{\varphi(y) > t \geq \varphi(y')\}} |v(y, \varphi(y) - t)| \\ &= 1_{\{\varphi(y) > t \geq \varphi(y')\}} |v(y, \varphi(y) - t) - v(y, 0)| \leq 1_{\{\varphi(y) > t \geq \varphi(y')\}} |v|_{\infty, \eta} |\varphi(y) - t|^\eta \\ &\leq 1_{\{\varphi(y) > t\}} |v|_{\infty, \eta} |\varphi(y) - \varphi(y')|^\eta \leq C_1 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} |v|_{\infty, \eta} \inf_{Y_j} \varphi^\eta d_{\theta_1}(y, y'). \end{aligned}$$

Hence in all cases,

$$|V(t)(y) - V(t)(y')| \leq 5C_1 1_{\{|1_{Y_j} \varphi|_{\infty} > t\}} \|v\|_{\theta, \eta} \inf_{Y_j} \varphi^\eta d_{\theta_1}(y, y').$$

By (3.1),

$$\|1_{Y_j} V(t)\|_{\theta_1} \leq 5C_1 \|v\|_{\theta, \eta} 1_{\{\inf_{Y_j} \varphi > t/(2C_1)\}} \inf_{Y_j} \varphi^\eta.$$

Hence by Corollary 7.2,

$$\begin{aligned} \|RV(t)\|_{\theta_1} &\leq 10C_1 C_2 \|v\|_{\theta, \eta} \sum \mu(Y_j) 1_{\{\inf_{Y_j} \varphi > t/(2C_1)\}} \inf_{Y_j} \varphi^\eta \\ &\leq 10C_1 C_2 \|v\|_{\theta, \eta} \int_Y 1_{\{\varphi > t/(2C_1)\}} \varphi^\eta d\mu. \end{aligned}$$

Now apply Proposition 8.5. \blacksquare

Corollary 8.15 *Let $\kappa : \mathbb{R} \rightarrow \mathbb{R}$ be \mathbb{C}^∞ with $|\kappa^{(k)}(b)| = O((b^2 + 1)^{-1})$ for all $k \in \mathbb{N}$. Then $\kappa R\widehat{V} \in \mathcal{R}(\|v\|_{\theta, \eta} t^{-q})$ in $\mathcal{F}_{\theta_1}(Y)$ for all $v \in \mathcal{F}_{\theta, \eta}(Y^\varphi)$.*

Proof Write $v(y, u) = v_0(y) + v_1(y, u)$ where $v_0(y) = v(y, 0)$. The result follows by combining Propositions 8.13 and 8.14 and applying Proposition 8.2. \blacksquare

8.3 Further estimates for dealing with the singularity at zero

By Propositions 7.4 and 8.7, there exists $\delta > 0$ such that $\widehat{R}(ib)$ has a C^q family of simple eigenvalues $\lambda(b)$, $|b| < \delta$, with $\lambda(0) = 1$ and $\lambda'(0) = -i|\varphi|_1$. Let $P(b)$, $|b| < \delta$, denote the corresponding C^q family of spectral projections, with $P(0)v = \int_Y v d\mu$ for $v \in L^1(Y)$.

Proposition 8.16 $b^{-1}P(0)R\widehat{V}(ib) \in \mathcal{R}(\|v\|_\infty t^{-(\beta-1)})$ for all $v \in L^\infty(Y^\varphi)$ with $\int_{Y^\varphi} v d\mu^\varphi = 0$.

Proof Let $\hat{g}(s) = s^{-1}P(0)R\widehat{V}(s) = s^{-1} \int_Y \int_0^{\varphi(y)} e^{-s(\varphi(y)-u)} v(y, u) du d\mu$. Since v has mean zero,

$$\hat{g}(s) = \int_Y \int_0^{\varphi(y)} s^{-1} (e^{-s(\varphi(y)-u)} - 1) v(y, u) du d\mu.$$

Hence

$$g(t) = - \int_Y \int_0^{\varphi(y)} 1_{\{\varphi(y) > t+u\}} v(y, u) du d\mu.$$

By Proposition 8.5,

$$|g(t)| \leq \|v\|_\infty \int_Y \int_0^\varphi 1_{\{\varphi > t\}} du d\mu = \|v\|_\infty \int_Y \varphi 1_{\{\varphi > t\}} d\mu \ll \|v\|_\infty t^{-(\beta-1)},$$

as required. \blacksquare

Define $\widetilde{R}(s) = s^{-1}(\widehat{R}(s) - \widehat{R}(0))$, $s \in \mathbb{H}$.

Proposition 8.17 *Let $q_1 > 0$. There exists a constant $C > 0$ such that*

$$\|\widetilde{R}^{(q_1)}(ib)\|_{\theta_1} \leq \begin{cases} C|b|^{-(1-\eta)} & q_1 < \beta - 2\eta \\ C & q_1 < \beta - 1 \end{cases} \quad \text{for all } b \in \mathbb{R} \setminus \{0\}.$$

Proof Write $\widetilde{R}(ib)v = R(f(b)v)$ where $f(b) = \int_0^\varphi e^{-ibt} dt$. This is the same function as in the proof of Proposition 8.13, leading to the same conclusion for $q_1 < \beta - 2\eta$.

The argument for $q_1 < \beta - 1$ is much simpler since $f^{(k)}(b) = \int_0^\varphi e^{-ibt} (-it)^k dt$ and φ^{q_1+1} is integrable. We omit the details. \blacksquare

For $b \in (-\delta, \delta)$, define

$$\widetilde{P}(b) = b^{-1}(P(b) - P(0)), \quad \widetilde{\lambda}(b) = (ib|\varphi|_1)^{-1}(1 - \lambda(b)).$$

Proposition 8.18 *The conclusion of Proposition 8.17 holds with $\|\widetilde{R}^{(q_1)}(ib)\|_{\theta_1}$ replaced by $\|\widetilde{P}^{(q_1)}(b)\|_{\theta_1}$ or $|\widetilde{\lambda}^{(q_1)}(b)|$ for all $|b| < \delta$.*

Proof Let $\Gamma \subset \mathbb{C}$ be a sufficiently small circle centered at 1. Then $P(b) = (2\pi i)^{-1} \int_{\Gamma} (\xi - \widehat{R}(ib))^{-1} db$ for all $|b| < \delta$. Hence

$$\widetilde{P}(b) = (2\pi i)^{-1} \int_{\Gamma} (\xi - \widehat{R}(ib))^{-1} \widetilde{R}(ib) (\xi - \widehat{R}(0))^{-1} d\xi.$$

By Proposition 8.7, $(\xi - \widehat{R})^{-1}$ is C^{q_1} uniformly in $\xi \in \Gamma$. Hence the estimates for $\widetilde{P}^{(q_1)}$ follow from Proposition 8.17.

Next, write

$$\widetilde{\lambda}(b)P(b) = \{-\widetilde{R}(ib)P(b) + (I - \widehat{R}(0))\widetilde{P}(b)\}/(i|\varphi|_1). \quad (8.3)$$

Since P is C^{q_1} , it follows from Proposition 8.17 and the estimates for $\widetilde{P}^{(q_1)}$ that $\|(\widetilde{\lambda}P)^{(q_1)}(b)\|_{\theta_1}$ also satisfies these estimates. Let $f(b) \in \mathcal{F}_{\theta_1}(Y)$ denote the eigenfunction corresponding to $\lambda(b)$ normalised so that $\int_Y f(b) d\mu = 1$. (If necessary, shrink δ so that $f(b) > 0$ ensuring that the normalisation exists.) Then $b \mapsto f(b)$ is C^{q_1} and

$$\widetilde{\lambda}(b) = \int_Y \widetilde{\lambda}(b) f(b) d\mu = \int_Y \widetilde{\lambda}(b) P(b) \lambda(b)^{-1} f(b) d\mu,$$

yielding the estimates for $\widetilde{\lambda}^{(q_1)}$. ■

8.4 Truncation

Given $N \geq 1$, we replace φ by $\varphi \wedge N = \min\{\varphi, N\}$. Consider the suspension semiflows F_t and $F_{N,t}$ on Y^φ and $Y^{\varphi \wedge N}$ respectively. Let $\rho_{v,w}$ and $\rho_{v,w}^{\text{trunc}}$ denote the respective correlation functions. In particular, $\rho_{v,w}^{\text{trunc}}(t) = \int_{Y^{\varphi \wedge N}} v w \circ F_{N,t} d\mu^{\varphi \wedge N} - \int_{Y^{\varphi \wedge N}} v d\mu^{\varphi \wedge N} \int_{Y^{\varphi \wedge N}} w d\mu^{\varphi \wedge N}$ where the observables $v, w : Y^{\varphi \wedge N} \rightarrow \mathbb{R}$ are the restrictions of $v, w : Y^\varphi \rightarrow \mathbb{R}$ to $Y^{\varphi \wedge N}$.

Proposition 8.19 *There are constants $C, t_0 > 0, N_0 \geq 1$ such that*

$$|\rho_{v,w}(t) - \rho_{v,w}^{\text{trunc}}(t)| \leq C|v|_\infty |w|_\infty (tN^{-\beta} + N^{-(\beta-1)}),$$

for all $v, w \in L^\infty(Y^\varphi)$, $N \geq N_0, t > t_0$. ■

Proof This proof follows [64, Section 3] and [65, Appendix A]. We begin with some definitions and preliminary estimates.

Let $F_{N,t} : Y^{\varphi \wedge N} \rightarrow Y^{\varphi \wedge N}$ denote the truncated flow. Define the top of the tower $T_N = Y^\varphi \setminus Y^{\varphi \wedge N}$ and the borderline region $\partial T_N = \{(y, N) \in Y^\varphi : \varphi(y) > N\}$. Also, we define the thickened borderline region

$$\partial T'_N = \{(y, u) \in Y^\varphi : \varphi(y) > N, N-1 \leq u \leq N\}.$$

Now, $\mu^\varphi(\partial T'_N) = |\varphi|_1^{-1} \int_Y 1_{\{\varphi > N\}} d\mu \ll N^{-\beta}$ and by Proposition 8.5,

$$\mu^\varphi(T_N) = |\varphi|_1^{-1} \int_Y 1_{\{\varphi > N\}} (\varphi - N) d\mu \leq |\varphi|_1^{-1} \int_Y 1_{\{\varphi > N\}} \varphi d\mu \ll N^{-(\beta-1)}.$$

We note the related estimate

$$|\varphi|_1 - |\varphi \wedge N|_1 = \int_Y (\varphi - (\varphi \wedge N)) d\mu = \int_Y 1_{\{\varphi > N\}} (\varphi - N) d\mu \ll N^{-(\beta-1)}.$$

Choosing N sufficiently large, we can suppose that $|\varphi \wedge N|_1 \geq \frac{1}{2}|\varphi|_1$ and hence that $||\varphi|_1^{-1} - |\varphi \wedge N|_1^{-1}| \ll N^{-(\beta-1)}$.

Fix $t \geq 1$. Note that if $F_s(y, u) \in T_N$ for some $s \in [0, t]$, then either $(y, u) \in T_N$ or $F_s(y, u) \in \partial T_N$ for some $s \in [0, t]$. Hence

$$\begin{aligned} \{(y, u) \in Y^{\varphi \wedge N} : F_t(y, u) \neq F_{N,t}(y, u)\} &\subset \{(y, u) \in Y^{\varphi \wedge N} : F_s(y, u) \in T_N \text{ for some } s \in [0, t]\} \\ &\subset T_N \cup \bigcup_{s \in [0, t+1]} F_s^{-1} \partial T_N = T_N \cup \bigcup_{j=0, \dots, [t+1]} F_j^{-1} \partial T'_N. \end{aligned}$$

It follows that

$$\begin{aligned} &\mu^\varphi \{(y, u) \in Y^{\varphi \wedge N} : F_t(y, u) \neq F_{N,t}(y, u)\} \\ &\leq \mu^\varphi(T_N) + \sum_{j=0}^{[t+1]} \mu^\varphi(F_j^{-1} \partial T'_N) = \mu^\varphi(T_N) + [t+1] \mu^\varphi(\partial T'_N) \ll N^{-(\beta-1)} + tN^{-\beta}. \end{aligned}$$

Now,

$$\int_{Y^\varphi} v w \circ F_t d\mu^\varphi - \int_{Y^{\varphi \wedge N}} v w \circ F_{N,t} d\mu^{\varphi \wedge N} = I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \int_{T_N} v w \circ F_t d\mu^\varphi, \quad I_2 = \int_{Y^{\varphi \wedge N}} v (w \circ F_t - w \circ F_{N,t}) d\mu^\varphi, \\ I_3 &= (|\varphi|_1^{-1} - |\varphi \wedge N|^{-1}) \int_Y \int_0^{\varphi \wedge N} v w \circ F_{N,t} du d\mu^{\varphi \wedge N}. \end{aligned}$$

It follows readily from the preceding calculations that

$$\begin{aligned} |I_1| &\leq |v|_\infty |w|_\infty \mu^\varphi(T_N) \ll |v|_\infty |w|_\infty N^{-(\beta-1)}, \quad |I_3| \ll |v|_\infty |w|_\infty N^{-(\beta-1)}, \\ |I_2| &\leq 2|v|_\infty |w|_\infty \mu^\varphi \{F_t(y, u) \neq F_{N,t}(y, u)\} \ll |v|_\infty |w|_\infty (N^{-(\beta-1)} + tN^{-\beta}). \end{aligned}$$

Hence $|\int_{Y^\varphi} v w \circ F_t d\mu^\varphi - \int_{Y^{\varphi \wedge N}} v w \circ F_{N,t} d\mu^{\varphi \wedge N}| \ll |v|_\infty |w|_\infty (N^{-(\beta-1)} + tN^{-\beta})$.

A simpler calculation shows that $|\int_{Y^\varphi} v d\mu^\varphi - \int_{Y^{\varphi \wedge N}} v d\mu^{\varphi \wedge N}| \ll |v|_\infty N^{-(\beta-1)}$ and hence that $|\int_{Y^\varphi} v d\mu^\varphi \int_{Y^\varphi} w d\mu^\varphi - \int_{Y^{\varphi \wedge N}} v d\mu^{\varphi \wedge N} \int_{Y^{\varphi \wedge N}} w d\mu^{\varphi \wedge N}| \ll |v|_\infty |w|_\infty N^{-(\beta-1)}$. This completes the proof. \blacksquare

Below we prove:

Lemma 8.20 *Assume absence of approximate eigenfunctions. There are constants $C > 0$, $m \geq 1$, $N_1 \geq 1$ such that*

$$|\rho_{v,w}^{\text{trunc}}(t)| \leq C \|v\|_{\theta, \eta} |w|_{\infty, m} t^{-(\beta-1)},$$

for all $v \in \mathcal{F}_{\theta, \eta}(Y^{\varphi \wedge N})$, $w \in L^{\infty, m}(Y^{\varphi \wedge N})$, $N \geq N_1$, $t > 1$.

Proof of Theorem 3.6 In general, $w \in L^{\infty,m}(Y^\varphi)$ need not restrict to $w \in L^{\infty,m}(Y^{\varphi \wedge N})$, but we can choose $w_N \in L^{\infty,m}(Y^{\varphi \wedge N})$ so that $|w_N|_{\infty,m} \ll |w|_{\infty,m}$ and $w_N \equiv w$ outside the set $S_N = \{(y, u) \in Y^{\varphi \wedge N} : \varphi(y) > N, u \in (N-1, N]\}$. Then

$$\begin{aligned} |\rho_{v,w}^{\text{trunc}}(t) - \rho_{v,w_N}^{\text{trunc}}(t)| &\leq |v|_\infty(|w|_\infty + |w_N|_\infty) \mu^{\varphi \wedge N}(F_{N,t}^{-1} S_N) \\ &= |v|_\infty(|w|_\infty + |w_N|_\infty) \mu^{\varphi \wedge N}(S_N) \ll |v|_\infty |w|_\infty \mu(\varphi > N) \ll |v|_\infty |w|_\infty N^{-\beta}. \end{aligned}$$

Taking $N = [t]$, the result follows directly from Proposition 8.19 and Lemma 8.20. \blacksquare

In the remainder of this subsection, we outline the strategy for proving Lemma 8.20.

By assumption, we can fix a finite union Z of partition elements such that the corresponding finite subsystem Z_0 does not support approximate eigenfunctions. Choose $N_1 \geq |1_Z \varphi|_\infty$.

For each fixed N , the truncated roof function $\varphi \wedge N$ is bounded and hence the results in Section 7 apply. In particular, $\hat{\rho}_{v,w}^{\text{trunc}}$ is C^∞ on $\bar{\mathbb{H}}$ and contours of integration can be moved to the imaginary axis. From now on we suppress the superscript “trunc” for sake of readability. The calculations in Proposition 6.3 and Corollary 6.4 now proceed on the imaginary axis in identical fashion to the calculation on \mathbb{H} . Hence

$$\rho_{v,w}(t) = \int_{-\infty}^{\infty} e^{ibt} \hat{\rho}_{v,w}(ib) db, \quad \hat{\rho}_{v,w} = \hat{J}_{0,v,w} + |\varphi|_1^{-1} \int_Y \hat{T} R \hat{V} \hat{w} d\mu,$$

where the constituent parts $\hat{J}_{0,v,w}$, $\hat{T} = (I - \hat{R})^{-1}$, \hat{V} , \hat{w} are C^∞ “truncated” versions of the originals.

It follows from rapid mixing for the truncated semiflow that $t \mapsto (\rho_{v,w})^{(m)}(t)$ lies in $L^1(\mathbb{R})$ for all $m \geq 0$, so we can use integration by parts to show that

$$\hat{\rho}_{v,w}(ib) = (ib)^{-m} \hat{\rho}_{v,\partial_t^m w}(ib) \quad \text{for all } b \neq 0, m \geq 0.$$

Choose $\psi : \mathbb{R} \rightarrow [0, 1]$ to be C^∞ and compactly supported such that $\psi \equiv 1$ on a neighborhood of zero. Let $\kappa_m(b) = (1 - \psi(b))(ib)^{-m}$. Note that

$$\psi, \kappa_m \in \mathcal{R}(t^{-p}) \quad \text{for all } p > 0, m \geq 2. \quad (8.4)$$

We have

$$\begin{aligned} \hat{\rho}_{v,w}(ib) &= \psi(b) \hat{\rho}_{v,w}(ib) + \kappa_m(b) \hat{\rho}_{v,\partial_t^m w}(ib) \\ &= \psi(b) \hat{J}_{0,v,w}(ib) + |\varphi|_1^{-1} \psi(b) \int_Y \hat{T}(ib) R \hat{V}(ib) \hat{w}(ib) d\mu \\ &\quad + \kappa_m(b) \hat{J}_{0,v,\partial_t^m w}(ib) + |\varphi|_1^{-1} \kappa_m(b) \int_Y \hat{T}(ib) R \hat{V}(ib) \widehat{\partial_t^m w}(ib) d\mu. \end{aligned} \quad (8.5)$$

It remains to estimate the inverse Fourier transform of each term in (8.5).

Proposition 8.21 *After truncation, uniformly in $N \geq 1$,*

$$\psi \hat{J}_{0,v,w} \in \mathcal{R}(|v|_\infty |w|_\infty t^{-(\beta-1)}), \quad \kappa_m \hat{J}_{0,v,w} \in \mathcal{R}(|v|_\infty |w|_\infty t^{-(\beta-1)}),$$

for all $m \geq 2$, $v \in L^\infty(Y^\varphi)$, $w \in L^\infty(Y^\varphi)$.

Proof By Corollary 8.6, $|\widehat{J}_{0,v,w}| \in \mathcal{R}(|v|_\infty |w|_\infty t^{-(\beta-1)})$ uniformly in N . Using (8.4) and Proposition 8.4, $\psi \widehat{J}_{0,v,w} \in \mathcal{R}(t^{-\beta} \star (|v|_\infty |w|_\infty t^{-(\beta-1)}) = \mathcal{R}(|v|_\infty |w|_\infty t^{-(\beta-1)})$. Similarly, for $\kappa_m \widehat{J}_{0,v,w}$. \blacksquare

The main two estimates are as follows. Recall that $q \in (\beta - 1, \beta)$.

Lemma 8.22 *There exists $N_1 \geq 1$ such that after truncation, uniformly in $N \geq N_1$, there exist $m \geq 2$ and ψ such that*

$$\psi \int_Y \widehat{T} R \widehat{V} \widehat{w} d\mu \in \mathcal{R}(\|v\|_{\theta,\eta} |w|_\infty t^{-(\beta-1)}),$$

for all $v \in \mathcal{F}_{\theta,\eta}(Y^\varphi)$ with $\int_{Y^\varphi} v d\mu^\varphi = 0$, $w \in L^\infty(Y^\varphi)$.

Lemma 8.23 *There exists $N_1 \geq 1$ such that after truncation, uniformly in $N \geq N_1$, there exist $m \geq 2$ such that*

$$\kappa_m \int_Y \widehat{T} R \widehat{V} \widehat{w} d\mu \in \mathcal{R}(\|v\|_{\theta,\eta} |w|_\infty t^{-q}),$$

for all $v \in \mathcal{F}_{\theta,\eta}(Y^\varphi)$, $w \in L^\infty(Y^\varphi)$.

Proof of Lemma 8.20 Substituting the results from Proposition 8.21 and Lemmas 8.22 and 8.23 into (8.5), we obtain that $\widehat{\rho}_{v,w} \in \mathcal{R}(\|v\|_{\theta,\eta} |w|_{\infty,m} t^{-(\beta-1)})$ uniformly in $N \geq N_1$ for all $v \in \mathcal{F}_{\theta,\eta}(Y^\varphi)$ with $\int_{Y^\varphi} v d\mu^\varphi = 0$, $w \in L^{\infty,m}(Y^\varphi)$. \blacksquare

The proofs of Lemmas 8.22 and 8.23 are presented in Subsections 8.5 and 8.6 respectively.

8.5 Small b : Proof of Lemma 8.22

Let $\delta > 0$ be as in Subsection 8.3. Temporarily, we introduce the notation $\widehat{R}_N(s)v = R(e^{-s(\varphi \wedge N)}v)$.

Proposition 8.24 $\lim_{N \rightarrow \infty} \|\widehat{R}_N^{(q)}(ib) - \widehat{R}^{(q)}(ib)\|_{\theta_1} = 0$ uniformly for $|b| < \delta$.

Proof Write $\widehat{R}(ib)v - \widehat{R}_N(ib)v = R(f(b)v)$ where $f(b) = g_1(b) - g_2(b)$, $g_1(b) = e^{-ib\varphi}$, $g_2(b) = e^{-ib(\varphi \wedge N)}$. Then $\widehat{R}^{(k)}(ib)v - \widehat{R}_N^{(k)}(ib)v = R(f^{(k)}(b)v)$ for all $k \in \mathbb{N}$ where

$$f^{(k)}(b) = g_1^{(k)}(b) - g_2^{(k)}(b), \quad g_1^{(k)}(b) = (-i)^k e^{-ib\varphi} \varphi^k, \quad g_2^{(k)}(b) = (-i)^k e^{-ib(\varphi \wedge N)} (\varphi \wedge N)^k.$$

If $\sup_{Y_j} \varphi \leq N$, then $f^{(k)}(b) \equiv 0$ on Y_j .

If $\sup_{Y_j} \varphi > N$, then $|1_{Y_j} g_i^{(k)}(b)|_\infty \leq \sup_{Y_j} \varphi^k \leq 2C_1 \inf_{Y_j} \varphi^k$ for $i = 1, 2$. By the MVT argument $|1_{Y_j} f^{(q)}(b)|_\infty \leq 4C_1 \inf_{Y_j} \varphi^q$.

Next, for $y_1, y_2 \in Y_j$, we have $g_1(b)(y_1) - g_1(b)(y_2) = I_1 + I_2$ where

$$I_1 = \{e^{-ib\varphi(y_1)} - e^{-ib\varphi(y_2)}\} \varphi(y_1)^k, \quad I_2 = e^{-ib\varphi(y_2)} \{\varphi(y_1)^k - \varphi(y_2)^k\}.$$

Note that

$$\begin{aligned} |I_1| &\leq 2|b|^\eta |\varphi(y_1) - \varphi(y_2)|^\eta \varphi(y_1)^k \leq 2|b|^\eta |1_{Y_j} \varphi|_\theta^\eta \sup_{Y_j} \varphi^k d_\theta(y_1, y_2)^\eta \ll \inf_{Y_j} \varphi^{k+\eta} d_{\theta_1}(y, y'), \\ |I_2| &\leq k|\varphi(y_1) - \varphi(y_2)| \sup_{Y_j} \varphi^{k-1} \leq k|1_{Y_j} \varphi|_\theta \sup_{Y_j} \varphi^{k-1} d_\theta(y_1, y_2) \ll k \inf_{Y_j} \varphi d_{\theta_1}(y, y'), \end{aligned}$$

so $|1_{Y_j} g_1^{(k)}(b)|_{\theta_1} \ll k \inf_{Y_j} \varphi^{k+\eta}$. By the MVT argument, $|1_{Y_j} g_1^{(q)}(b)|_{\theta_1} \ll \inf_{Y_j} \varphi^{q+\eta}$. Similarly $|1_{Y_j} g_2^{(q)}(b)|_{\theta_1} \ll \inf_{Y_j} \varphi^{q+\eta}$. Hence $\|1_{Y_j} f^{(k)}(b)\|_{\theta_1} \ll 1_{\{\sup_{Y_j} \varphi > N\}} \inf_{Y_j} \varphi^{k+\eta} \leq 1_{\{\inf_{Y_j} \varphi > N/(2C_1)\}} \inf_{Y_j} \varphi^{k+\eta}$.

By Remark 7.3,

$$\begin{aligned} \|\widehat{R}^{(q)}(ib) - \widehat{R}_N^{(q)}(ib)\|_{\theta_1} &\leq 2C_2 \sum \mu(Y_j) \|1_{Y_j} f^{(q)}(b)\|_{\theta_1} \\ &\ll \sum 1_{\{\inf_{Y_j} \varphi > N/(2C_1)\}} \mu(Y_j) \inf_{Y_j} \varphi^{q+\eta} \leq \int_Y 1_{\{\varphi > N/(2C_1)\}} \varphi^{q+\eta} d\mu. \end{aligned}$$

The result follows since $\varphi \in L^{q+\eta}(Y)$. ■

By Proposition 8.24, we can fix $\delta > 0$ and $N_1 \geq 1$ such that for all $N \geq N_1$ there exists a C^q family of simple eigenvalues $\lambda_N(b)$, $|b| < \delta$, for $\widehat{R}_N(ib)$ with $\lambda_N(0) = 1$ and $\lambda'_N(0) = 1$. There is also a corresponding C^q family of spectral projections $P_N(b)$ with $P_N(0)v = \int_Y v d\mu$.

From now on, we write \widehat{R} , P and λ instead of \widehat{R}_N , P_N and λ_N . Recall that $\widehat{T} = (I - \widehat{R})^{-1}$. Choose the C^∞ function $\psi : \mathbb{R} \rightarrow [0, 1]$ so that $\text{supp } \psi \in (-\delta, \delta)$, and write

$$\widehat{T}(ib)R\widehat{V}(ib) = (1 - \lambda(b))^{-1}P(b)R\widehat{V}(ib) + \widehat{T}(ib)(I - P(b))R\widehat{V}(ib). \quad (8.6)$$

We begin by dealing with the second term in (8.6).

Lemma 8.25 $\psi \int_Y \widehat{T}(I - P)R\widehat{V} \widehat{w} d\mu \in \mathcal{R}(\|v\|_{\theta, \eta} |w|_\infty t^{-q})$ for all $v \in \mathcal{F}_{\theta, \eta}(Y^\varphi)$, $w \in L^\infty(Y^\varphi)$.

Proof By definition of P and Proposition 8.7, $\widehat{T}(ib)(I - P(b))$ is C^q on $\mathcal{F}_{\theta_1}(Y)$ uniformly in N . By Proposition 8.2, $\psi \widehat{T}(I - P) \in \mathcal{R}(t^{-q})$.

Choose ψ_1 to be C^∞ with compact support such that $\psi_1 \equiv 1$ on $\text{supp } \psi$. By Corollary 8.15,

$$\psi \widehat{T}(I - P)R\widehat{V} = \{\psi \widehat{T}(I - P)\} \{\psi_1 R\widehat{V}\} \in \mathcal{R}(t^{-q} \star \|v\|_{\theta, \eta} t^{-q}),$$

in $\mathcal{F}_\theta(Y)$ and hence in $L^\infty(Y)$. Since $q > 1$, it follows from Proposition 8.4 that $\psi \widehat{T}(I - P)R\widehat{V} \in \mathcal{R}(\|v\|_{\theta, \eta} t^{-q})$ in $L^\infty(Y)$.

By Corollary 8.6, $\widehat{w} \in \mathcal{R}(|w|_\infty t^{-\beta})$ in $L^1(Y)$, so the result follows from Proposition 8.4. ■

Still working with the truncated roof function $\varphi \wedge N$, define

$$\widetilde{R}(s) = s^{-1}(\widehat{R}(s) - \widehat{R}(0)), \quad \widetilde{P}(b) = b^{-1}(P(b) - P(0)), \quad \widetilde{\lambda}(b) = (ib|\varphi \wedge N|_1)^{-1}(1 - \lambda(b)).$$

The estimates in Propositions 8.17 and 8.18 remain valid uniformly in N .

Proposition 8.26 $\psi \widetilde{\lambda}^{-1} \in \mathcal{R}(t^{-q})$.

Proof By Proposition 8.2, it suffices to show that $|(\widetilde{\lambda}^{-1})^{(q)}(b)| \ll |b|^{-(1-\eta)}$.

By Proposition 8.18, $|\widetilde{\lambda}^{(p)}(b)| \ll |b|^{-(1-\eta)}$ for $p < \beta - 2\eta$ and $|\widetilde{\lambda}^{(p)}(b)| \ll 1$ for $p < \beta - 1$. In particular, $\widetilde{\lambda}$ is Hölder uniformly in N and b . Recall that $\widetilde{\lambda}(0) = 1$. Shrinking δ , we obtain $|\widetilde{\lambda}(b)| \geq \frac{1}{2}$ for all $|b| < \delta$, $N \geq N_1$.

The case $\beta \geq 2$. Note that $(\tilde{\lambda}^{-1})^{(q)}$ is a finite linear combination of finite products with factors $\tilde{\lambda}^{-1}$ (each bounded by 2) and $\tilde{\lambda}^{(q_1)} \dots \tilde{\lambda}^{(q_k)}$ where $q_1 + \dots + q_k = q$. If $q_{i_j} \geq \beta - 1$ for distinct values of i_j , then $\beta \leq 2\beta - 2 \leq q_{i_1} + q_{i_2} \leq q$ contradicting the assumption that $q < \beta$. Hence there exists at most one i such that $q_i \geq \beta - 1$ and $|\tilde{\lambda}^{(q_1)} \dots \tilde{\lambda}^{(q_k)}| \ll |\tilde{\lambda}^{(q_i)}| \ll |b|^{-(1-\eta)}$. It follows that $|(\tilde{\lambda}^{-1})^{(q)}(b)| \ll |b|^{-(1-\eta)}$.

The case $\beta \in (1, 2)$. Let $q = 1 + r < \beta$. Then $(\tilde{\lambda}^{-1})' = -\tilde{\lambda}^{-2}\tilde{\lambda}'$ and $|(\tilde{\lambda}^{-1})'(b) - (\tilde{\lambda}^{-1})'(b')| \ll |\tilde{\lambda}'(b) - \tilde{\lambda}'(b')| \ll |b|^{-(1-\eta)}|b - b'|^r$ for $|b| \leq |b'|$. Hence $|(\tilde{\lambda}^{-1})^{(q)}(b)| \ll |b|^{-(1-\eta)}$. ■

Proof of Lemma 8.22 By Lemma 8.25, it remains to handle the first term in (8.6), namely

$$\begin{aligned} (1 - \lambda(b))^{-1}P(b)R\widehat{V}(ib) &= (ib|\varphi \wedge N|)^{-1}\tilde{\lambda}(b)^{-1}P(b)R\widehat{V}(ib) \\ &= (i|\varphi \wedge N|)^{-1}\tilde{\lambda}(b)^{-1}b^{-1}P(0)R\widehat{V}(ib) + (i|\varphi \wedge N|)^{-1}\tilde{\lambda}(b)^{-1}\tilde{P}(b)R\widehat{V}(ib). \end{aligned}$$

Choose $q \in (1, \beta)$, $q \geq \beta - 1$. By Propositions 8.2, 8.18 and 8.26, $\psi\tilde{P}$, $\psi\tilde{\lambda}^{-1} \in \mathcal{R}(t^{-q})$. By Proposition 8.16,

$$\psi(b)\tilde{\lambda}(b)^{-1}b^{-1}P(0)R\widehat{V}(ib) \in \mathcal{R}(t^{-q} \star |v|_{\infty} t^{-(\beta-1)}).$$

Choose ψ_1 to be C^∞ with compact support such that $\psi_1 \equiv 1$ on $\text{supp } \psi$. Using also Corollary 8.15,

$$\psi\tilde{\lambda}^{-1}\tilde{P}R\widehat{V} = \{\psi\tilde{\lambda}^{-1}\}\{\psi_1\tilde{P}\}\{\psi_1R\widehat{V}\} \in \mathcal{R}(t^{-q} \star t^{-q} \star \|v\|_{\theta, \eta} t^{-q}) \quad \text{in } L^\infty(Y).$$

Hence by Proposition 8.4, $\psi(1 - \lambda)^{-1}PR\widehat{V} \in \mathcal{R}(\|v\|_{\theta, \eta} t^{-(\beta-1)})$ in $L^\infty(Y)$. Combining with Corollary 8.6, we obtain $\psi \int_Y (1 - \lambda)^{-1}PR\widehat{V} \widehat{w} d\mu \in \mathcal{R}(\|v\|_{\theta, \eta} |w|_{\infty} t^{-(\beta-1)})$ as required. ■

8.6 Large b : Proof of Lemma 8.23

Define $\kappa_m(b) = (1 - \psi(b))(ib)^{-m}$ where ψ is chosen as in Lemma 8.22.

Proposition 8.27 *There exists $C > 0$ and $m \geq 2$ such that*

$$\kappa_m \widehat{T} \in \mathcal{R}(t^{-q}) \text{ in } \mathcal{B}(\mathcal{F}_{\theta_1}(Y)) \text{ for all } N \geq N_1, t > 1.$$

Proof By the choice of $N \geq N_1$, absence of approximate eigenfunctions passes over to the truncated semiflow. By Corollary 8.10, $\|\widehat{T}^q(ib)\|_{\theta_1} \leq C|b|^\alpha$ for $b \in \text{supp } \kappa_m$. Note that all constants entering into C and α are universal (C_2, C_3 , etc) or depend only on the values of φ on the finite subsystem Z_0 . In particular, C and α are independent of N for all $N \geq N_1$.

Hence $\|(\kappa_m \widehat{T})^{(q)}(b)\|_{\theta_1} \ll (|b| + 1)^{\alpha - m}$. Choosing $m \geq \alpha + 2$ ensures integrability and the result follows from Proposition 8.2. ■

Proof of Lemma 8.23 Write $\kappa_m = \kappa_{m-2}\kappa_2$, where κ_i is C^∞ , vanishes in a neighborhood of zero, and is $O(|b|^{-i})$, for $i = 2$ and $i = m - 2$.

By Corollaries 8.6 and 8.15, and Proposition 8.27, we obtain (for sufficiently large m)

$$\kappa_m \int_Y \widehat{T}R\widehat{V} \widehat{w} d\mu = \int_Y (\kappa_{m-2}\widehat{T})(\kappa_2R\widehat{V}) \widehat{w} d\mu \in \mathcal{R}(t^{-q} \star \|v\|_{\theta, \eta} t^{-q} \star |w|_{\infty} t^{-\beta}).$$

Since $q \in (1, \beta)$, the result follows from Proposition 8.4. ■

9 Some open questions

We conclude this review article with some open questions.

- Improve the Diophantine criterion for absence of approximate eigenfunctions by reducing the number of periods from three to two in Proposition 5.3. It would suffice to show that we can take $\psi_k = 1$ in (3.2) as is the case for uniformly hyperbolic systems [48].
(In particular, the proof of rapid mixing for two falling balls in [18] seems to rely on such an improvement. Alternatively, one could try to verify the good asymptotics condition [50] described in Section 5.2; this would also lead to a stronger conclusion (robust rapid mixing rather than almost sure) in [18].)
- For flows with polynomial decay of correlations, the decay rates in this article are optimal but the class of observables is not. An open problem is to remove the requirement that observables are smooth in the flow direction. In general, this is a currently intractable problem even in the superpolynomial case. However in situations where there is additional structure in which exponential decay methods have proven successful (smooth stable foliation or contact structure) there is the possibility of combining these methods with the truncation method in Section 8.4. A key example is the infinite horizon planar periodic Lorentz gas where the optimal decay rate $O(t^{-1})$ is obtained in [14] but for a restricted class of observables.
- The statistical properties for time-one maps of rapid mixing flows in Section 1.2 are restricted to observables that are sufficiently regular in the flow direction, and this is currently the best available result even when the flow is exponentially mixing. To fix ideas, consider the finite horizon planar periodic Lorentz gas. The time-one map is exponentially mixing for Hölder observables by [11], but currently this does not lead to any statistical properties. On the other hand, the proof of superpolynomial decay for sufficiently regular observables does lead to the statistical properties listed in Subsection 1.2. Hence a natural question is to investigate how to use the result or method in [11] to extract statistical properties. This is currently the topic of work in progress with Mark Demers and Matthew Nicol.
- As mentioned in Remark 5.7, Theorem 5.6 gives a simple criterion for absence of approximate eigenfunctions when $Z_0 \subset Y$ is connected, namely that the temporal distance function D is not identically zero. Such a nonintegrability property is not immediately of use in general: Z_0 is a Cantor set of positive Hausdorff dimension and Y is often connected, but D is only Hölder. On the other hand, Z_0 can be constructed using any finite subcollection of the partition elements Y_j , and so in some sense exhausts Y . The question is whether D has sufficient structure beyond being Hölder (which on its own is clearly insufficient) to imply that the lower box dimension of $D(Z_0 \times Z_0)$ is close to that of $D(Y \times Y)$ for suitable chosen Z_0 . This would rule out approximate eigenfunctions when Y is connected and D is not identically zero.
- Methods for suspension semiflows and flows can be adapted to toral extensions of maps, replacing roof functions $\varphi : Y \rightarrow \mathbb{R}^+$ by cocycles $\varphi : Y \rightarrow \mathbb{R}^d$. Rapid mixing for toral extensions (and general compact group extensions) of uniformly expanding/hyperbolic

maps is analysed in depth in [49]. Results on rapid and slow mixing for toral extensions of nonuniformly expanding maps are obtained in [67]. The results for skew product flows in Section 4.1 should also go over to toral extensions of nonuniformly hyperbolic transformations with φ constant along stable leaves.

For toral extensions that are not skew products, we expect that the methods described in [14] apply when there is exponential contraction along stable leaves (Section 4.2(i)), or when φ has bounded Hölder constants (Section 4.2(ii)), but there is no analogue of situation (iii) from Section 4.2. An open problem is to understand more fully toral extensions of nonuniformly hyperbolic transformations (and hence understand more fully nonuniformly hyperbolic flows).

- Continuing the previous question, although the results in [14] described here deal with many important classes of flows, the situation is still not as satisfactory as for semiflows. For example, Chernov & Zhang [41] consider a family of periodic dispersing billiards where the nonvanishing curvature hypothesis on scatterers is violated. The associated billiard maps exhibit polynomial decay rates $O(n^{-b})$ for any prescribed $b \in (1, \infty)$. However the flows do not seem to be covered by the methods in [14] even though the results in this article yield decay rates $O(t^{-b})$ at the semiflow level.

Acknowledgements This work was supported in part by a European Advanced Grant *StochExtHomog* (ERC AdG 320977). We are grateful to Péter Bálint and Oliver Butterley for very helpful discussions, and to the referees for making numerous suggestions that greatly improved the readability of the paper.

References

- [1] J. Aaronson. *An Introduction to Infinite Ergodic Theory*. Math. Surveys and Monographs **50**, Amer. Math. Soc., 1997.
- [2] J. Aaronson and M. Denker. Local limit theorems for partial sums of stationary sequences generated by Gibbs-Markov maps. *Stoch. Dyn.* **1** (2001) 193–237.
- [3] V. Araújo, O. Butterley and P. Varandas. Open sets of Axiom A flows with exponentially mixing attractors. *Proc. Amer. Math. Soc.* **144** (2016) 2971–2984.
- [4] J. F. Alves and V. Pinheiro. Slow rates of mixing for dynamical systems with hyperbolic structures. *J. Stat. Phys.* **131** (2008) 505–534.
- [5] V. Araújo and I. Melbourne. Exponential decay of correlations for nonuniformly hyperbolic flows with a $C^{1+\alpha}$ stable foliation, including the classical Lorenz attractor. *Ann. Henri Poincaré* **17** (2016) 2975–3004.
- [6] V. Araújo and I. Melbourne. Existence and smoothness of the stable foliation for sectional hyperbolic attractors. *Bull. London Math. Soc.* **49** (2017) 351–367.
- [7] V. Araújo and I. Melbourne. Mixing properties and statistical limit theorems for singular hyperbolic flows. Preprint, 2017.
- [8] V. Araújo, I. Melbourne and P. Varandas. Rapid mixing for the Lorenz attractor and statistical limit laws for their time-1 maps. *Comm. Math. Phys.* **340** (2015) 901–938.
- [9] V. Araujo, M. J. Pacifico, E. R. Pujals and M. Viana. Singular-hyperbolic attractors are chaotic. *Trans. Amer. Math. Soc.* **361** (2009) 2431–2485.
- [10] A. Avila, S. Gouëzel and J. Yoccoz. Exponential mixing for the Teichmüller flow. *Publ. Math. Inst. Hautes Études Sci.* **104** (2006) 143–211.

- [11] V. Baladi, M. Demers and C. Liverani. Exponential decay of correlations for finite horizon Sinai billiard flows. *Invent. Math.* **211** (2018) 39–177.
- [12] V. Baladi and A. Hachemi. A local limit theorem with speed of convergence for Euclidean algorithms and Diophantine costs. *Ann. Inst. Henri Poincaré Probab. Stat.* **44** (2008) 749–770.
- [13] V. Baladi and B. Vallée. Exponential decay of correlations for surface semi-flows without finite Markov partitions. *Proc. Amer. Math. Soc.* **133** (2005) 865–874.
- [14] P. Bálint, O. Butterley and I. Melbourne. Polynomial decay of correlations for flows, including Lorentz gas examples. Preprint, 2017.
- [15] P. Bálint, N. Chernov, D. Szász and I. P. Tóth. Geometry of multi-dimensional dispersing billiards. *Astérisque* (2003), no. 286, xviii, 119–150.
- [16] P. Bálint and S. Gouëzel. Limit theorems in the stadium billiard. *Comm. Math. Phys.* **263** (2006) 461–512.
- [17] P. Bálint and I. Melbourne. Decay of correlations and invariance principles for dispersing billiards with cusps, and related planar billiard flows. *J. Stat. Phys.* **133** (2008) 435–447.
- [18] P. Bálint and A. Némegy Varga. The flow of two falling balls mixes rapidly. *Nonlinearity* **29** (2016) 2537–2564.
- [19] P. Bálint and I. P. Tóth. Exponential decay of correlations in multi-dimensional dispersing billiards. *Ann. Henri Poincaré* **9** (2008) 1309–1369.
- [20] P. Bálint and I. P. Tóth. Example for exponential growth of complexity in a finite horizon multi-dimensional dispersing billiard. *Nonlinearity* **25** (2012) 1275–1297.
- [21] M. Benedicks and L. Carleson. The dynamics of the Hénon map. *Ann. of Math.* **133** (1991) 73–169.
- [22] M. Benedicks and L.-S. Young. Sinai-Bowen-Ruelle measures for certain Hénon maps. *Invent. Math.* **112** (1993) 541–576.
- [23] M. Benedicks and L.-S. Young. Markov extensions and decay of correlations for certain Hénon maps. *Astérisque* (2000), no. 261, 13–56.
- [24] R. Bowen and D. Ruelle. The ergodic theory of Axiom A flows. *Invent. Math.* **29** (1975) 181–202.
- [25] M. I. Brin. Topological transitivity of a certain class of dynamical systems, and flows of frames on manifolds of negative curvature. *Funkcional. Anal. i Priložen.* **9** (1975) 9–19.
- [26] M. I. Brin. The topology of group extensions of C -systems. *Mat. Zametki* **18** (1975) 453–465.
- [27] H. Bruin, M. Holland and I. Melbourne. Subexponential decay of correlations for compact group extensions of nonuniformly expanding systems. *Ergodic Theory Dynam. Systems* **25** (2005) 1719–1738.
- [28] H. Bruin, I. Melbourne and D. Terhesiu. Lower bounds on mixing for nonMarkovian flows. In preparation.
- [29] L. A. Bunimovič. The ergodic properties of billiards that are nearly scattering. *Dokl. Akad. Nauk SSSR* **211** (1973) 1024–1026.
- [30] L. A. Bunimovich. On the ergodic properties of nowhere dispersing billiards. *Comm. Math. Phys.* **65** (1979) 295–312.
- [31] L. A. Bunimovich, Y. G. Sinai and N. I. Chernov. Statistical properties of two-dimensional hyperbolic billiards. *Uspekhi Mat. Nauk* **46** (1991) 43–92.
- [32] K. Burns, H. Masur, C. Matheus and A. Wilkinson. Rates of mixing for the Weil–Petersson geodesic flow: exponential mixing in exceptional moduli spaces. *Geom. Funct. Anal.* **27** (2017) 240–288.
- [33] O. Butterley and K. War. Open sets of exponentially mixing Anosov flows. *J. Eur. Math. Soc.* To appear.
- [34] J.-R. Chazottes and S. Gouëzel. Optimal concentration inequalities for dynamical systems. *Comm. Math. Phys.* **316** (2012) 843–889.
- [35] N. Chernov. Decay of correlations and dispersing billiards. *J. Statist. Phys.* **94** (1999) 513–556.

- [36] N. Chernov. A stretched exponential bound on time correlations for billiard flows. *J. Stat. Phys.* **127** (2007) 21–50.
- [37] N. Chernov and R. Markarian. *Chaotic billiards*. Mathematical Surveys and Monographs **127**, American Mathematical Society, Providence, RI, 2006.
- [38] N. Chernov and L. S. Young. Decay of correlations for Lorentz gases and hard balls. *Hard ball systems and the Lorentz gas*, Encyclopaedia Math. Sci. **101**, Springer, Berlin, 2000, pp. 89–120.
- [39] N. I. Chernov. Markov approximations and decay of correlations for Anosov flows. *Ann. of Math.* **147** (1998) 269–324.
- [40] N. I. Chernov and H.-K. Zhang. Billiards with polynomial mixing rates. *Nonlinearity* **18** (2005) 1527–1553.
- [41] N. Chernov and H.-K. Zhang. A family of chaotic billiards with variable mixing rates. *Stoch. Dyn.* **5** (2005) 535–553.
- [42] N. I. Chernov and H.-K. Zhang. Improved estimates for correlations in billiards. *Comm. Math. Phys.* **77** (2008) 305–321.
- [43] C. Cuny and F. Merlevède. Strong invariance principles with rate for “reverse” martingales and applications. *J. Theor. Probab.* (2015) 137–183.
- [44] J. De Simoi and I. P. Tóth. An expansion estimate for dispersing planar billiards with corner points. *Ann. Henri Poincaré* **15** (2014) 1223–1243.
- [45] M. Denker and W. Philipp. Approximation by Brownian motion for Gibbs measures and flows under a function. *Ergodic Theory Dynam. Systems* **4** (1984) 541–552.
- [46] L. J. Díaz, J. Rocha and M. Viana. Strange attractors in saddle-node cycles: prevalence and globality. *Invent. Math.* **125** (1996) 37–74.
- [47] D. Dolgopyat. On the decay of correlations in Anosov flows. *Ann. of Math.* **147** (1998) 357–390.
- [48] D. Dolgopyat. Prevalence of rapid mixing in hyperbolic flows. *Ergodic Theory Dynam. Systems* **18** (1998) 1097–1114.
- [49] D. Dolgopyat. On mixing properties of compact group extensions of hyperbolic systems. *Israel J. Math.* **130** (2002) 157–205.
- [50] M. J. Field, I. Melbourne and A. Török. Stability of mixing and rapid mixing for hyperbolic flows. *Ann. of Math.* **166** (2007) 269–291.
- [51] B. Friedman and R. Martin. Behavior of the velocity autocorrelation function for the periodic Lorentz gas. *Phys. D* **30** (1988) 219–227.
- [52] S. Gouëzel. Central limit theorem and stable laws for intermittent maps. *Probab. Theory Relat. Fields* **128** (2004) 82–122.
- [53] S. Gouëzel. Sharp polynomial estimates for the decay of correlations. *Israel J. Math.* **139** (2004) 29–65.
- [54] S. Gouëzel. Private communication.
- [55] H. Hennion. Sur un théorème spectral et son application aux noyaux lipchitziens. *Proc. Amer. Math. Soc.* **118** (1993) 627–634.
- [56] H. Hu. Decay of correlations for piecewise smooth maps with indifferent fixed points. *Ergodic Theory Dynam. Systems* **24** (2004) 495–524.
- [57] A. Katok. Infinitesimal Lyapunov functions, invariant cone families and stochastic properties of smooth dynamical systems. *Ergodic Theory Dynam. Systems* **14** (1994) 757–785. With the collaboration of K. Burns.
- [58] A. Korepanov, Z. Kosloff and I. Melbourne. Explicit coupling argument for nonuniformly hyperbolic transformations. *Proc. Roy. Soc. Edinburgh A*. To appear.
- [59] C. Liverani. On contact Anosov flows. *Ann. of Math.* **159** (2004) 1275–1312.
- [60] C. Liverani, B. Saussol and S. Vaienti. A probabilistic approach to intermittency. *Ergodic Theory Dynam. Systems* **19** (1999) 671–685.

- [61] R. Markarian. Billiards with polynomial decay of correlations. *Ergodic Theory Dynam. Systems* **24** (2004) 177–197.
- [62] H. Matsuoka and R. F. Martin. Long-time tails of the velocity autocorrelation functions for the triangular periodic Lorentz gas. *J. Stat. Phys* **88** (1997) 81–103.
- [63] I. Melbourne. Rapid decay of correlations for nonuniformly hyperbolic flows. *Trans. Amer. Math. Soc.* **359** (2007) 2421–2441.
- [64] I. Melbourne. Decay of correlations for slowly mixing flows. *Proc. London Math. Soc.* **98** (2009) 163–190.
- [65] I. Melbourne and D. Terhesiu. Decay of correlations for nonuniformly expanding systems with general return times. *Ergodic Theory Dynam. Systems* **34** (2014) 893–918.
- [66] I. Melbourne and D. Terhesiu. Operator renewal theory for continuous time dynamical systems with finite and infinite measure. *Monatsh. Math.* **182** (2017) 377–431.
- [67] I. Melbourne and D. Terhesiu. Mixing properties for toral extensions of slowly mixing dynamical systems with finite and infinite measure. *J. Mod. Dyn.* To appear.
- [68] I. Melbourne and A. Török. Central limit theorems and invariance principles for time-one maps of hyperbolic flows. *Comm. Math. Phys.* **229** (2002) 57–71.
- [69] I. Melbourne and A. Török. Statistical limit theorems for suspension flows. *Israel J. Math.* **144** (2004) 191–209.
- [70] I. Melbourne and P. Varandas. A note on statistical properties for nonuniformly hyperbolic systems with slow contraction and expansion. *Stoch. Dyn.* **16** (2016) 1660012. 13 pages.
- [71] L. Mora and M. Viana. Abundance of strange attractors. *Acta Math.* **171** (1993) 1–71.
- [72] W. Philipp and W. F. Stout. *Almost Sure Invariance Principles for Partial Sums of Weakly Dependent Random Variables*. Memoirs of the Amer. Math. Soc. **161**, Amer. Math. Soc., Providence, RI, 1975.
- [73] M. Pollicott. A complex Ruelle-Perron-Frobenius theorem and two counterexamples. *Ergodic Theory Dynam. Systems* **4** (1984) 135–146.
- [74] M. Pollicott. On the rate of mixing of Axiom A flows. *Invent. Math.* **81** (1985) 413–426.
- [75] M. Pollicott. On the rate of mixing of Axiom A attracting flows and a conjecture of Ruelle. *Ergodic Theory Dynam. Systems* **19** (1999) 535–548.
- [76] Y. Pomeau and P. Manneville. Intermittent transition to turbulence in dissipative dynamical systems. *Comm. Math. Phys.* **74** (1980) 189–197.
- [77] M. Ratner. The central limit theorem for geodesic flows on n -dimensional manifolds of negative curvature. *Israel J. Math.* **16** (1973) 181–197.
- [78] D. Ruelle. Flows which do not exponentially mix. *C. R. Acad. Sci. Paris* **296** (1983) 191–194.
- [79] O. M. Sarig. Subexponential decay of correlations. *Invent. Math.* **150** (2002) 629–653.
- [80] Y. G. Sinaĭ. Dynamical systems with elastic reflections. Ergodic properties of dispersing billiards. *Uspehi Mat. Nauk* **25** (1970) 141–192.
- [81] L. Stoyanov. Spectrum of the Ruelle operator and exponential decay of correlations for open billiard flows. *Amer. J. Math.* **123** (2001) 715–759.
- [82] M. Tsujii. Quasi-compactness of transfer operators for contact Anosov flows. *Nonlinearity* **23** (2010) 1495–1545.
- [83] M. Tsujii. Exponential mixing for generic volume-preserving Anosov flows in dimension three. *J. Math. Soc. Japan* **70** (2018) 757–821.
- [84] M. Viana. Strange attractors in higher dimensions. *Bol. Soc. Brasil. Mat. (N.S.)* **24** (1993) 13–62.
- [85] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math.* **147** (1998) 585–650.
- [86] L.-S. Young. Recurrence times and rates of mixing. *Israel J. Math.* **110** (1999) 153–188.