

VARIANCE CONTINUITY FOR LORENZ FLOWS

Wael Bahsoun[†], Ian Melbourne^{*}, and Marks Ruziboev[‡]

ABSTRACT. The classical Lorenz flow, and any flow which is close to it in the C^2 -topology, satisfies a Central Limit Theorem (CLT). We prove that the variance in the CLT varies continuously.

1. INTRODUCTION

In 1963, Lorenz [20] introduced the following system of equations

$$\begin{cases} \dot{x} = -10x + 10y \\ \dot{y} = 28x - y - xz \\ \dot{z} = -\frac{8}{3}z + xy \end{cases} \quad (1)$$

as a simplified model for atmospheric convection. Numerical simulations performed by Lorenz showed that the above system exhibits sensitive dependence on initial conditions and has a non-periodic “strange” attractor. Since then, (1) became a basic example of a chaotic deterministic system that is notoriously difficult to analyse.

A rigorous mathematical framework of similar flows was initiated with the introduction of the so called geometric Lorenz attractors in [1, 15]. The papers [24, 25] provided a computer-assisted proof that the classical Lorenz attractor in (1) is indeed a geometric Lorenz attractor. In particular, it is a singularly hyperbolic attractor [23], namely a nontrivial robustly transitive attracting invariant set containing a singularity (equilibrium point). Moreover, there is a distinguished Sinai-Ruelle-Bowen (SRB) ergodic probability measure, see for example [8]. Statistical limit laws, in particular the central limit theorem (CLT) for Hölder observables, were first obtained in [16] for the classical Lorenz attractor, and were shown for general singular hyperbolic attractors in [5]. For further background and a complete list of references up to 2010 we refer the reader to the monograph of Araújo and Pacifico [7].

^{†,‡} DEPARTMENT OF MATHEMATICAL SCIENCES, LOUGHBOROUGH UNIVERSITY, LOUGHBOROUGH, LE11 3TU, UK

^{*} MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UK
E-mail addresses: [†] W.Bahsoun@lboro.ac.uk, ^{*} I.Melbourne@warwick.ac.uk,

[‡] M.Ruziboev@lboro.ac.uk.

Date: April 11, 2020.

1991 *Mathematics Subject Classification.* Primary 37A05, 37C10, 37E05.

Key words and phrases. Lorenz flow, Statistical stability, Central Limit Theorem.

WB and MR would like to thank The Leverhulme Trust for supporting their research through the research grant RPG-2015-346. The research of IM was supported in part by a European Advanced Grant StochExtHomog (ERC AdG 320977). We are very grateful to the referee for important comments that have led to a much more readable explanation of the method in this paper.

Let $X_\varepsilon : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$, $\varepsilon \geq 0$, be a continuous family of C^2 flows on \mathbb{R}^3 admitting a geometric Lorenz attractor with singularities x_ε and corresponding SRB measures μ_ε . Precise definitions are given in section 2.1; in particular, the framework includes the classical Lorenz attractor. By [2, 9], the flows X_ε are statistically stable: for any continuous $\psi : \mathbb{R}^3 \rightarrow \mathbb{R}$

$$\lim_{\varepsilon \rightarrow 0} \int \psi d\mu_\varepsilon = \int \psi d\mu_0.$$

The CLT in [5, 16] states that for fixed $\varepsilon \geq 0$ and $\psi : \mathbb{R}^3 \rightarrow \mathbb{R}$ Hölder, there exists $\sigma^2 = \sigma_{X_\varepsilon}^2(\psi) \geq 0$ such that

$$\frac{1}{\sqrt{t}} \left(\int_0^t \psi \circ X_\varepsilon(s) ds - t \int \psi d\mu_\varepsilon \right) \xrightarrow{\text{law}} \mathcal{N}(0, \sigma^2) \quad \text{as } t \rightarrow \infty. \quad (2)$$

By [16, Section 4.3], σ^2 is typically nonzero.

We prove continuity of the variance, namely that $\varepsilon \mapsto \sigma_{X_\varepsilon}^2(\psi)$ is continuous. At the same time, we obtain estimates on the dependence of the variance on ψ . We now state the main result of the paper. Define $\|\psi\| = \int |\psi| d\mu_0 + |\psi(x_0)|$.

Theorem 1.1. *Let $\psi, \psi' : \mathbb{R}^3 \rightarrow \mathbb{R}$ be Hölder observables. Then*

- (a) $\lim_{\varepsilon \rightarrow 0} \sigma_{X_\varepsilon}^2(\psi) = \sigma_{X_0}^2(\psi)$; and
- (b) *there exists a constant $C > 0$ (depending only on the Hölder norms of ψ and ψ') such that*

$$|\sigma_{X_0}^2(\psi) - \sigma_{X_0}^2(\psi')| \leq C \|\psi - \psi'\| (1 + |\log \|\psi - \psi'\||).$$

Remark 1.2. *In part (b), we obtain closeness of the variances provided the observables are close in L^1 with respect to both the SRB measure and the Dirac point mass at 0 (provided the individual Hölder norms are controlled). It is an easy consequence of the methods in this paper that the logarithmic factor can be removed if the norm $\|\cdot\|$ is replaced by the Hölder norm.*

Remark 1.3. *All results in this paper go through without change for continuous families of C^r flows, $r > 1$. We take $r = 2$ for notational convenience.*

Our technique is based on first proving variance continuity for the corresponding family of Poincaré maps and then passing the result to the family of flows. The main difficulty in passing from maps to flows lies in the fact that the return time function to the Poincaré section is unbounded. A key step in overcoming this hurdle is to show that for any Hölder observable on \mathbb{R}^3 vanishing at the singularity, the induced observable for the Poincaré map is piecewise Hölder. Related results for various classes of discrete time dynamical systems can be found in [11, 13, 19] using somewhat different methods, but there are no previous results for Lorenz attractors.

The paper is organised as follows. In section 2 we recall the basic setup and notation associated with (families of) geometric Lorenz attractors. In section 3 we show how to normalise the families of flows to obtain simplified coordinates for the proofs. Section 4 contains properties of one dimensional Lorenz maps. Section 5 studies the family of Poincaré maps. It starts by showing that the family of maps admit a uniform rate of correlations decay for piecewise Hölder functions, using suitable anisotropic norms. We

then use the Green-Kubo formula to show continuity of the variance for the family of Poincaré maps. In section 6 we prove a version of Theorem 1.1 for normalised families and use this to prove Theorem 1.1.

2. GEOMETRIC LORENZ ATTRACTORS

In this section, we recall the basic setup and notation associated with (families of) geometric Lorenz attractors. In subsections 2.1 and 2.2, we recall the class of (families of) geometric Lorenz attractors considered in the paper.

We begin with some notational preliminaries. Let $U \subset \mathbb{R}^m$ be open. Fix $\alpha \in (0, 1)$ and recall that $f : U \rightarrow \mathbb{R}^n$ is C^α if there exists $C > 0$ such that $|f_j(x) - f_j(y)| \leq C|x - y|^\alpha$ for all $x, y \in U$ and all $j = 1, \dots, n$. (Here $|x| = \sqrt{x_1^2 + \dots + x_m^2}$ denotes the Euclidean norm on \mathbb{R}^m .) Let $H_\alpha(f)$ be the least such constant C and define the Hölder norm $\|f\|_\alpha = |f|_\infty + H_\alpha(f)$. Then f is $C^{1+\alpha}$ if $Df : U \rightarrow \mathbb{R}^{n \times m}$ is C^α and we set $\|f\|_{1+\alpha} = |f|_\infty + \|Df\|_\alpha$. A family f_ε of $C^{1+\alpha}$ maps, $\varepsilon \geq 0$, is said to be continuous if $\lim_{\varepsilon \rightarrow \varepsilon_0} \|f_\varepsilon - f_{\varepsilon_0}\|_{1+\alpha} = 0$. Similarly, we speak of continuous families of C^2 flows, $C^{1+\alpha}$ diffeomorphisms, and so on. In the case of Lorenz flows, we are particularly interested in families of C^2 flows on \mathbb{R}^3 restricted to an open bounded region U of phase space; for convenience we suppress mentioning the subset U .

2.1. Definition of geometric Lorenz attractors. There are various notions of geometric Lorenz attractor in the literature depending on the properties being analysed. Roughly speaking, we take a geometric Lorenz attractor to be a singular hyperbolic attractor for a vector field on \mathbb{R}^3 with a single singularity x_0 and a connected global cross-section with a $C^{1+\alpha}$ stable foliation. As promised, we now give a precise description.

Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a C^2 vector field satisfying $G(x_0) = 0$ and let X be the associated flow. We suppose that the differential $DG(x_0)$ at the singularity has three real eigenvalues $\lambda_2 < \lambda_3 < 0 < \lambda_1$ satisfying $\lambda_1 + \lambda_3 > 0$ (Lorenz-like singularity).

Let Σ be a two-dimensional rectangular cross-section transverse to the flow chosen in a neighbourhood of the singularity x_0 , and let Γ be the intersection of Σ with the local stable manifold of x_0 . We suppose that there exists a well defined Poincaré map $F : \Sigma \setminus \Gamma \rightarrow \Sigma$. Moreover, we assume that the underlying flow is singular hyperbolic [23]. It follows [4, Theorem 4.2] that a neighbourhood of the attractor is foliated by one-dimensional C^2 stable leaves. We assume q -bunching for some $q > 1$ in [4, condition (4.2)]. By [4, Theorem 4.12 and Remark 4.13(b)], it follows that the stable foliation for the flow is C^q .

The foliation by stable leaves for the flow naturally induces (see for example [5, Section 3.1]) a C^q ($q > 1$) foliation inside Σ of a neighbourhood of the attractor intersected with Σ by one-dimensional C^2 stable leaves for the Poincaré map F . We denote this stable foliation for F by \mathcal{F} .

The stable leaves for the flow are exponentially contracting [4, Theorem 4.2(a)(3)] and this property is inherited by the stable leaves for F . This means that there exists $\rho \in (0, 1)$, $K > 0$ such that

$$|F^n \xi_1 - F^n \xi_2| \leq K \rho^n |\xi_1 - \xi_2|, \quad (3)$$

for ξ_1, ξ_2 in the same stable leaf in \mathcal{F} and $n \geq 1$.

Let $I \subset \Sigma$ be a smoothly (C^∞) embedded one-dimensional subspace transverse to the stable foliation and let $T : I \rightarrow I$ be the one-dimensional map obtained from F by quotienting along stable leaves. Let ξ_0 be the intersection of I with Γ .

Proposition 2.1. *T is a Lorenz-like expanding map. That is, T is monotone (without loss we take T to be increasing) and piecewise $C^{1+\alpha}$ on $I \setminus \{\xi_0\}$ for some $\alpha \in (0, 1)$ with a singularity at ξ_0 and one-sided limits $T(\xi_0^+) < 0$ and $T(\xi_0^-) > 0$. Also, T is uniformly expanding: there are constants $c > 0$ and $\theta > 1$ such that $(T^n)'(x) \geq c\theta^n$ for all $n \geq 1$, whenever $x \notin \bigcup_{j=0}^{n-1} T^{-j}(\xi_0)$.*

Proof. The map F is piecewise $C^{1+\alpha}$ and the foliation by stable leaves is $C^{1+\alpha}$, so T is piecewise $C^{1+\alpha}$ on $I \setminus \{\xi_0\}$. Uniform expansion follows from [5, Theorem 4.3]. The remaining properties are immediate. \square

The final part of the definition of geometric Lorenz attractor is that the one-dimensional map T is transitive on the interval $[T(\xi_0^+), T(\xi_0^-)]$. It is then standard [2, 5, 8, 17] that T has a unique absolutely continuous invariant probability measure (acip) $\bar{\mu}$ leading to a unique SRB measure μ for the geometric Lorenz attractor containing x_0 . The basin of μ has full Lebesgue measure in a neighbourhood of the attractor.

Remark 2.2. *The classical Lorenz attractor for the system of equations (1) (and for nearby equations) is an example of a geometric Lorenz attractor as defined above. Except for q -bunching, the assumptions above are verified in [25]. The q -bunching condition is verified in [6, Lemma 2.2]. (By [4, Section 5], the optimal value of q lies between 1.278 and 1.705; hence we have $C^{1+\alpha}$ regularity for the stable foliation as in [9] but not C^2 regularity as in [2].)*

2.2. Families of geometric Lorenz attractors. Let X_ε , $\varepsilon \geq 0$, be a continuous family of C^2 flows on \mathbb{R}^3 admitting a geometric Lorenz attractor as in subsection 2.1 with singularity x_ε . The constants K and ρ in (3) derive from the singular hyperbolic structure which varies continuously under C^1 perturbations. Hence K and ρ can be chosen independent of ε .

Making an initial C^2 change of coordinates (varying continuously in ε), we can suppose without loss that $x_\varepsilon \equiv 0$ and that Σ , Γ and I are given by $\Sigma = \{(x, y, 1) : -1 \leq x \leq 1, -\frac{1}{2} \leq y \leq \frac{1}{2}\}$, $\Gamma = \{(0, y, 1) : -\frac{1}{2} \leq y \leq \frac{1}{2}\}$ and $I = \{(x, 0, 1) : -\frac{1}{2} \leq x \leq \frac{1}{2}\} \cong [-\frac{1}{2}, \frac{1}{2}]$ for all ε . Throughout the paper, when we speak of a continuous family of C^2 flows admitting geometric Lorenz attractors, we assume that this initial change of coordinates has been performed.

Define the Poincaré return time to Σ ,

$$\tau_\varepsilon : \Sigma \rightarrow (0, \infty), \quad \tau_\varepsilon(\xi) = \inf\{t > 0 : X_\varepsilon(\xi, t) \in \Sigma\}.$$

Proposition 2.3. *The return time $\tau_\varepsilon : \Sigma \setminus \Gamma \rightarrow \Sigma$ is given by $\tau_\varepsilon(x, y) = -\frac{1}{\lambda_{1,\varepsilon}} \log |x| + \tau_{2,\varepsilon}(x, y)$ where $\tau_{2,\varepsilon}$ is a continuous family of C^2 functions.*

Proof. Applying the Hartman-Grobman theorem for fixed ε , we can linearise X_ε in a neighbourhood of 0 so that the linearised flow is given by $(x, y, z) \mapsto$

$(e^{\lambda_{1,\varepsilon}t}x, e^{\lambda_{2,\varepsilon}t}y, e^{\lambda_{3,\varepsilon}t}z)$. The time of flight in this neighbourhood is readily calculated in these coordinates to be $-\frac{1}{\lambda_{1,\varepsilon}} \log|x|$ for $x \in I$ and the same formula holds in the original coordinates. The remaining flight time $\tau_{2,\varepsilon}$ is a first hit-time for the C^2 flow away from the singularity at 0 and hence is C^2 . Since X_ε is a continuous family of C^2 flows, it follows that $\tau_{2,\varepsilon}$ is a continuous family of C^2 functions. \square

Theorem 2.4. *Let X_ε be a continuous family of C^2 flows on \mathbb{R}^3 admitting a geometric Lorenz attractor. Then there exists $\alpha > 0$ such that the one-dimensional maps $T_\varepsilon : I \rightarrow I$ form a continuous family of piecewise $C^{1+\alpha}$ maps.*

Proof. Recall that T_ε is obtained from the continuous family of piecewise $C^{1+\alpha}$ maps F_ε by quotienting along the stable foliation. Our assumption of q -bunching ($q > 1$) yields continuous families of C^q stable foliations [12]. Hence, T_ε is a continuous family of piecewise $C^{1+\alpha}$ maps. \square

3. NORMALISED GEOMETRIC LORENZ ATTRACTORS

Let X_ε , $\varepsilon \geq 0$, be a continuous family of C^2 flows on \mathbb{R}^3 admitting a geometric Lorenz attractor. In this section, we show how to normalise the families of flows to obtain simplified coordinates for carrying out the proofs.

Assume that the preliminary C^2 change of coordinates in section 2.2 has been performed. Let $F_\varepsilon : \Sigma \rightarrow \Sigma$ and $T_\varepsilon : I \rightarrow I$ be the associated families of Poincaré maps and one-dimensional piecewise expanding maps. Also, define $\tilde{\Sigma} = \{(x, y, 1) : -\frac{1}{2} \leq x, y \leq \frac{1}{2}\}$.

Proposition 3.1. *There exists a continuous family of $C^{1+\alpha}$ diffeomorphisms $\omega_\varepsilon : \Sigma \rightarrow \Sigma$ such that*

- (a) ω_ε restricts to the identity on I ; and
- (b) vertical lines in $\tilde{\Sigma}$ are transformed under ω_ε into stable leaves for $F_\varepsilon : \Sigma \rightarrow \Sigma$.

Proof. For ε fixed, this follows by definition of the smoothness of the stable foliation for F_ε . (An explicit formula is given in [4, Lemma 4.9] where Y and χ should be replaced by I and ω_ε , and the embedding is the identity.) Again, we obtain continuous families of $C^{1+\alpha}$ diffeomorphisms by [12]. \square

The change of coordinates ω_ε for the Poincaré map F_ε extends to a change of coordinates ϕ_ε for the flow X_ε . The extension is standard and essentially unique, though heavy on notation.

First, define the transformed Poincaré map and return time

$$\tilde{F}_\varepsilon = \omega_\varepsilon^{-1} \circ F_\varepsilon \circ \omega_\varepsilon : \tilde{\Sigma} \rightarrow \tilde{\Sigma}, \quad \tilde{\tau}_\varepsilon = \tau_\varepsilon \circ \omega_\varepsilon : \tilde{\Sigma} \rightarrow (0, \infty).$$

The set $U_\varepsilon = \{(X_\varepsilon(\xi, t) : \xi \in \Sigma, 0 \leq t \leq \tau_\varepsilon(\xi)\}$ defines a neighbourhood of the Lorenz attractor for X_ε . Note that $X_\varepsilon(\xi, \tau_\varepsilon(\xi)) = F_\varepsilon \xi$ for $\xi \in \Sigma$. Define the transformed flow $\tilde{X}_\varepsilon : U_\varepsilon \times [0, \infty) \rightarrow U_\varepsilon$ given by $\tilde{X}_\varepsilon(x, t) = X_\varepsilon(x, t)$ subject to the identifications $\tilde{X}_\varepsilon(\xi, \tilde{\tau}_\varepsilon(\xi)) = \tilde{F}_\varepsilon \xi$ for $\xi \in \tilde{\Sigma}$.

Finally, define $\phi_\varepsilon : U_\varepsilon \rightarrow U_\varepsilon$ by $\phi_\varepsilon(x) = X_\varepsilon(\omega_\varepsilon(\xi), t)$ for $x = X_\varepsilon(\xi, t) \in U_\varepsilon$, where $\xi \in \Sigma$ and $0 \leq t \leq \tau_\varepsilon(x)$. It follows from the definitions that $\phi_\varepsilon|_{\Sigma_\varepsilon} \equiv \omega_\varepsilon$ and $\tilde{X}_\varepsilon(x, t) = \phi_\varepsilon^{-1} \circ X_\varepsilon(\phi_\varepsilon(x), t)$. Hence ϕ_ε is the desired change of coordinates.

Corollary 3.2. *The changes of coordinates $\phi_\varepsilon : U \rightarrow U$ are continuous families of $C^{1+\alpha}$ diffeomorphisms.*

Proof. By assumption, X_ε is a continuous family of C^2 flows. Hence the result follows from Proposition 3.1. \square

Lemma 3.3. *The transformed flow $\tilde{X}_\varepsilon(x, t) = \phi_\varepsilon^{-1} \circ X_\varepsilon(\phi_\varepsilon(x), t)$ satisfies:*

- (i) *The Poincaré map $\tilde{F}_\varepsilon : \tilde{\Sigma} \setminus \Gamma \rightarrow \tilde{\Sigma}$ is a continuous family of piecewise $C^{1+\alpha}$ diffeomorphisms and has the form*

$$\tilde{F}_\varepsilon(x, y) = (T_\varepsilon x, g_\varepsilon(x, y)),$$

where $T_\varepsilon : I \rightarrow I$ is the family of Lorenz-like expanding maps in Theorem 2.4(a).

- (ii) *$|\tilde{F}_\varepsilon^n \xi_1 - \tilde{F}_\varepsilon^n \xi_2| \leq K\rho^n |\xi_1 - \xi_2|$ for all $\xi_1 = (x, y_1)$, $\xi_2 = (x, y_2)$ and $n \geq 1$,*
- (iii) *The return time $\tilde{\tau}_\varepsilon : \tilde{\Sigma} \setminus \Gamma \rightarrow (0, \infty)$ is given by $\tilde{\tau}_\varepsilon(x, y) = -\frac{1}{\lambda_{1,\varepsilon}} \log |x| + \tilde{\tau}_{2,\varepsilon}(x, y)$ where $\tilde{\tau}_{2,\varepsilon}$ is a continuous family of $C^{1+\alpha}$ functions.*
- (iv) *The SRB measures $\tilde{\mu}_\varepsilon$ for \tilde{X}_ε are given by $\tilde{\mu}_\varepsilon = \phi_\varepsilon^{-1} * \mu_\varepsilon$ and are statistically stable.*

Proof. (i) By Proposition 3.1(b), $\tilde{F}_\varepsilon = \omega_\varepsilon^{-1} \circ F_\varepsilon \circ \omega_\varepsilon$ has the form $\tilde{F}_\varepsilon(x, y) = (T_\varepsilon x, g_\varepsilon(x, y))$. By Proposition 3.1(a), the maps $T_\varepsilon : I \rightarrow I$ are unchanged by this change of coordinates. Also, ω_ε is a continuous family of $C^{1+\alpha}$ diffeomorphisms and F_ε is a continuous family of piecewise $C^{1+\alpha}$ diffeomorphisms, so \tilde{F}_ε a continuous family of piecewise $C^{1+\alpha}$ diffeomorphisms.

(ii) This is the uniform contraction condition (3) in the new coordinates.

(iii) By Proposition 2.3 and Proposition 3.1, $\tilde{\tau}_\varepsilon = \tau_\varepsilon \circ \omega_\varepsilon = -\frac{1}{\lambda_{1,\varepsilon}} \log |x| + \tau_{2,\varepsilon} \circ \omega_\varepsilon$ where $\tilde{\tau}_{2,\varepsilon} = \tau_{2,\varepsilon} \circ \omega_\varepsilon$ is a continuous family of $C^{1+\alpha}$ functions.

(iv) By Proposition 3.1(a), the acip $\bar{\mu}_\varepsilon$ for T_ε is unchanged by the change of coordinates and hence remains absolutely continuous. Using this and the construction of the SRB measure (see for example [2, 8] for the standard construction of μ_ε from $\bar{\mu}_\varepsilon$), we obtain that $\tilde{\mu}_\varepsilon = \phi_\varepsilon^{-1} * \mu_\varepsilon$ is the SRB measure for \tilde{X}_ε . Moreover, strong statistical stability [2, 9, 14] of the acips $\bar{\mu}_\varepsilon$ on I is preserved and hence the SRB measures $\tilde{\mu}_\varepsilon$ remain statistically stable. \square

In the following sections, we prove Theorem 1.1 for normalised families of geometric Lorenz attractors. At the end of section 6, we show how results for normalised families imply Theorem 1.1.

4. THE FAMILY OF ONE-DIMENSIONAL MAPS

In this section, we recall some functional-analytic properties associated with the family of one-dimensional Lorenz maps T_ε . For $p \geq 1$, we say $f : I \rightarrow \mathbb{R}$ is of (universally) bounded p -variation if

$$V_p(f) = \sup_{-\frac{1}{2} = x_0 < \dots < x_n = \frac{1}{2}} \left(\sum_{i=1}^n |f(x_i) - f(x_{i-1})|^p \right)^{1/p} < \infty.$$

We take $p \geq \frac{1}{\alpha}$.

Let $S_\rho(x) = \{y \in I : |x - y| < \rho\}$. For $f : I \rightarrow \mathbb{R}$, define

$$\text{osc}(f, \rho, x) = \text{esssup}\{|f(y_1) - f(y_2)| : y_1, y_2 \in S_\rho(x)\},$$

and

$$\text{osc}_1(f, \rho) = \|\text{osc}(f, \rho, x)\|_1,$$

where the essential supremum is taken with respect to Lebesgue measure on $I \times I$ and $\|\cdot\|_1$ is the L^1 -norm with respect to Lebesgue measure on I . Fix $\rho_0 > 0$ and let $BV_{1,1/p} \subset L^1$ be the Banach space equipped with the norm

$$\|f\|_{1,1/p} = V_{1,1/p}(f) + \|f\|_1, \quad \text{where} \quad V_{1,1/p}(f) = \sup_{0 < \rho \leq \rho_0} \frac{\text{osc}_1(f, \rho)}{\rho^{1/p}}.$$

(The space $BV_{1,1/p}$ does not depend on ρ_0 .) The fact that $BV_{1,1/p}$ is a Banach space is proved in [17]. Moreover, it is proved in [17] that $BV_{1,1/p}$ is embedded in L^∞ and compactly embedded in L^1 . In addition [17] shows that

$$V_{1,1/p}(f) \leq 2^{1/p} V_p(f). \quad (4)$$

We recall results from the literature that we use later in sections 5 and 6 of the paper. Recall from [17] that T_ε admits a unique acip $\bar{\mu}_\varepsilon$ for each $\varepsilon \geq 0$. Let h_ε denote the density for $\bar{\mu}_\varepsilon$. Let $P_\varepsilon : L^1(I) \rightarrow L^1(I)$ denote the transfer operator (Perron-Frobenius) associated with T_ε (so $\int P_\varepsilon f g d\text{Leb} = \int f g \circ T_\varepsilon d\text{Leb}$ for $f \in L^1(I)$, $g \in L^\infty(I)$).

Proposition 4.1. *There exists $C > 0$ and $\Lambda \in (0, 1)$ such that*

- (a) $\|h_\varepsilon\|_\infty < C$, and
- (b) $\|P_\varepsilon^n f - h_\varepsilon \int f d\text{Leb}\|_{1,1/p} \leq C\Lambda^n \|f\|_{1,1/p}$

for all $n \geq 1$, $\varepsilon \geq 0$, $f \in BV_{1,1/p}$.

Proof. By [9, Lemma 3.3], there exist $A_1, A_2 > 0$, $0 < \kappa < 1$, such that for all $n \geq 1$, $f \in BV_{1,1/p}$,

$$\|P_\varepsilon^n f\|_{1,1/p} \leq A_1 \kappa^n \|f\|_{1,1/p} + A_2 \|f\|_1.$$

Taking $f = h_\varepsilon$ and letting $n \rightarrow \infty$, we obtain $\|h_\varepsilon\|_{1,1/p} \leq A_2$ and part (a) follows. Moreover, it was proved in [9] that

$$\lim_{\varepsilon \rightarrow 0} \sup_{\|f\|_{1,1/p} \leq 1} \|(P_\varepsilon - P_0)f\|_1 = 0.$$

Thus, the Keller-Liverani stability result [18] implies that P_ε has a uniform (in ε) spectral gap on $BV_{1,1/p}$. This proves part (b). \square

5. VARIANCE CONTINUITY FOR THE POINCARÉ MAPS

In this section, we prove the analogue of Theorem 1.1 at the level of the Poincaré maps F_ε . Throughout, we work with normalised families as in section 3.

It is well known [8] that F_ε admits a unique SRB measure μ_{F_ε} for each $\varepsilon \geq 0$. Moreover, for any continuous $\psi : \Sigma \rightarrow \mathbb{R}$ we have

$$\lim_{\varepsilon \rightarrow 0} \int \psi d\mu_{F_\varepsilon} = \int \psi d\mu_{F_0};$$

i.e. the family F_ε is statistically stable [2, 9, 14]. We require the following strengthening of this property. In general, we say that $\Psi : \Sigma \rightarrow \mathbb{R}$ is piecewise continuous if it is uniformly continuous on the connected components of $\Sigma \setminus \Gamma$. Similarly Ψ is piecewise Hölder if it is uniformly Hölder on the connected components of $\Sigma \setminus \Gamma$.

Proposition 5.1. *Suppose that $\Psi : \Sigma \rightarrow \mathbb{R}$ is piecewise continuous and fix $n \geq 0$. Then $\lim_{\varepsilon \rightarrow 0} \int \Psi \cdot (\Psi \circ F_0^n) d\mu_{F_\varepsilon} = \int \Psi \cdot (\Psi \circ F_0^n) d\mu_{F_0}$.*

Proof. If $\Psi \cdot (\Psi \circ F_0^n)$ were continuous, this would be immediate from the statement of Proposition 3.3 in [2]. The proof in [2] already accounts for trajectories that visit Γ in finitely many steps, and it is easily checked that the same arguments apply here. \square

For fixed $\varepsilon \geq 0$, the CLT holds (see for instance [21]); i.e. for $\Psi : \Sigma \rightarrow \mathbb{R}$ piecewise Hölder there exists $\sigma^2 = \sigma_{F_\varepsilon}^2(\Psi)$ such that

$$\frac{1}{\sqrt{n}} \left(\sum_{i=0}^{n-1} \Psi \circ F_\varepsilon^i - n \int \Psi d\mu_{F_\varepsilon} \right) \xrightarrow{\text{law}} \mathcal{N}(0, \sigma^2) \quad \text{as } n \rightarrow \infty.$$

The variance satisfies the Green-Kubo formula

$$\sigma^2 = \int \hat{\Psi}_\varepsilon^2 d\mu_{F_\varepsilon} + 2 \sum_{n=1}^{\infty} \int \hat{\Psi}_\varepsilon \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon}, \quad (5)$$

where $\hat{\Psi}_\varepsilon = \Psi_\varepsilon - \int \Psi_\varepsilon d\mu_{F_\varepsilon}$.

5.1. Uniform decay of correlations. Let $\alpha \in (0, 1]$ and $p \geq \frac{1}{\alpha}$. Let $\Psi : \Sigma \rightarrow \mathbb{R}$ be piecewise Hölder with exponent α . Set

$$\|\Psi\|_{\alpha, s} = H_{\alpha, s}(\Psi) + \|\Psi\|_{\infty}, \quad H_{\alpha, s}(\Psi) = \sup_{\substack{x, y_1, y_2 \in I \\ y_1 \neq y_2}} \frac{|\Psi(x, y_2) - \Psi(x, y_1)|}{|y_2 - y_1|^\alpha}.$$

For $-\frac{1}{2} = x_0 < \dots < x_n = \frac{1}{2}$ and $y_i \in I$, $1 \leq i \leq n$, we define

$$\hat{V}_p(\Psi; (x_0, \dots, x_n); (y_1, \dots, y_n)) = \left(\sum_{1 \leq i \leq n} |\Psi(x_{i-1}, y_i) - \Psi(x_i, y_i)|^p \right)^{1/p},$$

and

$$\hat{V}_p(\Psi) = \sup \hat{V}_p(\Psi; (x_0, \dots, x_n); (y_1, \dots, y_n)),$$

where the supremum is taken over all finite partitions of I and all choices of $y_i \in I$. Finally, we define

$$(\Pi\Psi)(x) = \int_I \Psi(x, y) dy.$$

We state and prove the following Theorem about uniform (in ε) decay of correlations. Define

$$D_\Psi = \|\Pi\Psi\|_{1, 1/p} + \hat{V}_p(\Psi) + \|\Psi\|_{\alpha, s}.$$

Note that D_Ψ is finite.

Theorem 5.2. *Assume that there exists $K > 0$ such that*

$$\|\Pi(\Psi \circ F_\varepsilon^j)\|_{1,1/p} + \hat{V}_p(\Psi \circ F_\varepsilon^j) \leq D_\Psi K^j \quad (6)$$

for all $j \geq 1$, $\varepsilon \geq 0$. Then there exist $C > 0$ and $\theta \in (0, 1)$ such that

$$\left| \int \Psi \cdot (\Psi \circ F_\varepsilon^n) d\mu_{F_\varepsilon} - \left(\int \Psi d\mu_{F_\varepsilon} \right)^2 \right| \leq CD_\Psi \|\Psi\|_{\alpha,s} \theta^n$$

for all $n \geq 1$, $\varepsilon \geq 0$.

Proof. The proof follows from [3, Theorem 3]. The fact that C and θ do not depend on ε follows from the uniformity of the constants in Proposition 4.1(b) and the uniformity of the contraction on vertical fibres. \square

We now verify assumption (6) in Theorem 5.2.

Lemma 5.3. $V_{1,1/p}(\Pi\Psi) \leq 2^{1/p} \hat{V}_p(\Psi)$.

Proof. Fix $y \in I$ and let $x_0 < \dots < x_n$ be a partition of I . We have

$$\begin{aligned} \sum_{1 \leq i \leq n} |\Pi\Psi(x_{i-1}) - \Pi\Psi(x_i)|^p &= \sum_{1 \leq i \leq n} \left| \int (\Psi(x_{i-1}, y) - \Psi(x_i, y)) dy \right|^p \\ &\leq \sum_{1 \leq i \leq n} \int |\Psi(x_{i-1}, y) - \Psi(x_i, y)|^p dy \leq \hat{V}_p(\Psi)^p. \end{aligned}$$

Combining this with (4) we have $V_{1,1/p}(\Pi\Psi) \leq 2^{1/p} V_p(\Pi\Psi) \leq 2^{1/p} \hat{V}_p(\Psi)$ as required. \square

Recall that F_ε can be written in coordinates as $F_\varepsilon(x, y) = (T_\varepsilon x, g_\varepsilon(x, y))$.

Lemma 5.4. *Let $M = 4(1 + \sup_\varepsilon \|g_\varepsilon\|_{C^1}^\alpha)$. Then*

$$\hat{V}_p(\Psi \circ F_\varepsilon^j) \leq (2^j - 1)M \|\Psi\|_{\alpha,s} + 2^j \hat{V}_p(\Psi) \quad \text{for all } j \geq 1, \varepsilon \geq 0.$$

Proof. We suppress the dependence on ε . Fix $-\frac{1}{2} = x_0 \leq x_1 \leq \dots \leq x_n = \frac{1}{2}$ and $y_i \in I$, $1 \leq i \leq n$. Then

$$\begin{aligned} \hat{V}_p(\Psi \circ F; (x_0, \dots, x_n); (y_1, \dots, y_n))^p &= \sum_{1 \leq i \leq n} |\Psi \circ F(x_{i-1}, y_i) - \Psi \circ F(x_i, y_i)|^p \\ &\leq 2^{p-1} \sum_{1 \leq i \leq n} H_{\alpha,s}(\Psi)^p |g(x_{i-1}, y_i) - g(x_i, y_i)|^{\alpha p} \\ &\quad + 2^{p-1} \sum_{1 \leq i \leq n} |\Psi(Tx_{i-1}, g(x_{i-1}, y_i)) - \Psi(Tx_i, g(x_{i-1}, y_i))|^p \\ &\leq 2^{p-1} H_{\alpha,s}(\Psi)^p \|g\|_{C^1}^{\alpha p} \\ &\quad + 2^{p-1} \sum_{1 \leq i \leq n} |\Psi(Tx_{i-1}, g(x_{i-1}, y_i)) - \Psi(Tx_i, g(x_{i-1}, y_i))|^p. \end{aligned}$$

Let $x_0, \dots, x_k \in [-\frac{1}{2}, 0)$ and $x_{k+1} \notin [-\frac{1}{2}, 0)$. Since $T|_{[-\frac{1}{2}, 0)}$ is continuous and increasing, we get

$$\sum_{1 \leq i \leq k} |\Psi(Tx_{i-1}, g(x_{i-1}, y_i)) - \Psi(Tx_i, g(x_{i-1}, y_i))|^p \leq \hat{V}_p(\Psi)^p.$$

A similar estimate holds for $x_{k+1}, \dots, x_n \in (0, \frac{1}{2}]$. Moreover,

$$|\Psi(Tx_k, g(x_k, y_{k+1})) - \Psi(Tx_{k+1}, g(x_k, y_{k+1}))|^p \leq 2^p \|\Psi\|_{\infty}^p.$$

Consequently,

$$\begin{aligned}\hat{V}_p(\Psi \circ F)^p &\leq 2^{p-1} \left(H_{\alpha,s}(\Psi)^p \|g\|_{C^1}^{\alpha p} + 2\hat{V}_p(\Psi)^p + 2^p \|\Psi\|_{\infty}^p \right) \\ &\leq M^p \|\Psi\|_{\alpha,s}^p + 2^p \hat{V}_p(\Psi)^p \leq (M \|\Psi\|_{\alpha,s} + 2\hat{V}_p(\Psi))^p.\end{aligned}$$

Therefore,

$$\hat{V}_p(\Psi \circ F) \leq M \|\Psi\|_{\alpha,s} + 2\hat{V}_p(\Psi).$$

Using the last inequality repeatedly and the fact that $\|\Psi \circ F\|_{\alpha,s} \leq \|\Psi\|_{\alpha,s}$, we obtain the result. \square

For $\Psi : \Sigma \rightarrow \mathbb{R}$ piecewise C^α , we define

$$\|\Psi\|_{\alpha} = \|\Psi\|_{\infty} + H_{\alpha}(\Psi), \quad H_{\alpha}(\Psi) = \sup_{\xi_1 \neq \xi_2} \frac{|\Psi(\xi_2) - \Psi(\xi_1)|}{|\xi_2 - \xi_1|^{\alpha}},$$

where we restrict to the cases $\xi_1, \xi_2 \in \Sigma^+$ and $\xi_1, \xi_2 \in \Sigma^-$ in the supremum.

Corollary 5.5. *There exists $C > 0$ and $\theta \in (0, 1)$ such that*

$$\left| \int \Psi \cdot (\Psi \circ F_{\varepsilon}^n) d\mu_{F_{\varepsilon}} - \left(\int \Psi d\mu_{F_{\varepsilon}} \right)^2 \right| \leq C \|\Psi\|_{\alpha}^2 \theta^n$$

for all $n \geq 1$, $\varepsilon \geq 0$ and all piecewise C^α observables $\Psi : \Sigma \rightarrow \mathbb{R}$.

Proof. By Lemma 5.3,

$$\|\Pi(\Psi \circ F_{\varepsilon}^j)\|_{1,1/p} + \hat{V}_p(\Psi \circ F_{\varepsilon}^j) \leq \|\Psi\|_{\infty} + (2^{1/p} + 1)\hat{V}_p(\Psi \circ F_{\varepsilon}^j).$$

By Lemma 5.4, there is a constant $K_0 > 1$ such that

$$\|\Pi(\Psi \circ F_{\varepsilon}^j)\|_{1,1/p} + \hat{V}_p(\Psi \circ F_{\varepsilon}^j) \leq K_0 2^j (\|\Psi\|_{\alpha,s} + \hat{V}_p(\Psi)).$$

Hence assumption (6) holds with $K = 2K_0$. The result follows from Theorem 5.2 since $D_{\Psi} \leq 3\|\Psi\|_{\alpha}$ and $\|\Psi\|_{\alpha,s} \leq \|\Psi\|_{\alpha}$. \square

5.2. Variance continuity for the family of two dimensional maps.

We continue to fix $\alpha \in (0, 1]$.

Theorem 5.6. *Let Ψ_{ε} , $\varepsilon \geq 0$, be piecewise C^α with $\sup_{\varepsilon} \|\Psi_{\varepsilon}\|_{\alpha} < \infty$. Assume that*

$$\lim_{\varepsilon \rightarrow 0} \int |\Psi_{\varepsilon} - \Psi_0| d\mu_{F_{\varepsilon}} = 0. \quad (7)$$

Then $\lim_{\varepsilon \rightarrow 0} \sigma_{F_{\varepsilon}}^2(\Psi_{\varepsilon}) = \sigma_{F_0}^2(\Psi_0)$.

(The hypotheses of this result will be verified in section 6. In particular, condition (7) is addressed in Lemma 6.2.)

We first prove a lemma that will be used in the proof of Theorem 5.6.

Lemma 5.7. *Let $\Psi : \Sigma \rightarrow \mathbb{R}$ be piecewise continuous and fix $n \geq 0$. Then*

$$\lim_{\varepsilon \rightarrow 0} \int |\Psi \circ F_0^n - \Psi \circ F_0^{n-1} \circ F_{\varepsilon}| d\mu_{F_{\varepsilon}} = 0.$$

Proof. Let $\delta > 0$, $j \geq 0$, and define $E_{\delta,j} = \bigcup_{i=0}^j F_0^{-i}(B_{\delta})$ where B_{δ} is the δ -neighbourhood of Γ . Then $\mu_{F_0}(E_{\delta,n}) \leq (n+1)\mu_{F_0}(B_{\delta}) \leq M\delta$ where $M = 2(n+1)\|h_0\|_{\infty}$.

The closure of $E_{\delta,n}$ lies in the interior of $E_{2\delta,n}$, so there exists a continuous function $\chi : \Sigma \rightarrow [0, 1]$ supported in $E_{2\delta,n}$ and equal to 1 on $E_{\delta,n}$. By statistical stability, for ε sufficiently small, $\mu_{F_\varepsilon}(E_{\delta,n}) \leq \int \chi d\mu_{F_\varepsilon} \leq 2 \int \chi d\mu_{F_0} \leq 2\mu_{F_0}(E_{2\delta,n}) \leq 4M\delta$.

Let $E_{\delta,j}^c = \Sigma \setminus E_{\delta,j}$. Now $F_0(\xi) \in E_{\delta,n-1}^c$ for $\xi \in E_{\delta,n}^c$. Also, $F_\varepsilon \rightarrow F_0$ uniformly on $E_{\delta,n}^c$ as $\varepsilon \rightarrow 0$, so $F_0(\xi), F_\varepsilon(\xi) \in E_{\delta/2,n-1}^c$ for all $\xi \in E_{\delta,n}^c$ and all sufficiently small ε . Moreover, $\Psi \circ F_0^{n-1}$ is uniformly continuous on $E_{\delta/2,n-1}^c$. It follows from this and the uniform convergence of F_ε on $E_{\delta,n}^c$ that $S_\varepsilon = \sup_{E_{\delta,n}^c} |\Psi \circ F_0^{n-1} \circ F_\varepsilon - \Psi \circ F_0^n| \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Hence

$$\begin{aligned} \int |\Psi \circ F_0^n - \Psi \circ F_0^{n-1} \circ F_\varepsilon| d\mu_{F_\varepsilon} &\leq \int_{E_{\delta,n}} |\Psi \circ F_0^n - \Psi \circ F_0^{n-1} \circ F_\varepsilon| d\mu_{F_\varepsilon} \\ &\quad + \int_{E_{\delta,n}^c} |\Psi \circ F_0^n - \Psi \circ F_0^{n-1} \circ F_\varepsilon| d\mu_{F_\varepsilon} \\ &\leq 2\|\Psi\|_\infty \mu_{F_\varepsilon}(E_{\delta,n}) + S_\varepsilon \leq 8\|\Psi\|_\infty M\delta + S_\varepsilon \rightarrow 8\|\Psi\|_\infty M\delta. \end{aligned}$$

The result follows since $\delta > 0$ is arbitrary. \square

Proof of Theorem 5.6. We use the Green-Kubo formula (5). Recall that $\hat{\Psi}_\varepsilon = \Psi_\varepsilon - \int \Psi_\varepsilon d\mu_{F_\varepsilon}$. By Corollary 5.5, the series in (5) is absolutely convergent uniformly in ε . Therefore, it suffices to show that $\int \hat{\Psi}_\varepsilon \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon} \rightarrow \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_0^n) d\mu_{F_0}$ as $\varepsilon \rightarrow 0$ for each fixed $n \geq 0$. Now

$$\begin{aligned} &\int \hat{\Psi}_\varepsilon \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon} - \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_0^n) d\mu_{F_0} \\ &= \int \hat{\Psi}_\varepsilon \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon} - \int \hat{\Psi}_0 \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon} \\ &\quad + \int \hat{\Psi}_0 \cdot (\hat{\Psi}_\varepsilon \circ F_\varepsilon^n) d\mu_{F_\varepsilon} - \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_\varepsilon^n) d\mu_{F_\varepsilon} \\ &\quad + \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_\varepsilon^n) d\mu_{F_\varepsilon} - \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_0^n) d\mu_{F_0} \\ &\quad + \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_0^n) d\mu_{F_0} - \int \hat{\Psi}_0 \cdot (\hat{\Psi}_0 \circ F_0^n) d\mu_{F_0} \\ &= (I) + (II) + (III) + (IV). \end{aligned}$$

We have $|(II)| \leq \|\hat{\Psi}_0\|_\infty \int |\hat{\Psi}_\varepsilon - \hat{\Psi}_0| \circ F_\varepsilon^n d\mu_{F_\varepsilon} = \|\hat{\Psi}_0\|_\infty \int |\hat{\Psi}_\varepsilon - \hat{\Psi}_0| d\mu_{F_\varepsilon} \rightarrow 0$ by (7), and similarly $(I) \rightarrow 0$. Also, $(IV) \rightarrow 0$ by Proposition 5.1.

Finally, $|(III)| \leq \|\hat{\Psi}_0\|_\infty \int |\hat{\Psi}_0 \circ F_0^n - \hat{\Psi}_0 \circ F_\varepsilon^n| d\mu_{F_\varepsilon}$. Note that

$$\begin{aligned} \int |\hat{\Psi}_0 \circ F_0^n - \hat{\Psi}_0 \circ F_\varepsilon^n| d\mu_{F_\varepsilon} &\leq \sum_{i=1}^n \int |\hat{\Psi}_0 \circ F_0^i \circ F_\varepsilon^{n-i} - \hat{\Psi}_0 \circ F_0^{i-1} \circ F_\varepsilon^{n-i+1}| d\mu_{F_\varepsilon} \\ &= \sum_{i=1}^n \int |\hat{\Psi}_0 \circ F_0^i - \hat{\Psi}_0 \circ F_0^{i-1} \circ F_\varepsilon| d\mu_{F_\varepsilon}. \end{aligned}$$

Thus, by Lemma 5.7, $(III) \rightarrow 0$ as $\varepsilon \rightarrow 0$. \square

We end this section with the following result which gives explicit estimates in terms of Ψ as required for the proof of Theorem 1.1(b).

Proposition 5.8. *Let $\|\Psi\|_{1,\varepsilon} = \int |\Psi| d\mu_{F_\varepsilon}$. For all $H > 0$ and $\alpha > 0$, there exists $C > 0$ such that*

$$|\sigma_{F_\varepsilon}^2(\Psi) - \sigma_{F_\varepsilon}^2(\Psi')| \leq C \|\Psi - \Psi'\|_{1,\varepsilon} (1 + |\log \|\Psi - \Psi'\|_{1,\varepsilon}|)$$

for all Ψ, Ψ' piecewise Hölder with $\|\Psi\|_\alpha, \|\Psi'\|_\alpha \leq H$ and all $\varepsilon \geq 0$.

Proof. Let $\hat{\Psi} = \Psi - \int \Psi d\mu_{F_\varepsilon}$ and $\hat{\Psi}' = \Psi' - \int \Psi' d\mu_{F_\varepsilon}$. Let $N \geq 1$. It follows from the Green-Kubo formula (5) that

$$\begin{aligned} |\sigma_{F_\varepsilon}^2(\Psi) - \sigma_{F_\varepsilon}^2(\Psi')| &\leq 2 \sum_{n=0}^{N-1} \|\hat{\Psi}(\hat{\Psi} \circ F_\varepsilon^n) - \hat{\Psi}'(\hat{\Psi}' \circ F_\varepsilon^n)\|_{1,\varepsilon} \\ &+ 2 \sum_{n=N}^{\infty} \|\hat{\Psi}(\hat{\Psi} \circ F_\varepsilon^n)\|_{1,\varepsilon} + 2 \sum_{n=N}^{\infty} \|\hat{\Psi}'(\hat{\Psi}' \circ F_\varepsilon^n)\|_{1,\varepsilon}. \end{aligned}$$

Now

$$\begin{aligned} \|\hat{\Psi}(\hat{\Psi} \circ F_\varepsilon^n) - \hat{\Psi}'(\hat{\Psi}' \circ F_\varepsilon^n)\|_{1,\varepsilon} &\leq \|\hat{\Psi} - \hat{\Psi}'\|_{1,\varepsilon} \|\hat{\Psi} \circ F_\varepsilon^n\|_\infty \\ &\quad + \|\hat{\Psi}'\|_\infty \|\hat{\Psi} \circ F_\varepsilon^n - \hat{\Psi}' \circ F_\varepsilon^n\|_{1,\varepsilon} \\ &= (\|\hat{\Psi}\|_\infty + \|\hat{\Psi}'\|_\infty) \|\hat{\Psi} - \hat{\Psi}'\|_{1,\varepsilon} \leq 4(\|\Psi\|_\infty + \|\Psi'\|_\infty) \|\Psi - \Psi'\|_{1,\varepsilon}. \end{aligned}$$

Also, by Corollary 5.5, $\|\hat{\Psi}(\hat{\Psi} \circ F_\varepsilon^n)\|_{1,\varepsilon} \leq C\theta^n \|\hat{\Psi}\|_\alpha^2 \leq 4C\theta^n \|\Psi\|_\alpha^2$ and similarly for Ψ' . Hence

$$|\sigma_{F_\varepsilon}^2(\Psi) - \sigma_{F_\varepsilon}^2(\Psi')| \leq 8N(\|\Psi\|_\infty + \|\Psi'\|_\infty) \|\Psi - \Psi'\|_{1,\varepsilon} + C'\theta^N (\|\Psi\|_\alpha^2 + \|\Psi'\|_\alpha^2),$$

where $C' = 8C(1 - \theta)^{-1}$. Taking $N = \lceil q \log \|\Psi - \Psi'\|_{1,\varepsilon} \rceil$ for q sufficiently large yields the desired result. \square

6. VARIANCE CONTINUITY FOR THE FLOWS

By [5, 16], the CLT holds for Hölder observables for the Lorenz flows X_ε . In this section, we show how to obtain continuity of the variances, proving Theorem 1.1. During most of this section we continue to work with normalised families as in section 3, culminating in Corollary 6.5 which is an analogue of Theorem 1.1 for normalised families. We conclude by using Corollary 6.5 to prove Theorem 1.1. Throughout we fix $\beta \in (0, 1)$.

First, we recall some results from the literature. Let $\psi : \mathbb{R}^3 \rightarrow \mathbb{R}$ be C^β . Define

$$\tilde{\psi}(x) = \psi(x) - \psi(0).$$

Also, define the induced observables $\Psi_\varepsilon : \Sigma \setminus \Gamma \rightarrow \mathbb{R}$, $\varepsilon \geq 0$, by

$$\Psi_\varepsilon(\xi) = \int_0^{\tau_\varepsilon(\xi)} \tilde{\psi}(X_\varepsilon(\xi, t)) dt. \quad (8)$$

The left-hand side of (2) is identical for ψ and $\tilde{\psi}$, so

$$\sigma_{X_\varepsilon}^2(\psi) = \sigma_{X_\varepsilon}^2(\tilde{\psi}). \quad (9)$$

Proposition 6.1. *The variances for $\tilde{\psi}$ and Ψ_ε are related by*

$$\sigma_{X_\varepsilon}^2(\tilde{\psi}) = \frac{\sigma_{F_\varepsilon}^2(\Psi_\varepsilon)}{\int \tau_\varepsilon d\mu_{F_\varepsilon}}.$$

Proof. By [16], $\tilde{\psi}$ satisfies the CLT. The method of proof in [16] is to show that Ψ_ε and τ_ε satisfy the CLT, whereupon the CLT for $\tilde{\psi}$ follows from [22, Theorem 1.1]. The desired relation between the variances is given explicitly in [22, Theorem 1.1]. \square

By [2, Lemma 4.1],

$$\lim_{\varepsilon \rightarrow 0} \int \tau_\varepsilon d\mu_{F_\varepsilon} = \int \tau_0 d\mu_{F_0}. \quad (10)$$

Hence the main quantity to control is $\sigma_{F_\varepsilon}^2(\Psi_\varepsilon)$. First, we verify condition (7) in Theorem 5.6.

Lemma 6.2. $\lim_{\varepsilon \rightarrow 0} \int |\Psi_\varepsilon - \Psi_0| d\mu_{F_\varepsilon} = 0$.

Proof. Fix $N \geq 1$ and set $\tau_{\varepsilon,N}(\xi) = \min\{\tau_\varepsilon(\xi), N\}$. Define

$$\Psi_{\varepsilon,N}(\xi) = \int_0^{\tau_{\varepsilon,N}(\xi)} \tilde{\psi}(X_\varepsilon(\xi, t)) dt.$$

Then

$$|\Psi_0 - \Psi_\varepsilon| \leq |\Psi_0 - \Psi_{0,N}| + |\Psi_{0,N} - \Psi_{\varepsilon,N}| + |\Psi_{\varepsilon,N} - \Psi_\varepsilon|.$$

Now,

$$|\Psi_\varepsilon(\xi) - \Psi_{\varepsilon,N}(\xi)| = \left| \int_{\tau_{\varepsilon,N}(\xi)}^{\tau_\varepsilon(\xi)} \tilde{\psi}(X_\varepsilon(\xi, t)) dt \right| \leq 2\|\psi\|_\infty (\tau_\varepsilon(\xi) - \tau_{\varepsilon,N}(\xi)).$$

Recall that $\xi = (x, y, 1)$. Now $\tau_\varepsilon(\xi) - \tau_{\varepsilon,N}(\xi) \leq -C \log|x| - N$, and $\tau_\varepsilon - \tau_{\varepsilon,N}$ is supported on $B_\varepsilon = \{\xi : \tau_\varepsilon(\xi) > N\} \subset \{\xi : |x| < e^{-N/C}\}$. Hence,

$$\begin{aligned} \int |\Psi_\varepsilon - \Psi_{\varepsilon,N}| d\mu_{F_\varepsilon} &\leq 2\|h_\varepsilon\|_\infty \|\psi\|_\infty \int_{B_\varepsilon} (-C \log|x| - N) dx \\ &= 4\|h_\varepsilon\|_\infty \|\psi\|_\infty \int_0^{e^{-N/C}} (-C \log x - N) dx = 4C\|h_\varepsilon\|_\infty \|\psi\|_\infty e^{-N/C}. \end{aligned}$$

By Proposition 4.1(a), there exists $C' > 0$ such that $\int |\Psi_\varepsilon - \Psi_{\varepsilon,N}| d\mu_{F_\varepsilon} \leq C'e^{-N/C} \|\psi\|_\infty$. Similarly, $\int |\Psi_0 - \Psi_{0,N}| d\mu_{F_\varepsilon} \leq C'e^{-N/C} \|\psi\|_\infty$.

Next, $|\Psi_{\varepsilon,N}(\xi) - \Psi_{0,N}(\xi)| \leq (I) + (II)$ where

$$\begin{aligned} (I) &= \int_{\tau_{0,N}(\xi)}^{\tau_{\varepsilon,N}(\xi)} |\tilde{\psi}(X_\varepsilon(\xi, t))| dt \leq 2\|\psi\|_\infty |\tau_{\varepsilon,N}(\xi) - \tau_{0,N}(\xi)|, \\ (II) &= \int_0^{\tau_{0,N}(\xi)} |\tilde{\psi}(X_\varepsilon(\xi, t)) - \tilde{\psi}(X_0(\xi, t))| dt \\ &\leq \int_0^N |\psi(X_\varepsilon(\xi, t)) - \psi(X_0(\xi, t))| dt \leq H_\beta(\psi) \int_0^N |X_\varepsilon(\xi, t) - X_0(\xi, t)|^\beta dt. \end{aligned}$$

For $\delta > 0$ fixed sufficiently small, there exists $\varepsilon_0 > 0$ such that $\tau_{\varepsilon,N}(\xi) \equiv N$ for $|x| < \delta$ and $\varepsilon < \varepsilon_0$. Also, it follows from smoothness of the flow and boundedness of first hit times for $|x| \geq \delta$ that $\tau_{\varepsilon,N} \rightarrow \tau_{0,N}$ uniformly on $\{|x| \geq \delta\}$. Hence $\lim_{\varepsilon \rightarrow 0} \|\tau_{\varepsilon,N} - \tau_{0,N}\|_\infty = 0$ and so $\lim_{\varepsilon \rightarrow 0} \|(I)\|_\infty = 0$.

By continuity of the flow in initial conditions and parameters, $X_\varepsilon(\xi, t) \rightarrow X_0(\xi, t)$ uniformly in $\xi \in \Sigma$ and $t \in [0, N]$. Hence $\lim_{\varepsilon \rightarrow 0} \|(II)\|_\infty = 0$.

We have shown that $\lim_{\varepsilon \rightarrow 0} \int |\Psi_\varepsilon - \Psi_0| d\mu_{F_\varepsilon} \leq 2C'e^{-N/C}$. The result follows since N is arbitrary. \square

Next, we show that $\sup_\varepsilon \|\Psi_\varepsilon\|_\alpha < \infty$ for some $\alpha > 0$.

Lemma 6.3. *Define $\tilde{\Psi}_\varepsilon : \Sigma \rightarrow \mathbb{R}$ by setting*

$$\tilde{\Psi}_\varepsilon(x, y) = \int_0^{-\frac{1}{\lambda_{1,\varepsilon}} \log |x|} \tilde{\psi}(xe^{\lambda_{1,\varepsilon}t}, ye^{\lambda_{2,\varepsilon}t}, e^{\lambda_{3,\varepsilon}t}) dt.$$

Choose $0 < \alpha' < -\lambda_{3,\varepsilon}\beta/(\lambda_{1,\varepsilon} - \lambda_{3,\varepsilon})$. Then $\tilde{\Psi}_\varepsilon$ is piecewise $C^{\alpha'}$. Moreover, there is a constant $C > 0$ such that $\|\tilde{\Psi}_\varepsilon\|_{\alpha'} \leq C\|\psi\|_\beta$ for all $\varepsilon \geq 0$.

Proof. We suppress the dependence on ε . Write $\tilde{\psi}(x, y, z) = \psi_1(x) + \psi_2(x, y, z)$ where $\psi_1(x) = \tilde{\psi}(x, 0, 0)$. Then ψ_1 and ψ_2 are Hölder with $H_\beta(\psi_1) \leq H_\beta(\psi)$ and $H_\beta(\psi_2) \leq 2H_\beta(\psi)$. Also, $\psi_1(0) = 0$ and $\psi_2(x, 0, 0) \equiv 0$. Define

$$\tilde{\Psi}_1(x) = \int_0^{-\frac{1}{\lambda_1} \log x} \psi_1(xe^{\lambda_1 t}) dt, \quad \tilde{\Psi}_2(x, y) = \int_0^{-\frac{1}{\lambda_1} \log x} \psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, e^{\lambda_3 t}) dt.$$

Recall that $\lambda_2 < \lambda_3 < 0 < \lambda_1$.

First we carry out the estimates for $\tilde{\Psi}_1$ with $\alpha' = \beta$. By the change of variables $u = xe^{\lambda_1 t}$,

$$\tilde{\Psi}_1(x) = \frac{1}{\lambda_1} \int_x^1 \frac{\psi_1(u)}{u} du.$$

Now $|\psi_1(u)| = |\psi_1(u) - \psi_1(0)| \leq H_\beta(\psi)u^\beta$, so $|\tilde{\Psi}_1|_\infty \leq \frac{H_\beta(\psi)}{\lambda_1} \int_0^1 u^{-(1-\beta)} du = \frac{H_\beta(\psi)}{\lambda_1\beta}$. Also for $x_1 > x_2 > 0$,

$$\begin{aligned} |\tilde{\Psi}_1(x_1) - \tilde{\Psi}_1(x_2)| &\leq \frac{H_\beta(\psi)}{\lambda_1} \int_{x_2}^{x_1} u^{-(1-\beta)} du \\ &= \frac{H_\beta(\psi)}{\lambda_1\beta} (x_1^\beta - x_2^\beta) \leq \frac{H_\beta(\psi)}{\lambda_1\beta} (x_1 - x_2)^\beta. \end{aligned}$$

Here, we have used that $x_1^\beta - x_2^\beta \leq (x_1 - x_2)^\beta$ for all $\beta \in [0, 1]$. Hence $\|\tilde{\Psi}_1\|_\beta \leq \frac{1}{\lambda_1\beta} \|\psi\|_\beta$.

Next, we carry out the estimates for $\tilde{\Psi}_2$. Note that

$$\begin{aligned} |\psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, e^{\lambda_3 t})| &= |\psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, e^{\lambda_3 t}) - \psi_2(xe^{\lambda_1 t}, 0, 0)| \\ &= |\psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, e^{\lambda_3 t}) - \psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, 0)| \\ &\quad + |\psi_2(xe^{\lambda_1 t}, ye^{\lambda_2 t}, 0) - \psi_2(xe^{\lambda_1 t}, 0, 0)| \\ &\leq H_\beta(\psi_2)(e^{\beta\lambda_3 t} + |y|^\beta e^{\beta\lambda_2 t}) \leq 4H_\beta(\psi)e^{\beta\lambda_3 t}. \end{aligned} \quad (11)$$

In particular,

$$|\tilde{\Psi}_2|_\infty \leq 4H_\beta(\psi) \int_0^{-\frac{1}{\lambda_1} \log x} e^{\beta\lambda_3 t} dt = \frac{4H_\beta(\psi)}{\beta|\lambda_3|} (1 - x^{-\beta\lambda_3/\lambda_1}) \leq \frac{4H_\beta(\psi)}{\beta|\lambda_3|}.$$

Similarly,

$$|\tilde{\Psi}_2(x, y_1) - \tilde{\Psi}_2(x, y_2)| \leq \int_0^{-\frac{1}{\lambda_1} \log x} H_\beta(\psi_2) |y_1 - y_2|^\beta e^{\beta\lambda_2 t} dt \leq \frac{2H_\beta(\psi)}{\beta|\lambda_2|} |y_1 - y_2|^\beta.$$

For $x_1 > x_2 > 0$, we have $|\tilde{\Psi}_2(x_1, y) - \tilde{\Psi}_2(x_2, y)| \leq A + B$ where

$$A = \int_{-\frac{1}{\lambda_1} \log x_1}^{-\frac{1}{\lambda_1} \log x_2} \psi_2(x_2 e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t}) dt,$$

$$B = \int_0^{-\frac{1}{\lambda_1} \log x_1} |\psi_2(x_1 e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t}) - \psi_2(x_2 e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t})| dt.$$

By (11),

$$A \leq 4H_\beta(\psi) \int_{-\frac{1}{\lambda_1} \log x_1}^{-\frac{1}{\lambda_1} \log x_2} e^{\beta \lambda_3 t} dt = \frac{4H_\beta(\psi)}{\beta |\lambda_3|} (x_1^\gamma - x_2^\gamma) \leq \frac{4H_\beta(\psi)}{\beta |\lambda_3|} (x_1 - x_2)^\gamma$$

where $\gamma = -\lambda_3 \beta / \lambda_1$. Let

$$\alpha' = \delta \beta, \quad \lambda' = \delta \lambda_1 + (1 - \delta) \lambda_3,$$

where $0 < \delta < -\lambda_3 / (\lambda_1 - \lambda_3) < 1$. In particular, $\alpha' < \gamma$ and $\lambda' < 0$. Since the eigenvalues λ_1, λ_3 depend continuously on ε , we can choose δ independent of ε . The inequality $\min\{a, b\} \leq a^\delta b^{1-\delta}$ holds for all $a, b \geq 0$. By (11),

$$|\psi_2(x_1 e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t}) - \psi_2(x_2 e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t})|$$

$$\leq \min\{H_\beta(\psi_2)(x_1 - x_2)^\beta e^{\beta \lambda_1 t}, 8H_\beta(\psi) e^{\beta \lambda_3 t}\} \leq 8H_\beta(\psi)(x_1 - x_2)^{\delta \beta} e^{\delta \lambda' t}.$$

Hence $B \leq \frac{8}{\delta |\lambda'|} H_\beta(\psi)(x_1 - x_2)^{\alpha'}$ completing the proof. \square

Corollary 6.4. *There exists $\alpha \in (0, \beta)$ such that Ψ_ε is piecewise C^α . Moreover, there is a constant $C > 0$ such that $\sup_\varepsilon \|\Psi_\varepsilon\|_\alpha \leq C \|\psi\|_\beta$.*

Proof. After a C^α change of coordinates in a neighbourhood of the singularity, we may suppose without loss of generality that the flow X_ε is linear near the singularity. Here, $\alpha > 0$ can be chosen independent of ε with bounded Hölder constants for the linearisation [10]. We choose $\alpha < \alpha'$ where α' is as in Lemma 6.3

For the remainder of the proof we suppress the dependence on ε . The first hit time in this neighbourhood is given by $\tau_1(\xi) = -\frac{1}{\lambda_1} \log |x|$. Recall that $\xi = (x, y, 1)$ and set $\xi' = X(\xi, \tau_1(\xi)) = (1, yx^{-\lambda_2/\lambda_1}, x^{-\lambda_3/\lambda_1})$. Note that the dependence of ξ' on ξ is Hölder (also uniformly in ε). Then $\tau = \tau_1 + \tau_2$ where τ_2 is C^α in ξ' and hence ξ (uniformly in ε). Also, $\Psi = \tilde{\Psi} + \hat{\Psi}$ where

$$\tilde{\Psi}(\xi) = \int_0^{\tau_1(\xi)} \tilde{\psi}(x e^{\lambda_1 t}, y e^{\lambda_2 t}, e^{\lambda_3 t}) dt, \quad \hat{\Psi}(\xi) = \int_0^{\tau_2(\xi)} \tilde{\psi}(X(\xi', t)) dt. \quad (12)$$

Since $\|\tau_2\|_\infty$ is bounded, it is immediate that $\hat{\Psi}$ is C^α uniformly in ε . By Lemma 6.3, the same is true for $\tilde{\Psi}$. \square

We can now state and prove the analogue of Theorem 1.1 for normalised families. Define $\|\psi\|_{(\varepsilon)} = \int |\psi| d\mu_\varepsilon + |\psi(0)|$.

Corollary 6.5. *Let X_ε be a normalised family of flows admitting geometric Lorenz attractors. Then*

$$(a) \lim_{\varepsilon \rightarrow 0} \sigma_{X_\varepsilon}^2(\psi) = \sigma_{X_0}^2(\psi) \text{ for all } C^\beta \text{ observables } \psi : \mathbb{R}^3 \rightarrow \mathbb{R}$$

(b) For any $H > 0$, there exists $C > 0$ such that

$$|\sigma_{X_\varepsilon}^2(\psi) - \sigma_{X_\varepsilon}^2(\psi')| \leq C \|\psi - \psi'\|_{(\varepsilon)} (1 + |\log \|\psi - \psi'\|_{(\varepsilon)}|)$$

for all C^β observables $\psi, \psi' : \mathbb{R}^3 \rightarrow \mathbb{R}$ with $\|\psi\|_\beta, \|\psi'\|_\beta \leq H$, and all $\varepsilon \geq 0$.

Proof. By Corollary 6.4, there exists $C_1 > 0$, $\alpha \in (0, 1)$ such that $\Psi_\varepsilon, \Psi'_\varepsilon$ are piecewise C^α and

$$\sup_\varepsilon \|\Psi_\varepsilon\|_\alpha \leq C_1 \|\psi\|_\beta, \quad \sup_\varepsilon \|\Psi'_\varepsilon\|_\alpha \leq C_1 \|\psi'\|_\beta. \quad (13)$$

By (9) and Proposition 6.1,

$$\begin{aligned} |\sigma_{X_\varepsilon}^2(\psi) - \sigma_{X_{\varepsilon'}}^2(\psi')| &= |\sigma_{X_\varepsilon}^2(\tilde{\psi}) - \sigma_{X_{\varepsilon'}}^2(\tilde{\psi}')| = \left| \frac{\sigma_{F_\varepsilon}^2(\Psi_\varepsilon)}{\int \tau_\varepsilon d\mu_{F_\varepsilon}} - \frac{\sigma_{F_{\varepsilon'}}^2(\Psi'_{\varepsilon'})}{\int \tau_{\varepsilon'} d\mu_{F_{\varepsilon'}}} \right| \\ &\leq \sigma_{F_{\varepsilon'}}^2(\Psi'_{\varepsilon'}) \left| \frac{1}{\int \tau_\varepsilon d\mu_{F_\varepsilon}} - \frac{1}{\int \tau_{\varepsilon'} d\mu_{F_{\varepsilon'}}} \right| + \frac{1}{\int \tau_\varepsilon d\mu_{F_\varepsilon}} \left| \sigma_{F_\varepsilon}^2(\Psi_\varepsilon) - \sigma_{F_{\varepsilon'}}^2(\Psi'_{\varepsilon'}) \right|. \end{aligned}$$

First suppose that $\psi = \psi'$ and $\varepsilon' = 0$. By Lemma 6.2 and (13), we can apply Theorem 5.6 to deduce that $\lim_{\varepsilon \rightarrow 0} \sigma_{F_\varepsilon}^2(\Psi_\varepsilon) = \sigma_{F_0}^2(\Psi_0)$. Part (a) now follows from (10).

Next suppose that $\varepsilon = \varepsilon'$. By Proposition 5.8 and (13), there is a constant $C > 0$ (independent of ε) such that $|\sigma_{F_\varepsilon}^2(\Psi_\varepsilon) - \sigma_{F_\varepsilon}^2(\Psi'_\varepsilon)| \leq C \|\Psi_\varepsilon - \Psi'_\varepsilon\|_{1,\varepsilon} (1 + |\log \|\Psi_\varepsilon - \Psi'_\varepsilon\|_{1,\varepsilon}|)$ where

$$\begin{aligned} \|\Psi_\varepsilon - \Psi'_\varepsilon\|_{1,\varepsilon} &= \int |\Psi_\varepsilon - \Psi'_\varepsilon| d\mu_{F_\varepsilon} \leq \int \int_0^{\tau_\varepsilon(\xi)} |\tilde{\psi}(X_\varepsilon(\xi, t) - \tilde{\psi}'(X_\varepsilon(\xi, t))| dt d\mu_{F_\varepsilon} \\ &= \int \tau_\varepsilon d\mu_{F_\varepsilon} \int |\tilde{\psi} - \tilde{\psi}'| d\mu_\varepsilon \leq \int \tau_\varepsilon d\mu_{F_\varepsilon} \left(\int |\psi - \psi'| d\mu_\varepsilon + |\psi(0) - \psi'(0)| \right). \end{aligned}$$

This proves part (b). \square

Proof of Theorem 1.1. Let $\phi_\varepsilon : U \rightarrow U$ be the family of normalising conjugacies in section 3. Recall that $\tilde{X}_\varepsilon(x, t) = \phi_\varepsilon^{-1} \circ X_\varepsilon(\phi_\varepsilon(x), t)$. Define

$$\psi_\varepsilon = \psi \circ \phi_\varepsilon, \quad \psi'_\varepsilon = \psi' \circ \phi_\varepsilon, \quad \tilde{\mu}_\varepsilon = \phi_\varepsilon^{-1} * \mu_\varepsilon.$$

Then ψ_ε is Hölder and

$$\int \psi_\varepsilon d\tilde{\mu}_\varepsilon = \int \psi d\mu_\varepsilon, \quad \int_0^t \psi_\varepsilon \circ \tilde{X}_\varepsilon(s) ds = \left(\int_0^t \psi \circ X_\varepsilon(s) ds \right) \circ \phi_\varepsilon.$$

Hence $\int_0^t \psi \circ X_\varepsilon(s) ds - t \int \psi d\mu_\varepsilon$ and $\int_0^t \psi_\varepsilon \circ \tilde{X}_\varepsilon(s) ds - t \int \psi_\varepsilon d\tilde{\mu}_\varepsilon$ have the same distribution (relative to the probability measures μ_ε and $\tilde{\mu}_\varepsilon$ respectively) so it follows from (2) that $\sigma_{X_\varepsilon}^2(\psi) = \sigma_{\tilde{X}_\varepsilon}^2(\psi_\varepsilon)$. Similar comments apply to ψ' .

It follows from the definitions that \tilde{X}_ε is a normalised family. By Corollary 3.2, there exists $H > 0$ and $\beta \in (0, 1)$ such that $\sup_\varepsilon \|\psi_\varepsilon\|_\beta \leq H$ and $\sup_\varepsilon \|\psi'_\varepsilon\|_\beta \leq H$.

By Corollary 6.5(b), there is a constant $C > 0$ such that

$$\begin{aligned} |\sigma_{X_\varepsilon}^2(\psi) - \sigma_{X_0}^2(\psi)| &= |\sigma_{\tilde{X}_\varepsilon}^2(\psi_\varepsilon) - \sigma_{\tilde{X}_0}^2(\psi_0)| \\ &\leq |\sigma_{\tilde{X}_\varepsilon}^2(\psi_\varepsilon) - \sigma_{\tilde{X}_\varepsilon}^2(\psi_0)| + |\sigma_{\tilde{X}_\varepsilon}^2(\psi_0) - \sigma_{\tilde{X}_0}^2(\psi_0)| \\ &\leq C \|\psi_\varepsilon - \psi_0\|_\infty (1 + |\log \|\psi_\varepsilon - \psi_0\|_\infty|) + |\sigma_{\tilde{X}_\varepsilon}^2(\psi_0) - \sigma_{\tilde{X}_0}^2(\psi_0)|. \end{aligned}$$

By Corollary 6.5(a), the second term on the right-hand side converges to zero as $\varepsilon \rightarrow 0$. Also, $\lim_{\varepsilon \rightarrow 0} \|\psi_\varepsilon - \psi_0\|_\infty = \lim_{\varepsilon \rightarrow 0} \|\psi \circ \phi_\varepsilon^{-1} - \psi \circ \phi_0^{-1}\|_\infty = 0$ by Corollary 3.2. Hence $\lim_{\varepsilon \rightarrow 0} \sigma_{X_\varepsilon}^2(\psi) = \sigma_{X_0}^2(\psi)$ proving part (a).

Finally, by Corollary 6.5(b),

$$\begin{aligned} |\sigma_{X_0}^2(\psi) - \sigma_{X_0}^2(\psi')| &= |\sigma_{X_0}^2(\psi_0) - \sigma_{X_0}^2(\psi'_0)| \\ &\leq C \|\psi_0 - \psi'_0\|_{(0)} (1 + |\log \|\psi_0 - \psi'_0\|_{(0)}|) \\ &= C \|\psi - \psi'\| (1 + |\log \|\psi - \psi'\||) \end{aligned}$$

yielding part (b). \square

REFERENCES

- [1] V.S. Afraimovich, V.V. Bykov, L.P. Sil'nikov. The origin and structure of the Lorenz attractor. *Dokl. Akad. Nauk SSSR* 234 (1977), no. 2, 336–339.
- [2] J. F. Alves, M. Soufi. Statistical stability of geometric Lorenz attractors. *Fund. Math.* 224 (2014), no. 3, 219–231.
- [3] V. Araújo, S. Galatolo, M. J. Pacifico. Decay of correlations for maps with uniformly contracting fibers and logarithm law for singular hyperbolic attractors. *Math. Z.* 276 (2014), no. 3-4, 1001–1048.
- [4] V. Araújo, I. Melbourne. Existence and smoothness of the stable foliation for sectional hyperbolic attractors. *Bull. Lond. Math. Soc.* 49 (2017), no. 2, 351–367.
- [5] V. Araújo, I. Melbourne. Mixing properties and statistical limit theorems for singular hyperbolic flows without a smooth stable foliation. *Adv. Math.* 349 (2019), 212–245.
- [6] V. Araújo, I. Melbourne, P. Varandas. Rapid mixing for the Lorenz attractor and statistical limit laws for their time-1 maps. *Comm. Math. Phys.* 340 (2015), 901–938.
- [7] V. Araújo, M. J. Pacifico. *Three-dimensional flows*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics. Results in Mathematics and Related Areas. 3rd Series, vol. 53. Springer, Berlin, Heidelberg (2010).
- [8] V. Araújo, M. J. Pacifico, E. R. Pujals, M. Viana. Singular-hyperbolic attractors are chaotic. *Trans. Amer. Math. Soc.* 361 (5) (2009), 2431–2485.
- [9] W. Bahsoun, M. Ruziboev. On the statistical stability of Lorenz attractors with $C^{1+\alpha}$ stable foliation. *Ergodic Theory Dynam. Systems* 39 (2019), no. 12, 3169–3184.
- [10] L. Barreira and C. Valls. Hölder Grobman-Hartman linearization. *Discrete Contin. Dyn. Syst.* 18 (2007), 187–197.
- [11] T. Bomfim, A. Castro, P. Varandas. Differentiability of thermodynamical quantities in non-uniformly expanding dynamics. *Adv. Math.* 292 (2016), 478–528.
- [12] R. T. Bortolotti. Physical measures for certain partially hyperbolic attractors on 3-manifolds. *Ergodic Theory Dynam. Systems* 39 (2019), 74–104.
- [13] M.F. Demers, H.-K. Zhang, A functional analytic approach to perturbations of the Lorenz gas. *Commun. Math. Phys.* 324 (2013), 767–830.
- [14] S. Galatolo, R. Lucena. Spectral Gap and quantitative statistical stability for systems with contracting fibres and Lorenz like maps. *Discrete Contin. Dyn. Syst.* 40 (2020), no. 3, 1309–1360.
- [15] J. Guckenheimer, R.F. Williams. Structural stability of Lorenz attractors. *Inst. Hautes Études Sci. Publ. Math.* 50 (1979), 59–72.
- [16] M. Holland, I. Melbourne. Central limit theorems and invariance principles for Lorenz attractors. *J. Lond. Math. Soc.* (2) 76 (2007), no. 2, 345–364.
- [17] G. Keller. Generalized bounded variation and applications to piecewise monotonic transformations. *Z. Wahrsch. Verw. Gebiete* 69 (1985), no. 3, 461–478.
- [18] G. Keller, C. Liverani. Stability of the spectrum for transfer operators. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) 28 (1999), no. 1, 141–152.
- [19] A. Korepanov, Z. Kosloff, I. Melbourne. Martingalecoboundary decomposition for families of dynamical systems. *Ann. I. H. Poincaré Anal. Non Linéaire* 35 (2018), 859–885.

- [20] E. D. Lorenz. Deterministic nonperiodic flow. *J. Atmosph. Sci.* 20 (1963), 130–141.
- [21] I. Melbourne, M. Nicol. Almost sure invariance principle for nonuniformly hyperbolic systems. *Comm. Math. Phys.* 260 (2005), no. 1, 131–146.
- [22] I. Melbourne, A. Török. Statistical limit theorems for suspension flows. *Israel J. Math.* 144 (2004), 191–209.
- [23] C. A. Morales, M. J. Pacifico, E. R. Pujals. Robust transitive singular sets for 3-flows are partially hyperbolic attractors or repellers. *Ann. of Math. (2)* 160 (2004), no. 2, 375–432.
- [24] W. Tucker. The Lorenz attractor exists. *C. R. Acad. Sci. Paris Sér. I Math.*, 328, (1999), 1197–1202.
- [25] W. Tucker. A rigorous ODE solver and Smale’s 14th problem. *Found. Comput. Math.* 2 (2002), 53–117.